

# ANALYTIC ASYMPTOTIC EXPANSIONS OF THE RESHETIKHIN–TURAEV INVARIANTS OF SEIFERT 3–MANIFOLDS FOR SU(2)

SØREN KOLD HANSEN

**ABSTRACT.** We calculate the large quantum level asymptotic expansion of the RT–invariants associated to SU(2) of all oriented Seifert 3–manifolds  $X$  with orientable base or non-orientable base with even genus. Moreover, we identify the Chern–Simons invariants of flat SU(2)–connections on  $X$  in the asymptotic formula thereby proving the so-called asymptotic expansion conjecture (AEC) due to J. E. Andersen [An1], [An2] for these manifolds. For the case of Seifert manifolds with base  $S^2$  we actually prove a little weaker result, namely that the asymptotic formula has a form as predicted by the AEC but contains some extra terms which should be zero according to the AEC. We prove that these ‘extra’ terms are indeed zero if the number of exceptional fibers  $n$  is less than 4 and conjecture that this is also the case if  $n \geq 4$ . For the case of Seifert fibered rational homology spheres we identify the Casson–Walker invariant in the asymptotic formula.

Our calculations demonstrate a general method for calculating the large  $r$  asymptotics of a finite sum  $\sum_{k=1}^r f(k)$ , where  $f$  is a meromorphic function depending on the integer parameter  $r$  and satisfying certain symmetries. Basically the method, which is due to Rozansky [Ro1], [Ro3], is based on a limiting version of the Poisson summation formula together with an application of the steepest descent method from asymptotic analysis.

## CONTENTS

|                                                                                                |     |
|------------------------------------------------------------------------------------------------|-----|
| 1. Introduction                                                                                | 2   |
| 2. Asymptotic expansions og quantum type                                                       | 14  |
| 3. The Reshetikhin–Turaev invariants of Seifert manifolds for SU(2)                            | 19  |
| 4. The analytic asymptotic expansions of the Reshetikin–Turaev invariants of Seifert manifolds | 21  |
| 4.1. The moduli space of flat SU(2)–connections on Seifert manifolds                           | 30  |
| 4.2. The genus 0 case                                                                          | 35  |
| 4.3. Small Seifert manifolds with base $\mathbb{RP}^2$                                         | 39  |
| 4.4. The contribution from the trivial connection and the Casson–Walker invariant              | 39  |
| 5. Proof of Theorem 4.1                                                                        | 44  |
| 5.1. Calculation of asymptotics. A general case                                                | 48  |
| 6. Proof of Theorem 4.4                                                                        | 74  |
| 7. Appendices                                                                                  | 83  |
| 7.1. Appendix A. The periodicity lemma                                                         | 83  |
| 7.2. Appendix B. Proofs of Lemma 5.4–Lemma 5.6                                                 | 84  |
| 7.3. Appendix C. Proofs of Lemma 5.8 and Proposition 5.9                                       | 89  |
| 7.4. Appendix D. Proofs of Lemma 5.11, Lemma 5.13 and Proposition 5.14                         | 91  |
| 7.5. Appendix E. Proofs of Lemma 5.16 and Proposition 5.17                                     | 96  |
| 7.6. Appendix F. Proof of Lemma 6.1                                                            | 104 |
| References                                                                                     | 105 |

## 1. INTRODUCTION

In this paper we investigate the large quantum level asymptotics of the Reshetikhin–Turaev invariants of the Seifert manifolds. Here and elsewhere a 3–manifold is closed and oriented unless otherwise stated. In particular, a Seifert manifold is an oriented Seifert manifold.

It is now a well established fact that the Reshetikhin–Turaev approach [RT], [Tu] leads to a family of  $2 + 1$ –dimensional topological quantum field theories (TQFTs) indexed by a simply connected compact simple Lie group  $G$  and an integer  $r$  (the shifted quantum level) bigger than or equal to the dual Coxeter number  $h^\vee$  of  $G$ . In particular, we have invariants  $\tau_r^G$  of the 3–manifolds called the quantum  $G$ –invariants or the RT–invariants associated to  $G$ .

There are several other approaches to these quantum invariants and their underlying TQFTs, see e.g. [Tu], [BK] and references therein. For the following discussion we note that J. E. Andersen and K. Ueno [AU1] have recently constructed a family of TQFTs using ideas from conformal field theory, notably the works [TUY], [U], [KNTY]. They have so far proved that these TQFTs for the groups  $SU(n)$  coincide with the TQFTs of Reshetikhin and Turaev associated to these Lie groups, cf. [AU2]. Via work of Laszlo [La] one can combine the work of Andersen and Ueno with work of Axelrod, Della Pietra and Witten [APW], Hitchin [Hi], Faltings [Fa] and others on the geometric quantization of the moduli space of flat connections on surfaces to obtain an alternative gauge theoretic approach to the TQFTs of Andersen and Ueno. We refer to [At] and [An3] and references therein for details on this gauge theoretic approach which was originally outlined by Witten in [Wi].

The main aim of this paper is to prove the following conjecture for Seifert manifolds for  $G = SU(2)$ .

**Conjecture 1.1** (J. E. Andersen [An1], [An2]. The asymptotic expansion conjecture (AEC)).

<sup>1</sup> For any 3–manifold  $X$  there exist constants  $D_j \in \mathbb{Q}$  and  $b_j \in \mathbb{C}$  for  $j = 0, 1, \dots, n$  and  $a_l^j \in \mathbb{C}$  for  $j = 0, 1, \dots, n$  and  $l = 1, 2, \dots$  such that the asymptotic expansion of  $\tau_r^G(X)$  in the limit  $r \rightarrow \infty$  is given by

$$\tau_r^G(X) \sim \sum_{j=0}^n e^{2\pi i r q_j} r^{D_j} b_j \left( 1 + \sum_{l=1}^{\infty} a_l^j r^{-\theta_l} \right), \quad (1)$$

where  $q_0 = 0, q_1, \dots, q_n$  are the finitely many different values of the Chern–Simons functional on the moduli space of flat  $G$ –connections on  $X$ , and  $\{\theta_l\}$  is a strictly increasing sequence in  $\frac{1}{b}\mathbb{Z}_{>0}$ , where  $b$  is the least positive number such that  $bD_j \in \mathbb{Z}$  for  $j = 0, 1, \dots, n$ .

Here  $\sim$  means that there for all non-negative  $N$  is a  $C_N \in \mathbb{R}$  such that

$$\left| \tau_r^G(X) - \sum_{j=0}^n e^{2\pi i r q_j} r^{D_j} b_j \left( 1 + \sum_{l=1}^N a_l^j r^{-\theta_l} \right) \right| \leq C_N r^{D - \theta_{N+1}}$$

for all  $r \in \mathbb{Z}_{\geq h^\vee}$ , where  $D = \max\{D_0, \dots, D_n\}$ .

The formulation above seems a little awkward from the point of view of asymptotic analysis. We shall later on see a more compact formulation of the conjecture, cf. Conjecture 2.5. The main reason for the above formulation is to expose the quantities  $D_j$  and  $b_j$  which are believed to have interpretations in terms of certain topological/geometric quantities, cf. Conjecture 1.3 below and [An2].

Let us give a review of some of the results leading to this conjecture. We want by this review to stress what is known today from a rigorous mathematical point of view and what is based on arguments using techniques from theoretical physics not yet well understood from a

---

<sup>1</sup>We give the conjecture in a slightly different form than in [An2], where  $\theta_l = l$ .

mathematical point of view. In 1988 E. Witten [Wi] proposed new invariants  $Z_k^G(X, L)$  of a 3-manifold  $X$  with an embedded (colored) link  $L$ , which generalize the famous Jones polynomial [Jo] of knots in the 3-sphere. Here  $G$  is a nice Lie group as above and  $k$  is a non-negative integer called the (quantum) level. In case there is no link present,  $Z_k^G(X)$  is given (formally) by the following path integral over the infinite dimensional space of gauge equivalence classes of  $G$ -connections on  $X$

$$Z_k^G(X) = \int e^{2\pi\sqrt{-1}k\text{CS}(A)} \mathfrak{D}A, \quad (2)$$

where

$$\text{CS}(A) = \frac{1}{8\pi^2} \int_X \text{tr} \left( A \wedge dA + \frac{2}{3} A \wedge A \wedge A \right) \quad (3)$$

is the Chern–Simons functional associated to  $G$ . It is well-known that CS considered as a function into  $\mathbb{R}/\mathbb{Z}$  is invariant under gauge transformations if the Ad-invariant inner product  $\text{tr}$  on  $\mathfrak{g}$  (the Lie algebra of  $G$ ) is normalized properly, see e.g. [F, Sect. 2]. Although no rigorous mathematical definition of the path integral (2) has been given yet, see [JL, Sect. 20.2.A] for some comments, Witten was able using path integral techniques to give a recipe for calculating the invariant  $Z_k^G(X)$  via a surgery description of  $X$ . These ideas enabled people to obtain explicit (and rigorous) formulas for the invariants  $Z_k^G(X)$  for some classes of manifolds  $X$  such as lens spaces [FG], [Ga1], [J1] and more generally of Seifert manifolds with orientable base [FG], [Ga1], [Ro1], [Ro3].

In quantum field theory it is natural to study Feynman path integrals as in (2) and a main approach to understand them is via their behaviour in the large  $k$  limit. By using stationary phase approximation techniques together with path integral arguments, Witten was able [Wi] to express the leading large  $k$  asymptotics of  $Z_k^G(X)$  (also called the semiclassical approximation) as a sum over the set  $\mathcal{M}$  of stationary points of the Chern–Simons functional CS, the summand being given by such invariants as Chern–Simons invariants, spectral flows and Reidemeister torsions. Here  $\mathcal{M}$  is the moduli space of gauge equivalence classes of flat connections in the trivial  $G$ -bundle over  $X$  (and  $\mathcal{M}$  is assumed discrete hence finite in Witten’s derivation). Freed and Gompf [FG] and Jeffrey [J1] suggested different refinements to Witten’s semiclassical approximation. In particular Jeffrey allowed  $\mathcal{M}$  to have smooth components of non-zero dimension, and in these cases the sum over  $\mathcal{M}$  should be replaced by some integral over  $\mathcal{M}$ . Let us give some details. Let  $A$  be a flat  $G$ -connection on  $X$  and consider the elliptic complex  $d_A : \Omega^*(X, \mathfrak{g}) \rightarrow \Omega^{*+1}(X, \mathfrak{g})$ , where  $d_A f = df + [A \wedge f]$  is the covariant derivative in the adjoint representation. Let  $h_A^i$  be the dimension of the  $i$ th cohomology group  $H^i(X, d_A)$  of this complex. If  $\mathcal{M}$  is discrete and the covariant derivative complex is acyclic for all  $A$  (like in Witten’s derivation) then the conjectured formula for the semiclassical approximation of  $Z_k^G(X)$  is

$$Z_k^G(X) \sim_{k \rightarrow \infty} \frac{1}{|Z(G)|} e^{-\pi\sqrt{-1} \dim(G)(1+b_1(X))/4} \times \sum_{A \in \mathcal{M}} e^{2\pi\sqrt{-1}r\text{CS}(A)} r^{D_A} e^{-2\pi\sqrt{-1}(I_A/4+B_A)} \sqrt{\tau_X(A)}, \quad (4)$$

where  $r = k + h^\vee$ ,  $Z(G)$  is the center of  $G$  and  $b_1(X)$  is the first betti number of  $X$ . Moreover,  $\tau_X(A) \in \mathbb{R}_+$  is the Reidemeister torsion of the complex  $(\Omega^*(X, \mathfrak{g}), d_A)$  and  $I_A \in \mathbb{Z}/8\mathbb{Z}$  is the spectral flow of the family of operators  $\begin{pmatrix} *d_{A_t} & -d_{A_t}^* \\ d_{A_t}^* & 0 \end{pmatrix}$  on  $\Omega^1(X; \mathfrak{g}) \otimes \Omega^3(X; \mathfrak{g})$ , where  $A_t$  is a path in the affine space of  $G$ -connections on  $X$  from the trivial connection to the flat connection  $A$ . Under the above assumptions  $B_A = D_A = 0$  for all  $A$ , but the factor  $r^{D_A} e^{-2\pi\sqrt{-1}B_A}$  is included for the following discussion. (Here and elsewhere we follow the usual practice by denoting a connection and its gauge equivalence class by the same symbol.)

If we allow some of the  $A$ 's to be reducible (implying that  $h_A^0 > 0$ ) (still assuming that  $\mathcal{M}$  is finite) then it is conjectured that the above formula holds with  $D_A = -h_A^0/2$  and  $B_A = h_A^0/8$ . In particular, these reducible connections do not contribute to the leading large  $k$  asymptotics of  $Z_k^G(X)$  (which is given by the sum of terms with  $D_A = \max_{A \in \mathcal{M}} \{D_A\}$ ). If the gauge equivalence class of a flat connection  $A$  is non-isolated in  $\mathcal{M}$  then the situation becomes more complicated. It is conjectured that we in that case have  $D_A = (h_A^1 - h_A^0)/2$  and  $B_A = (h_A^0 + h_A^1)/8$ , and that the above sum over  $\mathcal{M}$  should be replaced by some integral over  $\mathcal{M}$ . In cases, where  $A$  is a non-acyclic connection, the interpretation of the factor  $\sqrt{\tau_X(A)}$  is not clear,  $A$  being called acyclic if  $(\Omega^*(X, \mathfrak{g}), d_A)$  is acyclic. (One can still define a torsion  $\in \mathbb{R}_+$  of the covariant derivative complex once a basis of  $H^*(X, d_A)$  has been fixed, but there seem to be some normalization problems even in the case of isolated reducible points in  $\mathcal{M}$ .) In the case of non-isolated points in  $\mathcal{M}$  the factor  $\sqrt{\tau_X(A)}$  should somehow be interpreted as a density function giving the needed measure on  $\mathcal{M}$ . We refer to [J1] and [An2] for some comments on this and also to [Ro1], [Ro2] for some further comments on the semiclassical approximation of  $Z_k^G(X)$ . We note that the Chern–Simons functional is constant on the connected components of  $\mathcal{M}$ , so in all cases one gets a (conjectured) formula for the leading large  $k$  asymptotics of  $Z_k^G(X)$  of the form

$$Z_k^G(X) \sim_{k \rightarrow \infty} \sum_{j=1}^m e^{2\pi\sqrt{-1}\text{rCS}_j} \int_{\mathcal{M}_j} r^{(h_A^1 - h_A^0)/2} e^{-2\pi\sqrt{-1}(I_A/4 + (h_A^0 + h_A^1)/8)} f(A), \quad (5)$$

where the sum is over the connected components of  $\mathcal{M}$ ,  $\text{CS}_j$  is the Chern–Simons invariant of the elements in the  $j$ th component  $\mathcal{M}_j$  of  $\mathcal{M}$  and the integral over  $\mathcal{M}_j$  is w.r.t. some density function  $f(A)$  generalizing the factor  $\frac{1}{|Z(G)|} e^{-\pi\sqrt{-1}\dim(G)(1+b_1(X))/4} \sqrt{\tau_X(A)}$  in (4).

A main first observation is that the expression in the right-hand side of (4) is mathematical rigorous in case  $\mathcal{M}$  is finite (although it is not clear what the meaning of the factor  $\sqrt{\tau_X(A)}$  should be in the non-acyclic case). Moreover, as stated above, one can in principal obtain a rigorous expression for the invariants  $Z_k^G(X)$  via a surgery description of  $X$ . Thus a kind of verification of (4) should be possible. This program was first carried out partly by Freed and Gompf [FG] presenting a large amount of computer calculations for the  $\text{SU}(2)$ -invariants of lens spaces and some 3-fibered Seifert manifolds, and about the same time by Jeffrey [J1] and Garoufalidis [Ga1] who independently gave exact calculations of the semiclassical approximation of the  $\text{SU}(2)$ -invariants of lens spaces, starting from explicit expressions for the invariants. Jeffrey also verified parts of the conjecture (4) for  $G$  arbitrary and  $X$  belonging to a class of mapping tori of the torus. Garoufalidis verified parts of the conjecture (4) for  $G = \text{SU}(2)$  and  $X$  any Seifert fibered integral homology sphere.

The perturbative (or asymptotic) expansion of the Chern–Simons path integral  $Z_k^G(X)$  is given by the semiclassical approximation and its so-called higher loop correction terms. This leads together with the semiclassical approximation (5) to a conjecture for the (full) perturbative expansion of  $Z_k^G(X)$  in the large  $k$  limit, see [An2, Conjecture 7.6] for a version of this conjecture. Roughly speaking one should obtain an expansion given by the right-hand side of (5) with each integrand being multiplied by an asymptotic series of the form  $1 + \sum_{l=1}^{\infty} c_l^A r^{-l}$  with the  $c_l^A$ 's given in terms of certain contributions of Feynman diagrams determined by the Feynman rules of the Chern–Simons theory.

S. Axelrod and I. M. Singer have considered the higher loop contributions, cf. [AS1], [AS2], [Ax]. Actually they were able to deduce rigorously defined formulas for the coefficients  $c_l^A$  in case  $A$  is an acyclic point in  $\mathcal{M}$  or an element in a smooth component of  $\mathcal{M}$ . Moreover, they showed that their  $c_l^A$  define topological invariants of  $(X, A)$ . There are, however, major difficulties which have not yet been worked out. We refer to the above works of Axelrod and Singer and [An2] for more comments and details.

It is generally believed that the TQFTs of Reshetikhin and Turaev are a mathematical realization of Witten's TQFTs, and one can say that the AEC, Conjecture 1.1, offers in a sense a converse point of view to the above works on the perturbative expansion of the Witten invariants, where one seeks to derive the final output of perturbation theory after all cancellations have been made (i.e. collect all terms with the same Chern–Simons value). We stress that, although the ideas leading to the AEC stem from Witten's approach, the AEC is a conjecture for the rigorously defined RT-invariants, and it is completely independent of Witten's Chern–Simons path integral approach. Seen in the above light we should also expect the following slightly different version of the AEC, where we have separated terms coming from different components of  $\mathcal{M}$  (even if there are different components with the same Chern–Simons value).

**Conjecture 1.2** (J. E. Andersen [An1], [An2]. The asymptotic expansion conjecture (AEC)). *For any 3-manifold  $X$  there exist constants  $\tilde{D}_j \in \mathbb{Q}$ ,  $I_j \in \mathbb{Q} \bmod 8$ ,  $v_j \in \mathbb{R}_+$  for  $j = 1, 2, \dots, m$  and  $\tilde{a}_l^j \in \mathbb{C}$  for  $j = 1, 2, \dots, m$  and  $l = 1, 2, \dots$  such that the asymptotic expansion of  $\tau_r^G(X)$  in the limit  $r \rightarrow \infty$  is given by*

$$\tau_r^G(X) \sim \sum_{j=1}^m e^{2\pi i r q_j} r^{\tilde{D}_j} v_j e^{\frac{\pi i}{4} I_j} \left( 1 + \sum_{l=1}^{\infty} \tilde{a}_l^j r^{-l} \right),$$

where the  $j$ -sum is over the connected components of the moduli space of flat  $G$ -connections on  $X$  and  $q_j$  is the value of the Chern–Simons functional on the component indexed by  $j$ .

Here  $\sim$  means that there for all non-negative  $N$  is a  $C_N \in \mathbb{R}$  such that

$$\left| \tau_r^G(X) - \sum_{j=1}^m e^{2\pi i r q_j} r^{\tilde{D}_j} v_j e^{\frac{\pi i}{4} I_j} \left( 1 + \sum_{l=1}^N \tilde{a}_l^j r^{-l} \right) \right| \leq C_N r^{D-N-1}$$

for all  $r \in \mathbb{Z}_{\geq h^\vee}$ , where  $D = \max\{\tilde{D}_1, \dots, \tilde{D}_m\}$ .

We note that Conjecture 1.1 follows from Conjecture 1.2, and that  $D$  is equal in the two versions of the conjecture. The AEC concerns (analytic) asymptotic expansions of (a class of) complex functions defined on the positive integers. As noted by Andersen [An1], [An2], such a function has at most one asymptotic expansion of the form (1) (up to some trivialities such as permutations of the terms in the asymptotic formula). (Here it is of course important that the  $q_j$ 's are mutually different.) This means that if the AEC is true, then the  $D_j$ 's,  $b_j$ 's and the  $a_l^j$ 's in (1) are all uniquely determined by the function  $r \mapsto \tau_r^G(X)$ , hence they are also topological invariants of  $X$ . In particular Andersen has proposed the following conjecture (compare with (4) and (5)):

**Conjecture 1.3** (J. E. Andersen [An1], [An2]. Topological interpretation of the  $D_j$ 's). *Let  $\mathcal{M}_j$  be the union of the components of the moduli space of flat  $G$ -connections on  $X$  having Chern–Simons value  $q_j$ . Then*

$$D_j = \frac{1}{2} \max_{A \in \mathcal{M}_j} (h_A^1 - h_A^0), \quad (6)$$

where max here means the maximum value of  $h_A^1 - h_A^0$  on a Zariski open subset of  $\mathcal{M}_j$ .

The results in [An1] showed the importance of being careful about the interpretation of the max in this conjecture. The  $\tilde{a}_l^j$ 's,  $v_j$ 's,  $\tilde{I}_j$ 's and  $\tilde{D}_j$ 's in Conjecture 1.2 are also expected to be topological invariants. In particular,  $\tilde{D}_j$  should be given by the right-hand side of (6) with  $\mathcal{M}_j$  now being the  $j$ th component of the moduli space.

Conjecture 1.3 implies that the sequence  $\{\theta_l\}$  in (1) is the sequence of positive half-integers  $\frac{1}{2}\mathbb{Z}_{>0}$ . This conjecture gives (in combination with Conjecture 1.1) a strong connection between

the fundamental group of  $X$  and the quantum invariants of  $X$ , recalling that the moduli space of flat  $G$ -connections on  $X$  is in bijection with  $\text{Hom}(\pi_1(X), G)/G$  (for  $X$  connected). Garoufalidis [Ga2] and Andersen [An4] have e.g. pointed out that the above growth rate in  $r$  for the invariants  $\tau_r^{\text{SU}(n)}$  leads to the fact that the colored Jones polynomials associated to  $\text{SU}(n)$  (for all colorings and all  $n$ ) can distinguish the unknot from all other knots. As also pointed out by Andersen [An2], Conjecture 1.3 implies together with a recent result of Kronheimer and Mrowka [KM] that the colored Jones polynomials associated to  $\text{SU}(2)$  alone can distinguish the unknot from all other knots.

Andersen was the first to give a complete proof of the AEC for a class of 3-manifolds. In fact he proved in [An1] the AEC (together with Conjecture 1.3) for the mapping tori of finite order diffeomorphisms of orientable surfaces of genus at least 2 using the gauge theoretic approach to the quantum invariants. These manifolds are Seifert manifolds with orientable base and Seifert Euler number equal to zero. Andersen proved the conjectures for  $G$  an arbitrary simply connected compact simple Lie group.

As already stated Jeffrey [J1] and Garoufalidis [Ga1] made completely rigorous calculations of the semiclassical approximation of the  $\text{SU}(2)$ -invariants of lens spaces. Actually these calculations contain a complete verification of the AEC for the lens spaces for  $\text{SU}(2)$ . Moreover, they confirm Conjecture 1.3 in this case. The paper [J1] also contains a proof of the AEC for a certain class of mapping tori over the torus (for the invariants associated to an arbitrary simply connected compact simple  $G$ ). Except for the identification of Chern–Simons invariants, the AEC has recently been proved in [HT] for all lens spaces and  $G$  an arbitrary simply connected compact simple Lie group.

In [Ro1], [Ro3] Rozansky calculated the Witten  $\text{SU}(2)$ -invariants of all Seifert manifolds with orientable base and carried through a rather technical analysis leading to a candidate for the full asymptotic expansion of these invariants. However, the error estimates required to prove that his calculations lead to the asymptotics of the invariants are missing. In this paper we establish those error estimates thereby proving that Rozansky’s calculations in [Ro3] really leads to the large  $r$  asymptotic expansion of the quantum  $\text{SU}(2)$ -invariants of the Seifert manifolds. In [Ha2, Sect. 8] it was proved that the Witten  $\text{SU}(2)$ -invariants of the Seifert manifolds calculated by Rozansky are equal to the RT-invariants associated to  $\text{SU}(2)$ .

For a certain subclass of the Seifert manifolds Lawrence and Rozansky [LR] have obtained a much shorter calculation of the asymptotic expansion of the  $\text{SU}(2)$ -invariants. In this calculation one does not have to apply the method of this paper to establish the required error estimates. These estimates directly follow from the method without further work. The method (as it stands) only works for the Seifert manifolds satisfying that the first coordinates of the oriented Seifert invariants are mutually coprime (the oriented Seifert invariants being pairs of coprime integers, one for each singular fiber). In [LR] the result is only stated for the Seifert manifolds with base  $S^2$ , but their method works for all Seifert manifolds with orientable base or nonorientable base with even genus satisfying the above coprime condition. However, it does not seem to be easy to extend this method to cases where the coprime condition is not satisfied.

In [Y2] Yoshida defines a family of invariants  $Y_k^{\text{SU}(2)}$ ,  $k \in \mathbb{Z}_{\geq 0}$ , of 3-manifolds via the abelianization of the  $\text{SU}(2)$  WZW model obtained in [Y1]. Moreover, he calculates the semiclassical approximation of his invariants for the integral homology spheres satisfying that the moduli space  $\mathcal{M}$  of flat  $\text{SU}(2)$ -connections is finite set of acyclic points. This calculation includes a complete description of the semiclassical formula in terms of Chern–Simons invariants, spectral flows and Reidemeister torsions. In fact, he finds that the semiclassical approximation of his invariants for these manifolds is given by the right-hand side of (4) with  $G = \text{SU}(2)$  (and

$B_A = D_A = 0$  for all  $A$ ). It is expected (but not yet proved) that the invariant  $Y_k^{\text{SU}(2)}$  is equal to the invariant  $\tau_{k+2}^{\text{SU}(2)}$ .

Let us finally mention that we via recent private communication have learned that J. E. Andersen for the groups  $G = \text{SU}(n)$  has proved Conjectures 1.1 and 1.3 for all 3-manifolds via the gauge theoretic approach, see [An3] for some comments.<sup>2</sup> The proof involves asymptotics of Hitchin's connection over Teichmüller space, approximations to all orders, of the boundary states of handle bodies and techniques similar to the ones presented in [An3]. Where Andersen works with the gauge theoretic definition of the quantum invariants, we work with the definition of Reshetikhin and Turaev and our proof of the AEC for the Seifert manifolds is very different from Andersen's general proof.

Next let us give a review of the content of this paper. The main result is

**Theorem 1.4.** *The AEC, Conjecture 1.1, is true for  $G = \text{SU}(2)$  and  $X$  any Seifert manifold with orientable base of positive genus or non-orientable base with even genus. If  $X$  is a Seifert manifold with base  $S^2$  and  $n$  exceptional fibers, then the AEC holds if  $n \leq 3$  and in case  $n > 3$  the large  $r$  asymptotic expansion of  $\tau_r(X)$  has the form (1) with a set  $\{q_j\}_{j=0}^n$  containing the image set of the Chern–Simons functional on the moduli space of flat  $\text{SU}(2)$ –connections on  $X$ . In all cases the sequence  $\{\theta_l\} \subseteq \frac{1}{2}\mathbb{Z}$  in accordance with Conjecture 1.3.*

A part of this theorem was proved in the authors thesis [Ha1], where the case of Seifert manifolds with orientable base was handled. In this paper we calculate the asymptotics of a class of functions generalizing the invariants of the Seifert manifolds with orientable base or non-orientable base with even genus. Unfortunately certain parts of the proof do not go through as they stand for the remaining Seifert manifolds, i.e. the ones with non-orientable base of odd genus. If  $X$  is a Seifert manifold with base  $S^2$  the asymptotic expansion of  $\tau_r(X)$  has the form (1) as stated above, but some of the  $q_j$ 's are not Chern–Simons values. If the number of exceptional fibers is  $n \leq 2$ , i.e. if  $X$  is a lens space, it is easy to prove that all the terms in our asymptotic formula, corresponding to  $q_j$ 's which are not Chern–Simons values, are zero, thus proving the AEC for those manifolds. We also prove that this is the case for the Seifert manifolds with 3 exceptional fibers, cf. Theorem 4.10. We do not in this paper carry through the analysis needed to prove the AEC for the cases of  $n \geq 4$  exceptional fibers, see Sec. 4.2 for more details. Let us finally mention that we also prove the AEC for some of the Seifert manifolds with non-orientable base with odd genus, namely for the Seifert manifold with base  $\mathbb{RP}^2$  and zero or one exceptional fiber, cf. Corollary 4.11, this indicating that nothing special happens in the case of a non-orientable base of odd genus.

The proof of Theorem 1.4 is long and technical, but each step in the proof only uses elementary mathematics. Let us for the convenience of the reader give an outline of the proof here. The method used follows closely ideas of Rozansky [Ro3] and this paper owes much to Rozansky's work. As stated earlier we supplement the work of Rozansky by establishing certain required error estimates. Unfortunately the calculations needed to establish those estimates take up a big space explaining to a great extend the length of this paper.

Let  $\tau_r = \tau_r^{\text{SU}(2)}$  and let  $X$  be a Seifert manifold. By [Ha2, Theorem 8.4] (see Theorem 3.1) we have

$$\tau_r(X) = a(r) \sum_{\gamma=1}^{r-1} \frac{(-1)^{\gamma dg} h(\gamma)}{\sin^{n+dg-2} \left( \frac{\pi}{r} \gamma \right)}, \quad (7)$$

---

<sup>2</sup>It should thus be a theorem that the family of colored Jones polynomials associated to  $\text{SU}(2)$  can distinguish the unknot from all other knots.

where  $a(r) = br^{dg/2-1}e^{ic/r}$  for some constants  $c \in \mathbb{R}$  and  $b \in \mathbb{C} \setminus \{0\}$  (depending on  $X$ ). Here  $n$  is the number of singular fibers,  $g$  is the genus of the base of  $X$  and  $d = 2$  if this base is orientable and  $d = 1$  if it is non-orientable. (Here we say that the non-orientable connected sum  $\#^k \mathbb{RP}^2$  has genus  $k$ .) The function  $h$  is extendable to an entire function with some nice symmetry properties (see (62)).

Because of the simple form of  $a(r)$  we can concentrate on  $Z(X; r) = \tau_r(X)/a(r)$ . For technical reasons (see Remarks 5.1 and 5.2) we will assume that  $dg$  is even, so from now on  $X$  is a Seifert manifold with orientable base or nonorientable base of even genus.

The first task is to get rid of the dependence of  $r$  in the summation limit in (7). To this end we use that a function  $f : \mathbb{Z} \rightarrow \mathbb{C}$  periodic with a period of  $P$  satisfies

$$\sum_{k=0}^{P-1} f(k) = P \lim_{\varepsilon \rightarrow 0+} \sqrt{\varepsilon} \sum_{k \in \mathbb{Z}} e^{-\pi \varepsilon k^2} f(k). \quad (8)$$

This leads together with some other symmetry considerations to the formula

$$Z(X; r) = r \lim_{\varepsilon \rightarrow 0+} \sqrt{\varepsilon} \lim_{\xi \rightarrow 0+} \sum_{\gamma \in \mathbb{Z}} e^{-\pi \varepsilon \gamma^2} \frac{h(\gamma)}{\sin^{n+dg-2} \left( \frac{\pi}{r} (\gamma - i\xi) \right)}, \quad (9)$$

where we have introduced an extra parameter  $\xi$  to avoid singularities. (If there are no singular fibers, i.e. if  $n = 0$ , then we have to include the extra factor

$$\frac{\sin \left( \frac{\pi}{r} \gamma \right)}{\sin \left( \frac{\pi}{r} (\gamma - i\xi) \right)}$$

in the summand.)

To proceed we change (9) to a sum of integrals by using the Poisson summation formula

$$\sum_{k \in \mathbb{Z}} \varphi(k) = \sum_{m \in \mathbb{Z}} \int_{-\infty}^{\infty} e^{2\pi i m x} \varphi(x) dx. \quad (10)$$

It is well-known that this formula is valid for functions  $\varphi$  in the Schwartz space  $\mathcal{S}(\mathbb{R})$  of smooth functions, that, together with their derivatives, are rapidly decreasing at infinity, see e.g. [Ho]. We use Poisson's formula with  $\varphi(\gamma) = e^{-\pi \varepsilon \gamma^2} \frac{h(\gamma)}{\sin^{n+dg-2} (\pi(\gamma - i\xi)/r)}$  (with fixed  $\xi$  and  $\varepsilon$ ). This is in  $\mathcal{S}(\mathbb{R})$  because  $h$  is periodic and (putting  $z = \gamma/r$ ,  $\delta = \varepsilon r^2$  and  $\eta = \xi/r$ ) we arrive at an expression of the form

$$Z(X; r) = r \lim_{\delta \rightarrow 0+} \sqrt{\delta} \lim_{\eta \rightarrow 0+} \sum_{m \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) \int_{-\infty}^{\infty} e^{-\pi \delta z^2} \tilde{\kappa}(z) e^{r(Q(z) + 2\pi i m z)} dz, \quad (11)$$

where  $\Lambda$  is a finite set (only depending on  $X$ ),  $Q$  is a polynomial in  $z$  with coefficients only depending on  $\lambda$  and  $X$ , and

$$\tilde{\kappa}(z) = \frac{e^{ibz}}{\sin^k (\pi(z - i\eta))},$$

where  $b \in \mathbb{R}$  only depends on  $\lambda$  and  $X$  and  $k = n + dg - 2$ . Put  $\kappa(z) = e^{-\pi \delta z^2} \tilde{\kappa}(z)$  and  $Q_m(z) = Q(z) + 2\pi i m z$  in the following. We note that  $Q_m$  is of degree 1 if the Seifert Euler number  $E$  of  $X$  is equal to zero and of degree 2 if  $E \neq 0$ . This makes the asymptotic analysis in the case  $E = 0$  different from (and also simpler than) that in the case  $E \neq 0$ . To make the notation in the following compatible with the notation used in Sec. 5 and 6, we put  $A = \pi \frac{E}{2}$ .

Let us first assume that  $E \neq 0$ . In this case the integrals  $I(r) = \int_{-\infty}^{\infty} \kappa(z) e^{rQ_m(z)} dz$  in (11) are approximated by using the steepest descent method. Thus, if  $z_{\text{st}}$  denotes the stationary point of  $Q_m$ , i.e.  $Q'_m(z_{\text{st}}) = 0$ , we deform the integration contour (here the real axis) to a new contour



$C$  passing through  $z_{\text{st}}$  in such a way that the large  $r$  asymptotics of  $\int_C \kappa(z) e^{rQ_m(z)} dz$  is easy to obtain by standard methods (such as the Laplace method), see e.g. [BH], [B], [Wo]. It can be shown that a good such contour is a contour  $C_{\text{sd}}$  along which the imaginary part of  $Q_m$  is constant. In the case of  $I(r)$  this so-called steepest descend contour is the straight line

$$\text{Im}(z) = \text{sign}(A)(\text{Re}(z) - z_{\text{st}}). \quad (12)$$

(We note that the stationary point  $z_{\text{st}}$  is non-degenerate, i.e.  $Q_m''(z_{\text{st}}) \neq 0$ , and that this point and thereby  $C_{\text{sd}}$  depend on  $(m, \lambda)$ .) To determine the asymptotic expansion of  $I(r)$  in the limit  $r \rightarrow \infty$  we need to calculate the large  $r$  asymptotics of the new contour integral  $J(r) = \int_{C_{\text{sd}}} \kappa(z) e^{rQ_m(z)} dz$ , and of the difference  $I(r) - J(r)$ . This difference is zero for  $k \leq 0$  and is for  $k > 0$  given by the infinite sum of residues in the poles of  $\kappa(z)$  crossed when deforming the real axes to  $C_{\text{sd}}$ , i.e. the poles  $z_l = l + i\eta$ ,  $l \in \mathbb{Z}$ , for which

$$\text{sign}(A)(l - z_{\text{st}}) > \eta. \quad (13)$$

(We refer to Lemma 5.4.) By further showing that certain infinite sums are absolutely convergent (see Lemma 5.5), we therefore end up with the following partition of  $Z(X; r)$ :

$$Z(X; r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} (Z_{\text{int},1}(\delta, \eta) + Z_{\text{int,polar}}(\delta, \eta) + Z_{\text{polar},1}(\delta, \eta)), \quad (14)$$

where

$$\begin{aligned} Z_{\text{int},1}(\delta, \eta) &= \sum_{(m,\lambda) \in W} g(\lambda) \int_{C_{\text{sd}}(m,\lambda)} \kappa(z) e^{rQ_m(z)} dz, \\ Z_{\text{int,polar}}(\delta, \eta) &= \sum_{\lambda \in \Lambda} g(\lambda) \sum_{\substack{m \in \mathbb{Z} \\ z_{\text{st}}(m,\lambda) \in \mathbb{Z}}} \int_{C_{\text{sd}}(m,\lambda)} \kappa(z) e^{rQ_m(z)} dz, \\ Z_{\text{polar},1}(\delta, \eta) &= 2\pi i \sum_{\lambda \in \Lambda} g(\lambda) \sum_{m \in \mathbb{Z}} \sum_{\substack{l \in \mathbb{Z} \\ \text{sign}(A)(l - z_{\text{st}}(m,\lambda)) > \eta}} \text{Res}_{z=z_l(\eta)} \left\{ \kappa(z) e^{rQ_m(z)} \right\}, \end{aligned} \quad (15)$$

where  $W = \{(m, \lambda) \in \Lambda \times \mathbb{Z} \mid z_{\text{st}}(m, \lambda) \notin \mathbb{Z}\}$ . (Here we suppress the dependency on  $r$  and also to some extent the dependency on  $\lambda$  in our notation for not making it too clumsy.) The above sums are all absolutely convergent. By rearranging the terms in the sum  $Z_{\text{polar},1}(\delta, \eta)$  we obtain an expression of the form (see Lemma 5.5)

$$Z_{\text{polar},1}(\delta, \eta) = 2\pi i \sum_{\lambda \in \Lambda} g(\lambda) \sum_{l \in \mathbb{Z}} \beta(l) \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi}l > \frac{|A|}{\pi}\eta}} \text{Res}_{z=i\eta} \left\{ \tilde{\phi}(z; l, \lambda) e^{2\pi i r m z} \right\},$$

where  $B/\pi$  is a certain rational number only depending on  $\lambda$  and  $X$ , and  $\tilde{\phi}$  is a meromorphic function independent of  $m$ . Along the same lines we obtain the expression

$$Z_{\text{int,polar}}(\delta, \eta) = \sum_{\lambda \in \Lambda} g(\lambda) \sum_{l \in \mathbb{Z}} \beta(l) \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi}l = 0}} \int_{C(\lambda, 0)} \tilde{\phi}(z; l, \lambda) e^{2\pi i r m z} dz,$$

where  $C(\lambda, \rho)$ ,  $\rho \in \mathbb{R}$ , is the line

$$\text{Im}(z) = \text{sign}(A)\text{Re}(z) - \rho.$$

We have a decomposition (see Lemma 5.6 for details)

$$Z_{\text{int,polar}}(\delta, \eta) = Z_{\text{polar},2}(\delta, \eta) + Z_{\text{int},2}(\delta, \eta),$$

where  $Z_{\text{polar},2}(\delta, \eta)$  is given by a sum of residues like  $Z_{\text{polar},1}(\delta, \eta)$ , in fact

$$Z_{\text{polar},2}(\delta, \eta) = \pi i \sum_{\lambda \in \Lambda} g(\lambda) \sum_{l \in \mathbb{Z}} \beta(l) \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi} l = 0}} \text{Res}_{z=i\eta} \left\{ \tilde{\phi}(z; l, \lambda) e^{2\pi i r m z} \right\},$$

and

$$Z_{\text{int},2}(\delta, \eta) = \sum_{\lambda \in \Lambda} g(\lambda) \sum_{\substack{l \in \mathbb{Z} \\ \frac{B}{2\pi} + \frac{A}{\pi} l \in \mathbb{Z}}} \beta(l) \int_{C(\lambda, \eta)} \tilde{F}(y; l, \lambda) dy, \quad (16)$$

where  $\tilde{F}(y)$  is a meromorphic function with poles in  $\mathbb{Z}$  (if  $k > 0$ ) (explaining the name  $Z_{\text{int},\text{polar}}$ ). The next observation is that there exists a  $\eta_0 > 0$  such that

$$Z_{\text{polar},1}(\delta, \eta) = 2\pi i \sum_{\lambda \in \Lambda} g(\lambda) \sum_{l \in \mathbb{Z}} \beta(l) \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi} l > 0}} \text{Res}_{z=i\eta} \left\{ \tilde{\phi}(z; l, \lambda) e^{2\pi i r m z} \right\},$$

for all  $\eta \in ]0, \eta_0]$ . By letting  $Z_{\text{polar}}(\delta, \eta) = Z_{\text{polar},1}(\delta, \eta) + Z_{\text{polar},2}(\delta, \eta)$  we therefore obtain

$$Z(X; r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} (Z_{\text{int},1}(\delta, \eta) + Z_{\text{int},2}(\delta, \eta) + Z_{\text{polar}}(\delta, \eta)), \quad (17)$$

where  $Z_{\text{int},1}(\delta, \eta)$  and  $Z_{\text{int},2}(\delta, \eta)$  are given respectively by (15) and (16) and

$$Z_{\text{polar}}(\delta, \eta) = 2\pi i \sum_{\lambda \in \Lambda} g(\lambda) \sum_{l \in \mathbb{Z}} \beta(l) \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi} l \geq 0}} \text{Res}_{z=i\eta} \left\{ \frac{\tilde{\phi}(z; l, \lambda) e^{2\pi i r m z}}{\text{Sym}_{\pm}(m + \frac{B}{2\pi} + \frac{A}{\pi} l)} \right\}. \quad (18)$$

Here we have introduced the function  $\text{Sym}_{\pm} : \mathbb{R} \rightarrow \{1, 2\}$  (borrowing notation from [Ro3]) which is 1 in all points except in zero where it is 2.

The next natural step would be to try to calculate the limit  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} f(\delta, \eta)$  (or at least the large  $r$  asymptotics of that limit) for  $f$  equal to each of the functions  $Z_{\text{int},1}$ ,  $Z_{\text{int},2}$ , and  $Z_{\text{polar}}$  (assuming these limits exist). Thus we start by showing that

$$Z_{\text{polar}}(r) := r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}(\delta, \eta) \quad (19)$$

exists by calculating this limit explicitly.

To handle the two other limits  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{int},\nu}(\delta, \eta)$ ,  $\nu = 1, 2$ , more care need to be taken. Since the limits on the right-hand sides of (17) and (19) exist we have that

$$Z_{\text{int}}(r) := r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{int}}(\delta, \eta) \quad (20)$$

exists, where  $Z_{\text{int}}(\delta, \eta) = Z_{\text{int},1}(\delta, \eta) + Z_{\text{int},2}(\delta, \eta)$ . However, we do not calculate this limit explicitly, but only give it an asymptotic description. This is of course sufficient for the proof of Theorem 1.4. To find the asymptotic expansion of  $Z_{\text{int}}(r)$  we show that we for each  $N \in \mathbb{Z}_{\geq 0}$  have a decomposition

$$Z_{\text{int},\nu}(\delta, \eta) = Z_{\text{int},\nu}(N; \delta, \eta) + R_{\nu}(N; \delta, \eta) \quad (21)$$

such that

$$Z_{\text{int},\nu}(r; N) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{int},\nu}(N; \delta, \eta)$$

exists and can be calculated explicitly,  $\nu = 1, 2$ . This implies that

$$\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} (R_1(N; \delta, \eta) + R_2(N; \delta, \eta))$$

exists. We will not show that  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} R_\nu(N; \delta, \eta)$  exists for each  $\nu = 1, 2$  separately. Instead we show that  $R_\nu(N; \delta, \eta)$  is bounded from above by a certain function  $A_\nu(N; \delta, \eta)$  for which

$$r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} A_\nu(N; \delta, \eta)$$

can easily be calculated and shown to be small in the correct asymptotic sense compared to  $Z_{\text{int}, \nu}(r; N)$  in the limit of large  $r$ ,  $\nu = 1, 2$ . This proves that  $Z_{\text{int}, 1}(r; N) + Z_{\text{int}, 2}(r; N)$  is the large  $r$  asymptotics of  $Z_{\text{int}}(r)$  to an order depending on  $N$ .

We observe that all the sums  $Z_{\text{polar}}(\delta, \eta)$ ,  $Z_{\text{int}, 1}(\delta, \eta)$ , and  $Z_{\text{int}, 2}(\delta, \eta)$  are infinite. (For the sum  $Z_{\text{int}, 2}(\delta, \eta)$  note here that  $A/\pi \in \mathbb{Q}$  and that  $B(\lambda)/(2\pi) \in \mathbb{Z}$  for some  $\lambda$ .) According to the AEC we should end up with a finite sum. Basically these infinite sums are changed to finite sums by using the periodicity result (8) ‘backwards’. The sum  $Z_{\text{polar}}(\delta, \eta)$  in (18) actually contains two infinite sums. We calculate one of these, namely the sum over  $m$ , explicitly. Let us give some further details. First we change  $Z_{\text{polar}}(\delta, \eta)$  to an expression of the form

$$Z_{\text{polar}}(\delta, \eta) = \sum_{l \in \mathbb{Z}} \sum_{\nu \in \Gamma(l)} \sum_{\substack{m \in \mathbb{Z} \\ m - V(\nu) \geq 0}} \frac{\text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} G(z; l, \nu, \eta) e^{2\pi i r m z} \right\}}{\text{Sym}_\pm(m - V(\nu))}, \quad (22)$$

where  $\Gamma(l)$  is a finite index set depending on  $l$  in a periodic way, and  $G$  is periodic w.r.t.  $l$ . We use certain symmetries to establish that. Now, if  $a \in \mathbb{R}$  and  $f : \mathbb{Z} \rightarrow \mathbb{C}$  then

$$\sum_{\substack{m \in \mathbb{Z} \\ m \geq a}} \frac{f(m)}{\text{Sym}_\pm(m - a)} = \sum_{m=0}^{\infty} \frac{f(m)}{\text{Sym}_\pm(m)} - \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |a|}} \frac{\text{sign}(a) f(\text{sign}(a)m)}{\text{Sym}_\pm(m) \text{Sym}_\pm(m - |a|)}.$$

in the sense that if the left-hand side is convergent, then the infinite sum in the right-hand side is also convergent and the identity holds. Using this identity on (22) together with

$$\sum_{m=0}^{\infty} \frac{e^{2\pi i r m z}}{\text{Sym}_\pm(m)} = \frac{i}{2} \cot(\pi r z),$$

valid for all  $z$  in the open disk in the complex plane with center  $i\eta$  and radius  $\eta$ , we thus obtain

$$\begin{aligned} Z_{\text{polar}}(\delta, \eta) &= \sum_{l \in \mathbb{Z}} \sum_{\nu \in \Gamma(l)} \left( \frac{i}{2} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} G(z; l, \nu, \eta) \cot(\pi r z) \right\} \right. \\ &\quad \left. - \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |V|}} \frac{\text{sign}(V) \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} G(z; l, \nu, \eta) e^{2\pi i r \text{sign}(V)m z} \right\}}{\text{Sym}_\pm(m) \text{Sym}_\pm(m - |V|)} \right). \end{aligned} \quad (23)$$

From this expression we obtain that  $Z_{\text{polar}}(\delta) := \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}(\delta, \eta)$  is equal to the right-hand side of (23) with  $\eta = 0$ . Finally  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} Z_{\text{polar}}(\delta)$  is calculated by using (8). We thus obtain an exact expression for  $Z_{\text{polar}}(r)$  equal to some Laurent polynomial in  $r$ .

Let us also give a few more details concerning the calculation of the asymptotic expansion of  $Z_{\text{int}}(r)$  in (20). First we observe that

$$Z_{\text{int}, 1}(\delta) := \lim_{\eta \rightarrow 0_+} Z_{\text{int}, 1}(\delta, \eta) = \sum_{(m, \lambda) \in W} g(\lambda) \int_{C_{\text{sd}}(m, \lambda)} \kappa_0(z) e^{r Q_m(z)} dz,$$

where  $\kappa_0(z)$  is equal to  $\kappa_\eta(z)$  with  $\eta = 0$ . (For  $\nu = 1$  we thus calculate this limit before making the decomposition (21).) For an arbitrary  $(m, \lambda) \in W$  we can expand  $\kappa_0$  as a power series in a small disk around  $z_{\text{st}}(m, \lambda)$  (with a radius independent of  $(m, \lambda)$ ). Let  $N$  be an arbitrary but

fixed non-negative integer, and let  $\kappa_0(z; N)$  be equal to the first  $N+1$  terms in this power series. Then we let

$$Z_{\text{int},1}(N; \delta) = \sum_{(m,\lambda) \in W} g(\lambda) \int_{C_{\text{sd}}(m,\lambda)} \kappa_0(z; N) e^{rQ_m(z)} dz.$$

The remainder term (to order  $N$ ) is then simply given by  $R_1(N; \delta) = Z_{\text{int},1}(\delta) - Z_{\text{int},1}(N; \delta)$ . Since the power series expansion of  $\kappa_0(z)$  is only valid in a small neighborhood of  $z_{\text{st}}(m, \lambda)$  this remainder term is not simply given by replacing  $\kappa_0(z)$  by the remaining terms in the power series expansion in the expression for  $Z_{\text{int},1}(\delta)$ , see Lemma 5.13 for details.

The decomposition (21) for  $\nu = 2$  is obtained in a similar way by representing a preexponential factor of the integrand  $\tilde{F}$  as a power series around  $y = 0$ . However, in this case the positive parameter  $\eta$  plays a crucial role and the limit  $\lim_{\eta \rightarrow 0_+}$  can only be calculated in the final step.

Let us finally say a few words about the proof of Theorem 1.4 in the case  $E = 0$ . Again we obtain the expression (11). However, in this case  $Q(z) = 2\pi i a z$ , where  $a$  is a constant depending on  $\lambda$  but not on  $m$ , and the steepest descent method can not be used to calculate the large  $r$  asymptotics of the integrals  $\int_{-\infty}^{\infty} \kappa(z) e^{rQ_m(z)} dz$  since  $Q_m(z) = Q(z) + 2\pi i m z$  has no stationary points in this case. Instead we show that there exist certain contours  $C(m, \lambda)$  so that

$$Z(X; r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} (Z_{\text{int}}(\delta, \eta) + Z_{\text{polar}}(\delta, \eta)),$$

where

$$\begin{aligned} Z_{\text{int}}(\delta, \eta) &= \sum_{\lambda \in \Lambda} g(\lambda) \sum_{m \in \mathbb{Z}} \int_{C(m,\lambda)} \kappa(z) e^{rQ_m(z)} dz, \\ Z_{\text{polar}}(\delta, \eta) &= 2\pi i \sum_{\lambda \in \Lambda} g(\lambda) \sum_{\substack{m \in \mathbb{Z} \\ m-a(\lambda) \geq 0}} \sum_{l \in \mathbb{Z}} \text{Res}_{z=z_l(\eta)} \left\{ \kappa(z) e^{rQ_m(z)} \right\}. \end{aligned}$$

By estimating the integrals inside the sum  $Z_{\text{int}}(\delta, \eta)$  it can be proved that

$$\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{int}}(\delta, \eta) = 0.$$

Hence

$$Z(X; r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}(\delta, \eta),$$

and this double limit can be calculated explicitly along the same lines as in the case  $E \neq 0$ .

From a general point of view the main problem when calculating the large  $r$  asymptotics of the quantum invariants as defined by Reshetikhin and Turaev, is that these invariants are given by a finite sum in which both the number of terms and the terms themselves depend on the level  $r$ . It has been speculated that there is some kind of number theoretic principle which can be used to calculate these asymptotics. In [J1] Jeffrey revealed that one can calculate the asymptotics of the  $\text{SU}(2)$ -invariants of lens spaces by an inductive argument using a reciprocity formula for Gauss sums. This reciprocity formula follows from an application of the Poisson summation formula, which is a special version of the Fourier transform (namely the Fourier transform of the sum of Dirac measures on some discrete subgroup of  $\mathbb{R}^n$ ). More generally Jeffrey speculated that the Fourier transform may be the basic mechanism behind identities needed for calculating the asymptotics.

Rozansky [Ro1], [Ro3] took up these ideas and showed how to use the Poisson summation formula to calculate the asymptotics for the broader class of Seifert manifolds with orientable base space. For a general Seifert manifold the main idea is as illustrated above to change the finite sum formula for the invariant to an infinite sum using certain periodicity results. This involves introducing certain limits with respect to certain parameters. One can then use the

Poisson summation formula on the infinite sum. This leads to a formula for the invariant given by an infinite sum of integrals. Each of these integrals can be given an asymptotic description via the steepest descent method. As proved in this paper this leads ultimately to an asymptotic formula like (1).

For the calculations of the asymptotics of the  $SU(2)$ -invariants of Seifert manifolds one only needs a one-dimensional Poisson summation formula (involving only a sum over  $\mathbb{Z}$ ), see (10). In [J1] Jeffrey also obtained a multi-dimensional reciprocity formula for Gauss sums using a multi-dimensional version of the Poisson formula (see also [J2] and [HT, Appendix A]). She then used this to calculate the asymptotics of the  $G$ -invariants of some mapping tori over the torus,  $G$  being any compact simply connected simple Lie group. In [HT] the author and Takata use the multi-dimensional reciprocity formula to calculate the asymptotics of the  $G$ -invariants of the lens spaces by a similar approach as in [J1].

In [AH], Andersen and the author investigate the asymptotics of the  $SU(2)$ -invariants of the 3-manifolds obtained by rational surgeries on the 3-sphere along the figure 8 knot. In these calculations the Fourier transform is not used. Instead the asymptotics is calculated by first obtaining a formula for the invariant given by a (finite) sum of certain contour integrals and then use the saddle point approximation method on each of the integrals.

It seems that a general principle behind calculating the large quantum level asymptotics of the RT-invariants of 3-manifolds is to obtain in some way or the other a formula of the invariants given by some sum of integrals, where the number of terms in the sum (allowed to be infinite) is independent of the level  $r$ , and where the integrals can be asymptotically described by using some well-known approximation method such as the method of steepest descent or more generally the saddle point method. The involved integrals will be of a form  $\int_C \psi(z) e^{r\phi(z)} dz$ , where  $C$  is some (multi-dimensional) contour in  $\mathbb{C}$  ( $\mathbb{C}^n$ ) and the phase function  $\phi$  and the function  $\psi$  are some complex functions independent of the level  $r$ . Note that even for the case of lens spaces and mapping tori over the torus, where a reciprocity formula for Gauss sums is used, one actually uses this approach, namely, the involved reciprocity formula can be proved by this approach, see e.g. [HT, Appendix]. The calculations in [AH] actually first lead to an expression for the invariants given by a finite sum of integrals, where the integrands are not of the nice form  $\int_C \psi(z) e^{r\phi(z)} dz$ . One has to make certain initial approximations to obtain an (approximative) expression for the invariants given by a finite sum of integrals of the nice form. This phenomenon should be expected in general.

The paper is organized as follows. In Sec. 2 we introduce the notions from asymptotic analysis needed. In this section we study from a general perspective asymptotic expansions of the form (1). These expansions constitute a certain subclass of a class of asymptotic expansions which we denote asymptotic expansions of generalized Poincaré type, since they are finite sums with each term being an asymptotic expansion of (ordinary) Poincaré type. A small but important result is Theorem 2.3, which extend the well-known uniqueness property Lemma 2.1 for asymptotics of Poincaré type to asymptotics of generalized Poincaré type. In Sec. 3 we introduce notation for the Seifert manifolds and state from [Ha2] formulas for the RT-invariants associated to  $SU(2)$  of all Seifert manifolds. In Sec. 4 we present detailed formulas for the large  $r$  asymptotic expansion of  $\tau_r^{SU(2)}(X)$ ,  $X$  a Seifert manifold as in Theorem 1.4. Moreover, we review results of Auckly [Au1] on the classical  $SU(2)$  Chern–Simons theory on Seifert manifolds. (We correct a small error in Auckly’s result, see Theorem 4.8 and below.) We demonstrate that Auckly’s results combine with our formulas for the asymptotic expansions to give a proof of Theorem 1.4. Finally, we identify for the Seifert fibered rational homology spheres the Casson–Walker invariant in the asymptotic formula using results of Lescop [Les]. In Sec. 5 and 6 the proofs of Theorems 4.1 and 4.4 are given. We focus on the main ideas deferring technicalities to appendices.

We end this introduction by a side remark. The part of the theory of quantum invariants concerned with asymptotic expansions is often denoted the perturbative theory, and (parts of) the asymptotic expansions of the RT-invariants (or Witten's invariants) are called perturbative quantum invariants. One should note, however, that the notion of perturbative invariants has different meanings in the mathematical literature. There are the analytic asymptotics studied in this paper, and there are the algebraically defined perturbative invariants of Ohtsuki [Oh1], [Oh2]. Note that Ohtsuki's perturbative invariant of a 3-manifold  $X$  is trivial unless  $X$  is a rational homology sphere [Oh2, Remark 1.3]. This is certainly not the case for the analytic asymptotics (as the results of this paper reveals). It is, however, generally believed that these two notions of perturbative invariants are closely related. It is speculated that the coefficients in the formal Ohtsuki series determine (some of) the coefficients in the analytic asymptotic expansion of the RT-invariants of rational homology spheres, see [Oh2, Example 1.6 and Sect. 5], [L], [Ro4], [Ro5] and [Oh3, Sect. 7] for some results and conjectures. The results in this paper should be useful to investigate this question for the Seifert fibered rational homology spheres. We will however not pursue this issue further in this paper. Recently K. Habiro has proposed new invariants for integral homology spheres which according to a conjecture of Habiro and Le should determine both Ohtsuki's perturbative invariants and the RT-invariants for these manifolds, see [Oh3, Conjecture 7.29]. (As stated in [Oh3, p. 492] this should now be a theorem of Habiro and Le.) Let us finally mention that there seem to be some very interesting number theoretic aspects of the asymptotics of the Seifert fibered integral homology spheres. We will not consider this issue in this paper but refer to papers of Lawrence and Zagier [LZ] and Hikami [H1]–[H3].

**Acknowledgements** The author thanks Institut de Recherche Mathématique Avancée (IRMA) at Université Louis Pasteur and C.N.R.S., Strasbourg, the School of Mathematics at the University of Edinburgh, the Max-Planck-Institut für Mathematik in Bonn, and the Department of Mathematical Sciences, University of Aarhus, for their hospitality during this work. He was supported by the European Commission, the Danish Natural Science Research Council and the Max-Planck-Institut für Mathematik. In addition this article is based upon work supported by the National Science Foundation under Grant No. EPS-0236913 and matching support from the State of Kansas through Kansas Technology Enterprise Corporation. The paper extends results obtained in the authors thesis [Ha1], and he would like to thank his thesis advisor J. E. Andersen for having given valuable comments to earlier versions of this paper, and also for the many helpful conversations about quantum invariants in general and asymptotic expansions of these in particular.

## 2. ASYMPTOTIC EXPANSIONS OF QUANTUM TYPE

In this section we study asymptotics of the form (1) from a general point of view. We shall call such asymptotics for asymptotic expansions of quantum type, see Definition 2.4. In this section we shall mention the few facts from asymptotic analysis needed to study them. For basic introductions to asymptotic analysis in general, see e.g. [BH], [B], [O], and [Wo].

Let in the following  $A = \mathbb{Z}_{\geq h} = \{m \in \mathbb{Z} | m \geq h\}$  for some fixed positive integer  $h$ . Recall the usual order-notation denoted by the  $O$ - and  $o$ -symbols: If  $f, g : A \rightarrow \mathbb{C}$  are two functions, then  $g = O(f)$  if there exists a constant  $C$  such that  $|f(k)| \leq C|g(k)|$  for all  $k \in A$ . Of course such a condition only puts limits on the large  $k$  behaviour of  $f(k)$ . We say that  $g = o(f)$  in the limit  $k \rightarrow \infty$  if for all  $\varepsilon > 0$  there exists a  $k_\varepsilon \in A$  such that  $|g(k)| \leq \varepsilon|f(k)|$  for all  $k \geq k_\varepsilon$ . If  $f(k) \neq 0$  for all  $k$  (or for all sufficiently large  $k$ ) this is equivalent to  $g(k)/f(k) \rightarrow 0$  as  $k \rightarrow \infty$ . As examples we have  $k^{-(n+1)} = o(k^{-n})$  in the limit  $k \rightarrow \infty$  and  $k^n + \log(k)k^{n-1} = O(k^n)$  for every  $n \in \mathbb{Z}$ . A sequence of functions  $\{\varphi_n : A \rightarrow \mathbb{C}\}_{n=0}^\infty$  is called an *asymptotic sequence* as

$k \rightarrow \infty$  if for all  $n \geq 0$

$$\varphi_{n+1}(k) = o(\varphi_n(k))$$

as  $k \rightarrow \infty$  and if we for all  $n$  have that  $\varphi_n(k) \neq 0$  for all sufficiently large  $k$ . For functions  $f : A \rightarrow \mathbb{C}$  and  $f_n : A \rightarrow \mathbb{C}$ , the formal series  $\sum_{n=0}^{\infty} f_n(k)$  is called a *generalized asymptotic expansion* of  $f(k)$  with respect to the asymptotic sequence  $\{\varphi_n(k)\}$ , as  $k \rightarrow \infty$ , if

$$f(k) = \sum_{n=0}^N f_n(k) + o(\varphi_N(k)) \quad (24)$$

in the limit  $k \rightarrow \infty$  for every  $N \geq 0$ . In this case we write

$$f(k) \sim \sum_{n=0}^{\infty} f_n(k); \quad \{\varphi_n\}, \quad k \rightarrow \infty. \quad (25)$$

When  $f_n(k) = a_n \varphi_n(k)$ ,  $a_n$  a fixed complex number, for every  $n$ , then the above expansion is said to be of *Poincaré type*. Furthermore, if the expansion is of this type, and  $\varphi_n(k) = (\xi(k))^{\lambda_n}$ ,  $\lambda_n$  a fixed complex number, the expansion is said to be of *power series type*. We have the following immediate uniqueness result for asymptotic expansions of Poincaré type (see e.g. [BH, pp. 16-17]).

**Lemma 2.1.** *Let  $\{\varphi_n\}_{n \in \mathbb{N}}$  be an asymptotic sequence and let  $f : A \rightarrow \mathbb{C}$ . Then there is at most one sequence of complex numbers  $a_0, a_1, a_2, \dots$  such that*

$$f(k) \sim \sum_{n=0}^{\infty} a_n \varphi_n(k); \quad \{\varphi_n\}, \quad k \rightarrow \infty.$$

□

The following definition gives a generalization of asymptotic expansions of Poincaré type and power series type suitable for our needs.

**Definition 2.2.** A function  $f : A \rightarrow \mathbb{C}$  is said to have an *asymptotic expansion of generalized Poincaré type* (w.r.t. the asymptotic sequence  $\{\varphi_n(k)\}$ ) if we for all nonnegative integers  $N$  have

$$f(k) = \left[ \sum_{j=1}^M e^{2\pi i k q_j} \sum_{m=0}^N c_m^j \varphi_m(k) \right] + o(\varphi_N(k)) \quad (26)$$

in the limit  $k \rightarrow \infty$ , where  $M$  is a non-negative integer,  $q_j \in [0, 1[$  and  $c_m^j \in \mathbb{C}$ . If, moreover,  $\varphi_m(k) = (\xi(k))^{\lambda_m}$ ,  $\lambda_m$  fixed complex numbers, then we will say that  $f$  has an *asymptotic expansion of generalized power series type*. The asymptotic expansion in (26) is called *trivial* if  $c_m^j = 0$  for all  $j$  and  $m$  (this case being equivalent to the case  $M = 0$ , i.e. the case where the sum  $\sum_{j=1}^M$  is put equal to zero).

By the notation (25), the identities (26) can also be written

$$f(k) \sim \sum_{m=0}^{\infty} \left( \sum_{j=1}^M e^{2\pi i k q_j} c_m^j \right) \varphi_m(k); \quad \{\varphi_m\}, \quad k \rightarrow \infty.$$

We immediately get that the set of functions  $f : A \rightarrow \mathbb{C}$  having an asymptotic expansion of generalized Poincaré type w.r.t. a fixed asymptotic sequence is a subspace of the  $\mathbb{C}$ -vector space of all functions  $f : A \rightarrow \mathbb{C}$ . (The zero function  $0 : \mathbb{N} \rightarrow \mathbb{C}$  has trivially a trivial asymptotic expansion of generalized Poincaré type (w.r.t. any asymptotic sequence).)

It is natural not to distinguish between asymptotic expansions of generalized Poincaré type which can be obtained from each other by trivial means. We will thus say that two such asymptotics are equivalent if the one can be obtained from the other by a finite sequence of the following operations: i) interchange two terms in the sum  $\sum_{j=1}^M$  in (26), ii) remove or add a term in  $\sum_{j=1}^M$  with  $c_m^j = 0$  for all  $m$  and iii) collect two terms with the same  $q_j$ -value. These operations also lead to a kind of minimal representation for each equivalence class of asymptotic expansions of generalized Poincaré type, namely if we have such an asymptotics we will say that it is on minimal form if it has no terms with  $c_m^j = 0$  for all  $m$  and if the  $q_j$ 's are mutually different. Thus, up to the summation order in  $\sum_{j=1}^M$  each equivalence class has a unique representative on minimal form. In particular, a trivial expansion is on minimal form if and only if  $M = 0$  in (26). We shall not distinguish between an asymptotic expansion of generalized Poincaré type and its equivalence class.

The following theorem extends the uniqueness property for asymptotic expansions of Poincaré type, Lemma 2.1, to asymptotic expansions of generalized Poincaré type. The proof is very short due to an idea of Pieter Moree.

**Theorem 2.3.** *Let  $f : A \rightarrow \mathbb{C}$  be an arbitrary function. Then  $f$  has at most one asymptotic expansion of generalized Poincaré type w.r.t. a given asymptotic sequence. The zero function has only the trivial asymptotic expansions of generalized Poincaré type.*

Said in another way: If  $f : A \rightarrow \mathbb{C}$  has an asymptotic expansion of generalized Poincaré type as in (26) and if this expansion is on minimal form, then  $M$  and the  $q_j$ 's and  $c_m^j$ 's are uniquely determined by  $f$ .

**Proof.** Since a linear combination of asymptotic expansions of generalized Poincaré type w.r.t. a fixed asymptotic sequence is again an asymptotic expansion of generalized Poincaré type w.r.t. this asymptotic sequence it is enough to consider the case where  $f = 0$ . Assume therefore that we for all  $N \geq 0$  have

$$0 = \left[ \sum_{j=1}^M e^{2\pi i k q_j} \sum_{m=0}^N c_m^j \varphi_m(k) \right] + o(\varphi_N(k)) \quad (27)$$

in the limit  $k \rightarrow \infty$ , where the  $q_j$ 's are pairwise different numbers in  $[0, 1[$  and  $c_m^j \in \mathbb{C}$ . By multiplying by  $e^{-2\pi i k q_1}$  we can assume that  $q_1 = 0$  and  $q_j \in ]0, 1[$ ,  $j = 2, 3, \dots, M$ . We have to show that all the  $c_m^j$ 's are zero. If  $M = 1$  this trivially follows from Lemma 2.1, so we can assume that  $M > 1$ . By putting  $N = 0$  we see that  $g(k) := c_1 + \sum_{j=2}^M c_j e^{2\pi i k q_j}$  converges to zero as  $k \rightarrow \infty$ , where we have put  $c_j = c_0^j$ . Thus  $|g(k)|^2 \rightarrow 0$  as  $k \rightarrow \infty$  implying that  $\frac{1}{L} \sum_{k=0}^{L-1} |g(k)|^2$  converges to zero as  $L \rightarrow \infty$ . Here

$$|g(k)|^2 = |c_1|^2 + |c_2|^2 + \dots + |c_M|^2 + \sum_{\substack{i,j \in \{1, \dots, M\} \\ i \neq j}} c_i \bar{c}_j e^{2\pi i k (q_i - q_j)},$$

and therefore

$$\frac{1}{L} \sum_{k=0}^{L-1} |g(k)|^2 = |c_1|^2 + |c_2|^2 + \dots + |c_M|^2 + \frac{1}{L} \sum_{\substack{i,j \in \{1, \dots, M\} \\ i \neq j}} c_i \bar{c}_j \frac{1 - e^{2\pi i L (q_i - q_j)}}{1 - e^{2\pi i (q_i - q_j)}}$$

so  $\frac{1}{L} \sum_{k=0}^{L-1} |g(k)|^2$  converges to  $|c_1|^2 + |c_2|^2 + \dots + |c_M|^2$  as  $L \rightarrow \infty$ , hence  $c_j = 0$  for all  $j$ .



The proof is finalized by induction: Assume that  $c_m^j = 0$  for all  $j$  and all  $m = 0, 1, \dots, N-1$  and get from (27) that  $c_N^1 + \sum_{j=2}^M c_N^j e^{2\pi i k q_j}$  converges to zero as  $k \rightarrow \infty$ . As before we then have  $c_N^j = 0$  for all  $j$ .  $\square$

Let us see how the above fits into the theory of quantum invariants. Therefore, let  $f : A \rightarrow \mathbb{C}$  and assume that there exist constants  $q_j \in [0, 1[$ ,  $D_j \in \mathbb{Q}$  and  $b_j \in \mathbb{C}$  for  $j = 1, 2, \dots, n$ , and  $a_m^j \in \mathbb{C}$  for  $j = 1, 2, \dots, n$  and  $m = 1, 2, \dots$  such that we for all  $N \in \{0, 1, 2, \dots\}$  have

$$f(k) = \sum_{j=1}^M b_j e^{2\pi i k q_j} k^{D_j} \left( 1 + \sum_{m=1}^N a_m^j k^{-\theta_m} \right) + o(k^{D-\theta_N}) \quad (28)$$

in the limit  $k \rightarrow \infty$ , where  $D = \max\{D_1, \dots, D_M\}$  and  $\{\theta_m\}$  is a strictly increasing sequence in  $\frac{1}{b}\mathbb{Z}_{>0}$ , where  $b$  is the least positive integer such that  $bD_j \in \mathbb{Z}$  for all  $j$ , compare with Conjecture 1.1. By adding terms of the form  $0k^{-l/b}$  we can assume that  $\{\theta_m\}_{m=1}^\infty = \frac{1}{b}\mathbb{Z}_{>0}$ . It is straightforward to see then that we for any  $N \in \{0, 1, 2, \dots\}$  have

$$f(k) = \sum_{j=1}^M e^{2\pi i k q_j} \sum_{m=0}^N c_m^j k^{\lambda_m} + o(k^{\lambda_N}) \quad (29)$$

in the limit  $k \rightarrow \infty$ , where  $c_m^j = 0$  for  $m = 0, 1, \dots, e_j - 1$ ,  $e_j = b(D - D_j)$ , and  $c_m^j = b_j a_{m-e_j}^j$  for  $m = e_j, e_j + 1, \dots$ , and  $\lambda_m = D - \theta_m$ ,  $m = 0, 1, \dots$ . Here we have put  $a_0^j = 1$  and  $\theta_0 = 0$ . Thus  $f$  has an asymptotic expansion of generalized power series type with  $\xi = \text{id}$  and  $\{\lambda_m\}$  a strictly decreasing sequence equal to  $\frac{1}{b}\mathbb{Z}_{\leq bD}$  as a set.

If oppositely  $f : A \rightarrow \mathbb{C}$  has an asymptotic expansion like in (29) with  $\{\lambda_m\}$  a strictly decreasing sequence equal to  $\frac{1}{b}\mathbb{Z}_{\leq a}$  as a set,  $a \in \mathbb{Z}$  and  $b$  some positive integer, then it is straightforward to bring the expansion on a form like in (28). Simply put  $D_j = \lambda_{e_j}$ , where

$$e_j = \min\{m \in \mathbb{Z}_{\geq 0} \mid c_m^j \neq 0\}$$

and  $b_j = c_{e_j}^j$ ,  $a_m^j = c_{m+e_j}^j / b_j$  and  $\theta_m = D - \lambda_{m+e_j}$ , where  $e = \min\{e_1, \dots, e_M\}$ . (We assume here that there are no  $j$ 's with  $c_m^j = 0$  for all  $m$ . If  $c_m^j = 0$  for all  $m$ , simply ignore that term.)

Because of the above, we will pay special attention to the following asymptotic expansions of generalized power series type.

**Definition 2.4.** A function  $f : A \rightarrow \mathbb{C}$  is said to have an *asymptotic expansion of quantum type* w.r.t.  $b \in \mathbb{N}$  or a quantum  $b$ -expansion for short if  $f$  has an asymptotic expansion of generalized power series type as in Definition 2.2 with  $\xi = \text{id}$  and  $\{\lambda_m\} \subseteq \frac{1}{b}\mathbb{Z}$ . The expansion is said to be on minimal form if the expansion is on minimal form as an asymptotic expansion of generalized power series type.

We can assume w.l.o.g. that  $\{\lambda_m\}_{m=0}^\infty = \frac{1}{b}\mathbb{Z}_{\leq a}$ , where  $a = \lambda_0 b$ . Moreover, it is obvious that a quantum  $b$ -expansion is also a quantum  $b'$ -expansion for any  $b'$  divisible by  $b$ . We note that the set  $\mathcal{Q}$  of functions  $f : A \rightarrow \mathbb{C}$  having an asymptotic expansion of quantum type is a subalgebra of the  $\mathbb{C}$ -algebra of all complex functions on  $A$ : It is obvious that a scalar times a  $f \in \mathcal{Q}$  is again in  $\mathcal{Q}$ . If  $f_i \in \mathcal{Q}$  has a quantum  $b_i$ -expansion,  $i = 1, 2$ , then both  $f_1$  and  $f_2$  have a quantum  $b$ -expansion, where  $b = \text{lcm}(b_1, b_2)$ . Now the sum and product of quantum  $b$ -expansions are again quantum  $b$ -expansions, so  $f_1 + f_2$  and  $f_1 f_2$  are both in  $\mathcal{Q}$ .

We can now restate the AEC, Conjecture 1.1, in the following form:

**Conjecture 2.5.** *For any 3-manifold  $X$  the function  $r \mapsto \tau_r^G(X)$ ,  $\mathbb{Z}_{\geq h^\vee} \rightarrow \mathbb{C}$ , has an asymptotic expansion of quantum type as in (29) with  $\{q_j\}_{j=1}^M$  being the image set of the Chern–Simons functional on the moduli space of flat  $G$ -connections on  $X$ .*

By combining the AEC and Conjecture 1.3 we get the stronger conjecture that  $r \mapsto \tau_r^G(X)$  has a quantum 2-expansion. Note that the fact that  $\mathcal{Q}$  is multiplicative stable is well in accordance with the AEC and the fact that the RT-invariants behave multiplicatively w.r.t. connected sums, see e.g. [Tu, Chap. II].

We end this section by giving miscellaneous facts about asymptotic expansions of quantum type. First we make the trivial remark that not every function  $f : A \rightarrow \mathbb{C}$  has an asymptotic expansion of quantum type. If namely  $f$  has such an expansion there exists a positive constant  $C$  and a rational number  $\lambda$  such that

$$|f(k)| \leq Ck^\lambda$$

for all  $k \in A$ . Thus it is easy to give examples of classes of functions which do not have an asymptotic expansion of quantum type. One such class is given by  $f(k) = P(k)e^{Q(k)}$ , where  $P(k)$  is a Laurent polynomial in  $k$  (different from zero), and  $Q(k)$  is a polynomial in  $k$  of degree  $\geq 1$  with positive leading coefficient.

Theorem 2.3 shows that a function having an asymptotic expansion of quantum type determines the quantities building up the expansion (if the expansion is on minimal form). The opposite is obviously not the case. That is, two different functions  $f, g : A \rightarrow \mathbb{C}$  can have the same asymptotic expansion of quantum type. If namely  $f$  has an expansion like in (29), if  $g = f + h$ , and if  $h = o(k^{\lambda_m})$  as  $k \rightarrow \infty$  for all  $m$ , then  $g$  has the same asymptotic expansion of quantum type as  $f$ . As an example one can let  $h(k) = \exp(\beta k)$  with  $\beta$  a negative real number or more generally  $h(k) = P(k)\exp(Q(k))$ , where  $P(k)$  is a (non-zero) Laurent polynomial and  $Q(k)$  is a polynomial of degree  $\geq 1$  with negative leading coefficient. In particular, such a function  $h$  has a trivial asymptotic expansion of quantum type (without being zero).

Consider a function  $g : A \rightarrow \mathbb{C}$  given by an expression of the form

$$g(k) = \sum_{j=1}^M e^{2\pi i k q_j} k^{D_j} b_j g_j(k), \quad (30)$$

where  $b_j \in \mathbb{C} \setminus \{0\}$ ,  $D_j \in \mathbb{Q}$  and  $q_j \in [0, 1[$ , and  $g_j(k)$  are polynomials in  $k^{-1/b}$  with constant term 1, where  $b$  is the least positive integer such that  $bD_j \in \mathbb{Z}$  for all  $j$ . Then  $f$  has obviously an asymptotic expansion of quantum type. More generally this is true if the functions  $g_j$  satisfy asymptotic identities

$$g_j(k) \sim \sum_{m=0}^{\infty} a_m^j k^{-m/b}; \quad \{k^{-m/b}\}, \quad k \rightarrow \infty \quad (31)$$

with  $a_0^j = 1$  for all  $j$ . A particular example of the above is given by the case where

$$g_j(k) \sim \sum_{m=0}^{\infty} \tilde{a}_m^j k^{-m}; \quad \{k^{-m}\}, \quad k \rightarrow \infty.$$

If we put  $a_{bm}^j = \tilde{a}_m^j$  for  $m \in \mathbb{Z}_{\geq 0}$  and  $a_m^j = 0$  for  $m$  not divisible by  $b$  we get (31).

For any function  $f : A \rightarrow \mathbb{C}$  having an asymptotic expansion of quantum type there exists a function  $g : A \rightarrow \mathbb{C}$  given as in (30) and satisfying that  $f$  and  $g$  have the same asymptotic expansion of quantum type. In fact, assume that the expansion of  $f$  is non-trivial and assume it is on minimal form (to avoid terms with  $c_m^j = 0$  for all  $m$ ). By our comments in connection to (29) we can find constants  $b_j$ ,  $D_j$  and  $a_m^j$  and a sequence  $\{\theta_m\}$  such that (28) is satisfied.

According to [O, Sec. 1.9] we can find functions  $g_j : A \rightarrow \mathbb{C}$  such that (31) is satisfied. (In fact, if  $0 < a < 1$ ,  $0 < \sigma < \pi$  and  $\mathbb{S} = \{z \in \mathbb{C} : |z| \geq a, |\text{Arg}(z)| \leq \sigma\}$  then we can find analytic functions  $f_j : \mathbb{S} \rightarrow \mathbb{C}$  such that

$$f_j(z) \sim \sum_{m=0}^{\infty} a_m^j z^{-m}; \quad \{z^{-m}\}, \quad z \rightarrow \infty \text{ in } \mathbb{S}.$$

By letting  $g_j(k) = f_j(k^{1/b})$ ,  $k \in \mathbb{N}$ , we get the desired functions.) Now, if we let  $g : A \rightarrow \mathbb{C}$  be given by (30) with the  $b_j$ 's,  $D_j$ 's and the  $q_j$ 's coming from the expansion (28) of  $f$ , then  $g$  has an asymptotic expansion of quantum type equal to that of  $f$ . The functions  $g_j$  are of course not uniquely determined by the numbers  $a_m^j$ . We see that the function  $g$  can even be chosen to be analytic in any sector shaped region in the complex plane containing the positive integers. However, from an asymptotic point of view we do of course not gain anything. The asymptotic behaviour is solely determined by the coefficients  $c_m^j$ , the numbers  $q_j$  and the sequence  $\{\lambda_m\}$ .

Assume  $f : A \rightarrow \mathbb{C}$  has a non-trivial asymptotic expansion of quantum type as in (28), and let  $J = \{j \in \{1, \dots, M\} \mid d_j = D\}$ . Then

$$\sum_{j \in J} b_j e^{2\pi i k q_j} k^D \quad (32)$$

is called the leading term of the asymptotic expansion of  $f$ . For a 3-manifold  $X$  with discrete moduli space of flat  $G$ -connections the semiclassical approximation of  $Z_k^G(X)$ , i.e. the right-hand side of (4), is (conjecturally) the leading order large  $k$  asymptotics of  $Z_k^G(X)$  if and only if all the  $D_A$ 's are equal. In general the leading asymptotics is given by the sum of terms with maximal  $D_A$ .

### 3. THE RESHETIKHIN–TURAEV INVARIANTS OF SEIFERT MANIFOLDS FOR $\text{SU}(2)$

We use the notation introduced by Seifert in his classification results for the Seifert manifolds (or rather fibrations), see [Se1], [Se2], [Ha2, Sect. 2]. That is,  $(\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  is the Seifert manifold with orientable base of genus  $g \geq 0$  if  $\epsilon = o$  and non-orientable base of genus  $g > 0$  if  $\epsilon = n$  (where the genus of the non-orientable connected sum  $\#^k \mathbb{RP}^2$  is  $k$ ). (In [Se1], [Se2]  $(\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  is denoted  $(O, \epsilon; g \mid b; \alpha_1, \beta_1; \dots; \alpha_n, \beta_n)$  but we leave out the  $O$ , since we are only dealing with oriented Seifert manifolds.) The pair  $(\alpha_j, \beta_j)$  of coprime integers is the oriented Seifert invariant of the  $j$ 'th exceptional (or singular) fiber. We have  $0 < \beta_j < \alpha_j$ . The integer  $-b$  is equal to the Euler number of the Seifert fibration  $(\epsilon; g \mid b)$  (which is a locally trivial  $S^1$ -bundle). The sign is chosen so that the Euler number of the spherical (or unit) tangent bundle over an orientable surface  $\Sigma$  is equal to the Euler characteristic of  $\Sigma$ , see [M, Chap. 1 and 4], [Sc, Sec. 3]. More generally, the Seifert Euler number of  $(\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  is  $E = -\left(b + \sum_{j=1}^n \beta_j / \alpha_j\right)$ . We note that except for a small class of Seifert manifolds, the Seifert invariants are actually topological invariants, i.e. they classify the Seifert manifolds up to orientation preserving homeomorphisms, see e.g. [Or, Chap. 5 Theorem 6 p. 97 and Sec. 5.4] or [JN, Theorem 5.1 p. 32]. The exceptions are the lens spaces, the prism manifolds  $(o; 0 \mid b; (2, 1), (2, 1), (\alpha, \beta))$ , and the manifolds  $(n; 2 \mid 0)$ ,  $(o; 0 \mid -2; (2, 1), (2, 1), (2, 1), (2, 1))$ ,  $(n; 1 \mid b)$  with  $b \neq 0$  and  $(n; 1 \mid b; (\alpha, \beta))$ . These exceptions constitute a proper subclass of the 'small Seifert manifolds' listed in [Or, pp 91-92]. The lens spaces are the Seifert manifolds with  $\epsilon = o$ ,  $g = 0$  and  $n \leq 2$ . A Seifert manifold which is not small is sometimes called a large Seifert manifold, see [Or, p. 92].

For the convenience of the reader we will state all results in the following both in terms of the Seifert invariants and in terms of the so-called non-normalized Seifert invariants due to W. D. Neumann. For a Seifert manifold  $X$  with non-normalized Seifert invariants  $\{\epsilon; g; (\alpha_1, \beta_1), \dots,$

$(\alpha_n, \beta_n)\}$  the invariants  $\epsilon$  and  $g$  are as above. The  $(\alpha_j, \beta_j)$ 's are here pairs of coprime integers with  $\alpha_j > 0$  but not necessarily with  $0 < \beta_j < \alpha_j$ . These pairs are actually not invariants of the fibration  $X$ . Thus  $\{\epsilon; g; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\}$  and  $\{\epsilon; g; (\alpha'_1, \beta'_1), \dots, (\alpha'_m, \beta'_m)\}$  are two sets of non-normalized Seifert invariants of  $X$  if and only if the set of pairs  $(\alpha'_j, \beta'_j)$  can be obtained from the set of pairs  $(\alpha_i, \beta_i)$  by a finite number of the following two moves: i) add or delete a pair  $(1, 0)$ , ii) replace each  $(\alpha_i, \beta_i)$  by  $(\alpha_i, \beta_i + K_i \alpha_i)$  provided  $\sum K_i = 0$ . For details, see [JN]. The Seifert Euler number of  $X$  is  $-\sum_{i=1}^n \beta_i / \alpha_i$  (which is an invariant of the Seifert fibration  $X$ ). The main advantage obtained by working with the non-normalized invariants is to make more 'symmetric' expressions, since the invariant  $b$  is treated formally as a Seifert invariant  $(1, b)$  of an exceptional fiber. Thus the Seifert manifold  $(\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  is equal to  $\{\epsilon; g; (1, b), (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\}$ . Of course the number of exceptional fibers is a constant for a Seifert manifold. This number can be read off directly from the (normalized) Seifert invariants, but can also be read off from a set of non-normalized invariants as the number of pairs  $(\alpha_j, \beta_j)$  with  $\alpha_j > 1$ .

In [JN] the authors work with a more general class of oriented fibered spaces, denoted Generalized Seifert fibrations. According to [JN, Theorem 5.1] these spaces are (up to orientation preserving homeomorphism) the classical oriented Seifert manifolds as considered here and connected sums of lens spaces (considering  $S^1 \times S^2$  as a lens space). Since the RT-invariant of a connected sum of 3-manifolds is the product of the RT-invariants of the 3-manifolds in that connected sum (up to a normalization factor), the results of [J1] actually show the AEC for these connected sums of lens spaces and  $G = \text{SU}(2)$ .

To state the next theorem we need some notation. For a pair of coprime integers  $c, d$  with  $|c| \geq 1$  the Dedekind sum is given by

$$s(d, c) = \frac{1}{4|c|} \sum_{j=1}^{|c|-1} \cot \frac{\pi j}{c} \cot \frac{\pi d j}{c} \quad (33)$$

for  $|c| > 1$  and  $s(d, \pm 1) = 0$ . We refer to [RG] for a comprehensive description of this function. The Dedekind symbol is given by

$$S(d/c) = 12 \text{sign}(c) s(d, c). \quad (34)$$

Multi-indices are denoted by an underline (e.g.  $\underline{n}$ ). For  $\underline{k} = (k_1, \dots, k_n), \underline{l} = (l_1, \dots, l_n) \in \mathbb{Z}^n$ ,  $\underline{k} < \underline{l}$  if and only if  $k_j < l_j$  for all  $j = 1, \dots, n$ . We let  $\underline{1} = (1, \dots, 1)$ . For  $\underline{k} = (k_1, \dots, k_n) \in \mathbb{Z}_+^n$  we write  $\sum_{\underline{m}=\underline{0}}^{\underline{k}}$  for  $\sum_{m_1=0}^{k_1} \dots \sum_{m_n=0}^{k_n}$  etc. Let  $a_o = 2$  and  $a_n = 1$ . Given pairs of coprime integers  $\alpha_j, \beta_j$  we choose integers  $\rho_j, \sigma_j$  such that  $\alpha_j \sigma_j - \beta_j \rho_j = 1$ .

**Theorem 3.1** ([Ha2, Theorem 8.4]). *The RT-invariant at level  $r-2$  of the Seifert manifold  $X$  with (normalized) Seifert invariants  $(\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  is*

$$\begin{aligned} \tau_r(X) &= (-1)^{a_\epsilon g} \frac{r^{a_\epsilon g/2-1}}{2^{a_\epsilon g/2-1}} \frac{1}{\sqrt{\mathcal{A}}} e^{i \frac{3\pi}{4} (1-a_\epsilon) \text{sign}(E)} \\ &\quad \times \exp \left( \frac{i\pi}{2r} \left[ 3(a_\epsilon - 1) \text{sign}(E) - E - \sum_{j=1}^n S \left( \frac{\beta_j}{\alpha_j} \right) \right] \right) Z(X; r), \end{aligned} \quad (35)$$

where  $\mathcal{A} = \prod_{j=1}^n \alpha_j$  and

$$Z(X; r) = \left( \frac{i}{2} \right)^n \sum_{\gamma=1}^{r-1} \frac{(-1)^{\gamma a_\epsilon g} h(\gamma)}{\sin^{n+a_\epsilon g-2} \left( \frac{\pi}{r} \gamma \right)}, \quad (36)$$

where

$$h(\gamma) = \exp\left(\frac{i\pi}{2r}E\gamma^2\right) \sum_{\mu \in \{\pm 1\}^n} \sum_{\underline{n}=0}^{\underline{\alpha}-1} \left( \prod_{j=1}^n \mu_j \right) \quad (37)$$

$$\times \exp\left(2\pi i \sum_{j=1}^n \frac{\rho_j}{\alpha_j} [rn_j^2 + \mu_j n_j]\right) \exp\left(-\frac{i\pi}{r}\gamma \sum_{j=1}^n \frac{2rn_j + \mu_j}{\alpha_j}\right).$$

The RT-invariant at level  $r-2$  of the Seifert manifold  $X$  with non-normalized Seifert invariants  $\{\epsilon; g; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\}$  is given by the same expression.

The theorem is also valid in case  $n = 0$ , where one has to put  $\sum_{j=1}^n S(\beta_j/\alpha_j) = 0$  and  $\mathcal{A} = 1$  in (35) and let  $h(\gamma) = \exp\left(\frac{i\pi}{2r}E\gamma^2\right)$  in (36). The reason for including the factor  $(i/2)^n$  in the expression for  $Z(X; r)$  is that then  $Z(X; r)$  does not depend on the choice of non-normalized Seifert invariants for  $X$ . In fact,  $Z(X; r)$  only depends on the  $\beta_j$ 's through the Seifert Euler number  $E$ , and if we add a pair  $(\alpha, \beta) = (1, 0)$  each of the terms in the sum-expression for  $Z(X; r)$  changes by a factor  $-2i$ , hence  $Z(X; r)$  does not change.

#### 4. THE ANALYTIC ASYMPTOTIC EXPANSIONS OF THE RESHETIKIN-TURAEV INVARIANTS OF SEIFERT MANIFOLDS

Let  $X$  be a Seifert manifold with (normalized) Seifert invariants  $(\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  or non-normalized Seifert invariants  $\{\epsilon; g; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\}$  with  $\epsilon = 0$  or  $g$  even. Let  $\beta_0 = b$  if  $X$  is given by normalized Seifert invariants and  $\beta_0 = 0$  otherwise. We will below present detailed expressions for the asymptotic expansions of the RT-invariants of  $X$ . It will follow that all parts of the asymptotics are expressible by the Seifert invariants.

By the Theorems 4.1 and 4.4 below,  $Z(X; r)$  has an asymptotic expansion of the form (1), and according to Theorem 3.1

$$\tau_r(X) = br^{a_\epsilon g/2-1} \exp\left(\frac{ia}{r}\right) Z(X; r),$$

where  $a \in \mathbb{R}$  and  $b \in \mathbb{C} \setminus \{0\}$ , so  $\tau_r(X)$  has also an asymptotic expansion of the form (1).

Before stating Theorems 4.1 and 4.4 we need some preliminaries. We treat the case  $n = 0$  as the case  $n = 1$  with  $(\alpha_1, \beta_1) = (1, 0)$ . For  $E \neq 0$  we let  $z_{\text{st}} : \mathbb{Z} \times \mathbb{Z}^n \rightarrow \mathbb{R}$  be given by

$$z_{\text{st}}(m, \underline{n}) = -\frac{2}{E} \left( m - \sum_{j=1}^n \frac{n_j}{\alpha_j} \right). \quad (38)$$

Moreover, let

$$q(m, \underline{n}) = \sum_{j=1}^n \frac{\rho_j}{\alpha_j} n_j^2 - \frac{1}{4} E z_{\text{st}}(m, \underline{n})^2 \pmod{\mathbb{Z}}, \quad (m, \underline{n}) \in \mathbb{Z} \times \mathbb{Z}^n, \quad (39)$$

$$q(l, \underline{n}') = -\frac{1}{4} \beta_0 l^2 + \sum_{j=1}^n \left( \frac{\rho_j}{\alpha_j} n_j'^2 - \frac{1}{4} \sigma_j \beta_j l^2 \right) \pmod{\mathbb{Z}}, \quad (l, \underline{n}') \in M, \quad (40)$$

where  $M = \{ (l, \underline{n}') \in \mathbb{Z} \times (\frac{1}{2}\mathbb{Z})^n \mid \underline{n}' \in \mathbb{Z}^n + \frac{1}{2}l\underline{\beta} \}$ . We let

$$\mathcal{S} = \{ \underline{n} \in \mathbb{Z}^n \mid \underline{0} \leq \underline{n} \leq \underline{\alpha} - \underline{1} \} \quad (41)$$

and

$$\mathcal{I}_1 = \{ (m, \underline{n}) \in \mathbb{Z} \times \mathcal{S} \mid 0 < z_{\text{st}}(m, \underline{n}) < 1 \}, \quad (42)$$

$$\mathcal{I}_2 = \left\{ (l, \underline{n}') \in M \mid l \in \{0, 1\}, 0 \leq \underline{n}' \leq \frac{1}{2}\underline{\alpha} \right\}. \quad (43)$$

Above  $q_{(m,n)}$  and  $\mathcal{I}_1$  are of course only defined for  $E \neq 0$ . We need a partition

$$\mathcal{I}_2^a = \left\{ (l, \underline{n}') \in \mathcal{I}_2 \mid \exists \underline{\mu}' \in \{\pm 1\}^n : \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \in \mathbb{Z} \right\}$$

and  $\mathcal{I}_2^b = \mathcal{I}_2 \setminus \mathcal{I}_2^a$  of  $\mathcal{I}_2$ . We will need to write 1 and 2 in a sophisticated way, namely let

$$\text{Sym}_{\pm}(x) = \begin{cases} 1 & , x \neq 0, \\ 2 & , x = 0, \end{cases} \quad (44)$$

and

$$\text{Sym}_{\mathbb{Z}_{\pm}}(x) = \begin{cases} 1 & , x \in \mathbb{R} \setminus \frac{1}{2}\mathbb{Z}, \\ 2 & , x \in \frac{1}{2}\mathbb{Z}. \end{cases} \quad (45)$$

These functions have a group theoretical explanation, see [Ro3, p. 36]. However, this can be neglected here. Basically these functions are used to keep track of how many times certain terms contribute to certain sums being part of the asymptotic expansion of  $Z(X; r)$ .

As already indicated above we have to consider the cases  $E = 0$  and  $E \neq 0$  separately. Theorem 4.1 concerns the case  $E \neq 0$  while Theorem 4.4 takes care of the case  $E = 0$ .

**Theorem 4.1.** *Assume the Seifert Euler number  $E \neq 0$ . Then*

$$Z(X, r) = Z_{\text{polar}}(X; r) + Z_{\text{int}}(X; r),$$

where  $Z_{\text{polar}}(X; r)$  is a sum of residues, while the other term  $Z_{\text{int}}(X; r)$  is given by certain limits of sums of certain integrals along contours in the complex plane. If  $n + a_{\epsilon}g - 2 \leq 0$  then  $Z_{\text{polar}}(X; r) = 0$ . For  $n + a_{\epsilon}g - 2 > 0$  we have

$$Z_{\text{polar}}(X; r) = \sum_{(l, \underline{n}') \in \mathcal{I}_2} b_{(l, \underline{n}')} \exp(2\pi i r q_{(l, \underline{n}')} r) \left( Z_0^{(l, \underline{n}')} (r) + Z_1^{(l, \underline{n}')} (r) \right), \quad (46)$$

where

$$b_{(l, \underline{n}')} = \frac{2^n (-1)^{(n + \sum_{j=1}^n \sigma_j)l}}{\prod_{j=1}^n \text{Sym}_{\mathbb{Z}_{\pm}}\left(\frac{n'_j}{\alpha_j}\right)}, \quad (47)$$

$q_{(l, \underline{n}')}$  is given by (40), and the functions  $Z_0^{(l, \underline{n}')}$  and  $Z_1^{(l, \underline{n}')}$  are given by

$$\begin{aligned} Z_0^{(l, \underline{n}')} (r) &= -\frac{\pi}{2} \text{Res}_{z=0} \left\{ \frac{\exp\left(\frac{i\pi r}{2} E z^2\right)}{\sin^{n+a_{\epsilon}g-2}(\pi z)} \cot(\pi r z) \right. \\ &\quad \times \prod_{j=1}^n \left[ i \sin\left(2\pi \frac{\rho_j}{\alpha_j} n'_j\right) \cos\left(\pi \frac{z}{\alpha_j}\right) \sin\left(2\pi r z \frac{n'_j}{\alpha_j}\right) \right. \\ &\quad \left. \left. + \cos\left(2\pi \frac{\rho_j}{\alpha_j} n'_j\right) \sin\left(\pi \frac{z}{\alpha_j}\right) \cos\left(2\pi r z \frac{n'_j}{\alpha_j}\right) \right] \right\} \end{aligned} \quad (48)$$

and

$$\begin{aligned}
Z_1^{(l, \underline{n}')} (r) &= -\frac{\pi i}{(-2)^n} \sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |a|}} \frac{\text{sign}(a)}{\text{Sym}_{\pm}(m) \text{Sym}_{\pm}(m - |a|)} \\
&\times \text{Res}_{z=0} \left\{ \frac{\exp \left[ \frac{i\pi r}{2} E z^2 + 2\pi i r \text{sign}(a) (m - |a|) z \right]}{\sin^{n+a_{\epsilon}g-2}(\pi z)} \right. \\
&\times \left. \prod_{j=1}^n \mu'_j \sin \left[ 2\pi \left( \frac{\rho_j}{\alpha_j} n'_j - \frac{z}{2} \frac{\mu'_j}{\alpha_j} \right) \right] \right\},
\end{aligned}$$

where  $a = a(\underline{\mu}', \underline{n}') = \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j}$ . In particular,  $Z_0^{(l, \underline{n}')} (r) = 0$  if  $g = 0$ .

Let  $\mathcal{I} = \mathcal{I}_1 \cup \mathcal{I}_2^a$ , and let  $k_0 = 0$  if  $n$  is even and  $k_0 = 1$  otherwise. Moreover, let  $k_2 = \min\{0, k_1\}$ , where  $k_1 = (k_0 - n - a_{\epsilon}g + 2)/2$ . Then

$$r^{-1/2} Z_{\text{int}}(X; r) \sim \sum_{k=k_2}^{\infty} \left( \sum_{\gamma \in \mathcal{I}} \exp(2\pi i r q_{\gamma}) c_k^{\gamma} \right) r^{-k}; \quad \{r^{-k}\}, \quad r \rightarrow \infty,$$

where the quantities  $q_{\gamma}$  and  $c_k^{\gamma}$  are given as follows: For  $\gamma = (m, \underline{n}) \in \mathcal{I}_1$ ,  $q_{\gamma}$  is given by (39). Moreover,

$$c_k^{\gamma} = (-1)^n e^{\frac{i\pi}{4} \text{sign}(E)} \sqrt{\frac{2}{|E|}} \frac{1}{k!} \left( \frac{i}{2\pi E} \right)^k \partial_z^{(2k)} \frac{\prod_{j=1}^n \sin \left( \frac{\pi}{\alpha_j} (2\rho_j n_j - z) \right)}{\sin^{n+a_{\epsilon}g-2}(\pi z)} \Big|_{z=z_{\text{st}}(m, \underline{n})} \quad (49)$$

for  $k \geq 0$  and  $c_k^{\gamma} = 0$  for  $k < 0$  (in case  $k_2 < 0$ ). Here  $z_{\text{st}}(m, \underline{n})$  is given by (38). For  $\gamma = (l, \underline{n}') \in \mathcal{I}_2^a$ ,  $q_{\gamma}$  is given by (40). For  $k \in \{k_1, k_1 + 1, \dots\}$  we put  $k' = 2k + n + a_{\epsilon}g - 2$  and have

$$c_k^{\gamma} = \frac{b(l, \underline{n}')}{(-2)^n \pi^{n+a_{\epsilon}g-2}} \frac{e^{\frac{i\pi}{4} \text{sign}(E)} \Gamma(k + \frac{1}{2})}{\sqrt{2\pi|E|} k!} \left( \frac{2i}{\pi E} \right)^k \sum_{\substack{\underline{\mu}' \in \{\pm 1\}^n : \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \in \mathbb{Z}}} \partial_z^{(k')} f(z; l, \underline{n}', \underline{\mu}') \Big|_{z=0}, \quad (50)$$

where  $b(l, \underline{n}')$  is given by (47) and

$$f(z; l, \underline{n}', \underline{\mu}') = \left( \frac{\pi z}{\sin(\pi z)} \right)^{n+a_{\epsilon}g-2} \prod_{j=1}^n \sin \left( \pi \frac{2\rho_j \mu'_j n'_j - z}{\alpha_j} \right).$$

We note that  $k_1 \leq 1$ , and  $k_1 = 1$  if and only if  $n = k_0 = 1$ ,  $\epsilon = 0$  and  $g = 0$ . In that case  $c_0^{\gamma} = 0$ .

We have in the above theorem focused on the asymptotic expansion of  $Z(X; r)$  and have therefore chosen not to give an explicit expression for the term  $Z_{\text{int}}(X; r)$ . Such an explicit expression can be found in the proof of Theorem 4.1, see (107) together with (100) and (102), but is not relevant here and is not very informative.

**Remark 4.2.** The above theorem is also true in case  $n = 0$  if we as usual put all products  $\prod_{j=1}^n$  equal to 1 and all sums  $\sum_{j=1}^n$  equal to 0 and if we make the following natural adjustments: The

function  $z_{\text{st}}$  in (38) and the  $q$ -function in (39) only depend on  $m \in \mathbb{Z}$ , namely

$$\begin{aligned} z_{\text{st}}(m) &= -\frac{2m}{E}, \\ q_m &= -\frac{m^2}{E} \pmod{\mathbb{Z}}. \end{aligned}$$

The index sets  $\mathcal{I}_1$ ,  $\mathcal{I}_2$  and  $\mathcal{I}_2^a$  become

$$\begin{aligned} \mathcal{I}_1 &= \{ m \in \mathbb{Z} \mid 0 < z_{\text{st}}(m) < 1 \}, \\ \mathcal{I}_2 &= \mathcal{I}_2^a = \{0, 1\}. \end{aligned}$$

The  $q$ -function in (40) degenerates to  $q_l = -\frac{1}{4}\beta_0 l^2 \pmod{\mathbb{Z}}$ ,  $l = 0, 1$ . The sum

$$\sum_{\underline{\mu}' \in \{\pm 1\}^n : \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \in \mathbb{Z}} \partial_z^{(k')} f(z; l, \underline{n}', \underline{\mu}') \Big|_{z=0}$$

in (50) simply becomes

$$\partial_z^{(k')} \left( \frac{\pi z}{\sin(\pi z)} \right)^{a_\epsilon g - 2} \Big|_{z=0}.$$

The coefficients  $b_{(l, \underline{n}')}$  in (47) become  $b_l = 1$ ,  $l = 0, 1$ . Finally the functions  $Z_\nu^{(l, \underline{n}')}$ ,  $\nu = 0, 1$ , degenerate to  $Z_1^l(r) = 0$  and

$$Z_0^l(r) = -\frac{\pi}{2} \text{Res}_{z=0} \left\{ \frac{\exp\left(\frac{i\pi r}{2} E z^2\right)}{\sin^{a_\epsilon g - 2}(\pi z)} \cot(\pi r z) \right\}$$

for  $l \in \mathcal{I}_2$ , both actually independent of  $l$ . Thus

$$Z_{\text{polar}}(X; r) = -\pi r \text{Res}_{z=0} \left\{ \frac{\exp\left(\frac{i\pi r}{2} E z^2\right)}{\sin^{a_\epsilon g - 2}(\pi z)} \cot(\pi r z) \right\}.$$

To verify the above claims recall that we in Theorem 4.1 have treated the case  $n = 0$  formally as the case  $n = 1$  with  $(\alpha_1, \beta_1) = (1, 0)$ . Putting  $(\rho_1, \sigma_1) = (0, 1)$  we immediately get the claims above, see also the proof of Proposition 4.5. The case  $n = 0$  and  $E \neq 0$  only occurs if we work with normalized Seifert invariants, namely for the spaces  $X = (\epsilon; g \mid b)$ ,  $b \neq 0$ . If we work with non-normalized Seifert invariants there will always be a  $(\alpha, \beta)$ -pair of the form  $(1, \beta)$  coming from (modifications of) the pair  $(1, b)$ .

There is a peculiar thing in the above theorem. Namely if  $X$  is described by non-normalized Seifert invariants we can always add as many  $(\alpha, \beta)$ -pairs  $(1, 0)$  as we want. In that way we can always obtain that the criterion  $n + a_\epsilon g - 2 > 0$  is satisfied. We will see that this does not change anything. In fact both terms  $Z_{\text{polar}}(X; r)$  and  $Z_{\text{int}}(X; r)$  are independent of the choice of non-normalized Seifert invariants for  $X$ , see Proposition 4.5 for the precise statement.

Before considering the case  $E = 0$  let us take a more careful look at  $Z_{\text{polar}}(X; r)$ .

**Proposition 4.3.** *Assume that  $n + a_\epsilon g - 2 > 0$ . For each  $\gamma = (l, \underline{n}') \in \mathcal{I}_2$  and  $\nu \in \{0, 1\}$  the function  $Z_\nu^\gamma(r)$  is zero or is a Laurent polynomial in  $r$  of the form  $\sum_{k=-\infty}^{n+a_\epsilon g-3} a_k^\gamma r^k$ , where the coefficients  $a_k^\gamma = 0$  for all but finitely many  $k$ .*

**Proof.** The proof is straightforward. Let  $(l, \underline{n}') \in \mathcal{I}_2$  be fixed and put  $\lambda_j = \sin\left(\pi \frac{2\rho_j n'_j}{\alpha_j}\right)$  and



$\kappa_j = \cos\left(\pi \frac{2\rho_j n'_j}{\alpha_j}\right)$ . We start by analyzing  $Z_0^{(l, \underline{n}')}$ . Put

$$\begin{aligned} \tau(z) &= \prod_{j=1}^n \left[ i \sin\left(2\pi \frac{\rho_j}{\alpha_j} n'_j\right) \cos\left(\pi \frac{z}{\alpha_j}\right) \sin\left(2\pi r z \frac{n'_j}{\alpha_j}\right) \right. \\ &\quad \left. + \cos\left(2\pi \frac{\rho_j}{\alpha_j} n'_j\right) \sin\left(\pi \frac{z}{\alpha_j}\right) \cos\left(2\pi r z \frac{n'_j}{\alpha_j}\right) \right] \end{aligned}$$

and write  $\tau(z) = \sum_{\nu=0}^n i^\nu \sum_{1 \leq j_1 < \dots < j_\nu \leq n} g_{(j_1, \dots, j_\nu)}(z)$ , where

$$\begin{aligned} g_{(j_1, \dots, j_\nu)}(z) &= \prod_{j \in \{j_1, \dots, j_\nu\}} \lambda_j \cos\left(\pi \frac{z}{\alpha_j}\right) \sin\left(2\pi r z \frac{n'_j}{\alpha_j}\right) \\ &\quad \times \prod_{j \in \{1, \dots, n\} \setminus \{j_1, \dots, j_\nu\}} \kappa_j \sin\left(\pi \frac{z}{\alpha_j}\right) \cos\left(2\pi r z \frac{n'_j}{\alpha_j}\right). \end{aligned}$$

Note that  $g_{(j_1, \dots, j_\nu)}$  has a zero in 0 of order  $n$  or is constantly zero. We have

$$g_{(j_1, \dots, j_\nu)}(z) = r^\nu \sum_{k=n}^{\infty} e_k^{(j_1, \dots, j_\nu)}(r) z^k,$$

where  $e_k^{(j_1, \dots, j_\nu)}(r)$  is a real polynomial in  $r$  of order at most  $k - n$ . This description gives the result for  $Z_0^{(l, \underline{n}')} (r)$ . For the analysis of  $Z_1^{(l, \underline{n}')} (r)$  we fix  $\underline{\mu}' \in \{\pm 1\}^n$  and put

$$\sigma(z) = \prod_{j=1}^n \mu'_j \sin\left[2\pi \left(\frac{\rho_j}{\alpha_j} n'_j - \frac{z}{2} \frac{\mu'_j}{\alpha_j}\right)\right] = \prod_{j=1}^n \left[ \mu'_j \lambda_j \cos\left(\pi \frac{z}{\alpha_j}\right) - \kappa_j \sin\left(\pi \frac{z}{\alpha_j}\right) \right]$$

and write  $\sigma(z) = \sum_{\nu=0}^n \sum_{1 \leq j_1 < \dots < j_\nu \leq n} h_{(j_1, \dots, j_\nu)}(z)$ , where

$$h_{(j_1, \dots, j_\nu)}(z) = (-1)^{n-\nu} \prod_{j \in \{j_1, \dots, j_\nu\}} \mu'_j \lambda_j \cos\left(\pi \frac{z}{\alpha_j}\right) \prod_{j \in \{1, \dots, n\} \setminus \{j_1, \dots, j_\nu\}} \kappa_j \sin\left(\pi \frac{z}{\alpha_j}\right).$$

We note that  $h_{(j_1, \dots, j_\nu)}(z)$  is independent of  $r$  and has a zero in 0 of order  $n - \nu$  or is constantly zero.  $\square$

Let  $\lambda_j$  and  $\kappa_j$  be as in the above proof. We note that  $\lambda_j = 0$  if and only if  $n'_j \in \{0, \alpha_j/2\}$  and  $\kappa_j = 0$  if and only if  $n'_j = \alpha_j/4$  (so this happens only in case  $\alpha_j$  is even and  $l$  is odd). If  $\lambda_j = 0$  for a  $j \in \{1, 2, \dots, n\}$  then the highest degree term in both  $Z_0^{(l, \underline{n}')} (r)$  and  $Z_1^{(l, \underline{n}')} (r)$  is of an order strictly less than  $n + a_\epsilon g - 3$ . If  $\alpha_j = 1$  for a  $j \in \{1, 2, \dots, n\}$  we have  $\lambda_j = 0$  for all  $(l, \underline{n}') \in \mathcal{I}_2$ . If  $\lambda_j = 0$  for  $m$  indices  $j \in \{1, 2, \dots, n\}$  then the highest degree term in both  $Z_0^{(l, \underline{n}')} (r)$  and  $Z_1^{(l, \underline{n}')} (r)$  is of an order less than or equal to  $n + a_\epsilon g - 3 - m$ .

In the following theorem the case  $n = 0$  is treated as the case  $n = 1$  with  $(\alpha_1, \beta_1) = (1, 0)$ . For that case, see also the remarks following the theorem.

**Theorem 4.4.** *Assume the Seifert Euler number  $E = 0$ . If  $n + a_\epsilon g - 2 \leq 0$  then  $X = S^2 \times S^1$  (so  $\tau_r(X) = 1$ ). For  $n + a_\epsilon g - 2 > 0$  we have*

$$Z(X; r) = \sum_{(l, \underline{n}') \in \mathcal{I}_2} b_{(l, \underline{n}')} \exp(2\pi i r q_{(l, \underline{n}')} r) \left( Z_0^{(l, \underline{n}')} (r) + \tilde{Z}_1^{(l, \underline{n}')} (r) \right),$$

where  $b_{(l, \underline{n}')}$  is as in (46),  $q_{(l, \underline{n}')}$  is given by (40),  $Z_0^{(l, \underline{n}')}$  is given by (48) with  $E = 0$ , and

$$\begin{aligned} \tilde{Z}_1^{(l, \underline{n}')} (r) &= -\pi i \left( \frac{i}{4} \right)^n \sum_{\underline{\mu} \in \{\pm 1\}^n} \sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |a|}} \left( \prod_{j=1}^n \mu_j \right) \\ &\quad \times \frac{\text{sign}(a)}{\text{Sym}_{\pm}(m) \text{Sym}_{\pm}(m - |a|)} \exp \left( 2\pi i \sum_{j=1}^n \mu_j \frac{\rho_j}{\alpha_j} \mu'_j n'_j \right) \\ &\quad \times \text{Res}_{z=0} \left\{ \frac{\exp[2\pi i r \text{sign}(a)](m - |a|)z}{\sin^{n+a_\epsilon g-2}(\pi z)} \right\}, \end{aligned}$$

where  $a = a(\underline{\mu}, \underline{\mu}', \underline{n}') = \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} + \frac{1}{2r} \sum_{j=1}^n \frac{\mu_j}{\alpha_j}$ . Moreover, if  $\sum_{j=1}^n \frac{1}{\alpha_j} < n + a_\epsilon g - 2$  we have that  $Z(X; r) = Z_{\text{polar}}(X; r)$  for all levels  $r$ , where  $Z_{\text{polar}}(X; r)$  is given by (46) with  $E = 0$ . Finally, in all cases there exists a positive number  $r(\underline{\alpha}, \underline{\beta})$  such that

$$Z(X; r) = Z_{\text{polar}}(X; r) + r Z_{\text{spec}}(X; r)$$

for all  $r \geq r(\underline{\alpha}, \underline{\beta})$ , where  $Z_{\text{polar}}(X; r)$  is given by (46) with  $E = 0$  and

$$\begin{aligned} Z_{\text{spec}}(X; r) &= \pi i \left( \frac{i}{4} \right)^n \sum_{(l, \underline{n}') \in \mathcal{I}_2^a} b_{(l, \underline{n}')} \exp(2\pi i r q_{(l, \underline{n}')} ) \sum_{\substack{\underline{\mu}' \in \{\pm 1\}^n \\ \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \in \mathbb{Z}}} \sum_{\substack{\underline{\mu} \in \{\pm 1\}^n \\ \sum_{j=1}^n \frac{\mu_j}{\alpha_j} < 0}} \left( \prod_{j=1}^n \mu_j \right) \\ &\quad \times \exp \left( 2\pi i \sum_{j=1}^n \mu_j \frac{\rho_j}{\alpha_j} \mu'_j n'_j \right) \text{Res}_{z=0} \left\{ \frac{f(z; \underline{\mu})}{\sin^{n+a_\epsilon g-2}(\pi z)} \right\}, \end{aligned}$$

where

$$f(z; \underline{\mu}) = \begin{cases} \cos \left( \pi \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right) & , n \text{ odd}, \\ -i \sin \left( \pi \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right) & , n \text{ even}. \end{cases}$$

In particular,  $Z_{\text{spec}}(X; r) = 0$  for all  $r \geq r(\underline{\alpha}, \underline{\beta})$  if  $\sum_{j=1}^n \frac{1}{\alpha_j} < n + a_\epsilon g - 2$ .

Let us look at the special case  $n = 0$ , i.e.  $X$  is given by the Seifert invariants  $(\epsilon; g \mid 0)$  or non-normalized Seifert invariants  $\{\epsilon; g \mid \}$ . As stated above this case is treated as the case  $n = 1$  with  $(\alpha_1, \beta_1) = (1, 0)$ . If  $a_\epsilon g \geq 4$  we get from the above theorem that

$$Z(X; r) = Z_{\text{polar}}(X; r) = -\pi r \text{Res}_{z=0} \left\{ \frac{\cot(\pi r z)}{\sin^{a_\epsilon g-2}(\pi z)} \right\}. \quad (51)$$

If  $a_\epsilon g = 2$  we have  $X = (o; 1 \mid 0)$  or  $X = (n; 2 \mid 0)$ . For this case we use that  $r(\underline{\alpha}, \underline{\beta})$  in Theorem 4.4 can be put equal to 2 since  $n = 1$ . Thus we get for all levels  $r$  that

$$Z(X; r) = Z_{\text{polar}}(X; r) + r Z_{\text{spec}}(X; r).$$

Here  $Z_{\text{polar}}(X; r)$  is given by (51), i.e.  $Z_{\text{polar}}(X; r) = -\pi r \text{Res}_{z=0}(\cot(\pi r z)) = -1$ . Moreover  $Z_{\text{spec}}(X; r) = 1$  so  $Z_{\text{polar}}(X; r) = r - 1$ . Note that this infact follows directly from (36). Finally, if  $a_\epsilon g = 0$ , i.e.  $\epsilon = o$  and  $g = 0$ , then  $X = S^2 \times S^1$ .

By the above theorems we see that

$$Z(X; r) = \left[ \sum_{j \in \mathcal{I}} e^{2\pi i r q_j} \sum_{l=l_0}^N c_l^j r^{-l/2} \right] + o(r^{-N/2}) \quad (52)$$

for every  $N \geq l_0$  in the limit  $r \rightarrow \infty$ , where  $l_0$  is some fixed integer (depending on  $X$ ), and  $\mathcal{I} = \mathcal{I}_1 \amalg \mathcal{I}_2$  if  $E \neq 0$  and  $\mathcal{I} = \mathcal{I}_2$  if  $E = 0$ . By Proposition 4.3 the term  $Z_{\text{polar}}(X; r)$  has a structure like (30). In Theorem 4.1 we have separated the ‘exact part’  $Z_{\text{polar}}(X; r)$  from a part  $Z_{\text{int}}(X; r)$ , which is only given an asymptotic description. However, one should note that in the expression (52) there are contributions to the terms with  $j \in \mathcal{I}_2^a$  coming both from  $Z_{\text{polar}}(X; r)$  and  $Z_{\text{int}}(X; r)$ .

If we work with non-normalized Seifert invariants, the different quantities involved in the above Theorems 4.1 and 4.4 potentially depends on the actual choice of invariants. The precise situation is described in the following proposition, where we remind the reader that for the large Seifert manifolds the Seifert invariants in fact classify the Seifert manifolds up to an orientation preserving homeomorphism. Thus all quantities independent of the choice of non-normalized Seifert invariants are topological invariants for these large Seifert manifolds. Moreover, these topological invariants can then be calculated using any set of non-normalized Seifert invariants for a given large Seifert manifold.

**Proposition 4.5.** *We have*

$$Z(X; r) = Z_{\text{polar}}(X; r) + Z_{\text{int}}(X; r)$$

where  $Z_{\text{polar}}(X; r)$  and  $Z_{\text{int}}(X; r)$  are as in Theorem 4.1 if  $E \neq 0$  and  $Z_{\text{int}}(X; r) = 0$  if  $E = 0$ . In both cases

$$Z_{\text{polar}}(X; r) = \sum_{(l, \underline{n}') \in \mathcal{I}_2} \exp(2\pi i r q(l, \underline{n}')) Q^{(l, \underline{n}')} (r)$$

where  $q(l, \underline{n}')$  is given by (40) and the  $Q^{(l, \underline{n}')} (r)$  are Laurent polynomials in  $r$ . The expressions  $Z_{\text{polar}}(X; r)$  and  $Z_{\text{int}}(X; r)$  do not depend on the choice of non-normalized Seifert invariants for  $X$ . More specifically, let  $\Delta = \{\epsilon; g; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\}$  and  $\tilde{\Delta} = \{\epsilon; g; (\tilde{\alpha}_1, \tilde{\beta}_1), \dots, (\tilde{\alpha}_{\tilde{n}}, \tilde{\beta}_{\tilde{n}})\}$  be two sets of non-normalized Seifert invariants for  $X$ , let  $\mathcal{I}_1$  and  $\mathcal{I}_2$  be the index sets related to  $\Delta$  given by (42) and (43) respectively and let  $\tilde{\mathcal{I}}_1$  and  $\tilde{\mathcal{I}}_2$  be the corresponding index sets related to  $\tilde{\Delta}$ . Then we have a one-one correspondence between  $\mathcal{I}_j$  and  $\tilde{\mathcal{I}}_j$  for  $j = 1, 2$ , and under these correspondences the functions  $q$  in (39) and (40) are preserved. Moreover, the large  $r$  asymptotic expansion of  $Z_{\text{int}}(X; r)$ , i.e. the coefficients in (49) and (50) are preserved. Finally the Laurent polynomials  $Q^{(l, \underline{n}')} (r)$  are preserved if  $E \neq 0$  or  $E = 0$  and  $\sum_{j=1}^n \frac{1}{\alpha_j} < n + a_\epsilon g - 2$ .

**Proof.** The proof is simply a matter of routine checks. We first consider the case  $E = -\sum_{j=1}^n \frac{\beta_j}{\alpha_j} \neq 0$ . Let us write a tilde on each quantity or set relating to  $\tilde{\Delta}$ . Thus we write  $\tilde{\mathcal{S}}$  instead of  $\mathcal{S}$  etc. Let us first assume that  $\tilde{\Delta}$  is obtained from  $\Delta$  by adding a trivial  $(\alpha, \beta)$ -pair  $(1, 0)$ . Thus  $\tilde{n} = n + 1$  and  $(\tilde{\alpha}_j, \tilde{\beta}_j) = (\alpha_j, \beta_j)$  for  $j = 1, 2, \dots, n$  and  $(\tilde{\alpha}_{n+1}, \tilde{\beta}_{n+1}) = (1, 0)$ . We then have  $\tilde{\mathcal{S}} = \mathcal{S} \times \{0\}$  and  $\tilde{z}_{\text{st}}(m, (\underline{n}, 0)) = z_{\text{st}}(m, \underline{n})$  for all  $(m, \underline{n}) \in \mathbb{Z} \times \mathcal{S}$ . Thus we also have  $\tilde{\mathcal{I}}_1 = \mathcal{I}_1 \times \{0\}$ . We let  $(\tilde{\rho}_j, \tilde{\sigma}_j) = (\rho_j, \sigma_j)$  for  $j = 1, 2, \dots, n$  and put  $(\tilde{\rho}_{n+1}, \tilde{\sigma}_{n+1}) = (0, 1)$  and find that  $\tilde{q}_{(m, (\underline{n}, 0))} = q_{(m, \underline{n})}$  for all  $(m, \underline{n}) \in \mathcal{I}_1$ . For  $(m, \underline{n}) \in \mathcal{I}_1$  and  $k = 0, 1, 2, \dots$  it follows immediately from (49) that  $\tilde{c}_k^{(m, (\underline{n}, 0))} = c_k^{(m, \underline{n})}$ . Next, let us look at the set  $\tilde{\mathcal{I}}_2$  and the related quantities. We have  $\tilde{M} = M \times \mathbb{Z}$  and then  $\tilde{\mathcal{I}}_2 = \mathcal{I}_2 \times \{0\}$ . Thus we also have that  $\tilde{\mathcal{I}}_2^a = \mathcal{I}_2^a \times \{0\}$ . For  $(l, \underline{n}') \in \mathcal{I}_2$  we immediately get that  $\tilde{q}_{(l, (\underline{n}', 0))} = q_{(l, \underline{n}')}$ ,  $\tilde{b}_{(l, (\underline{n}', 0))} = b_{(l, \underline{n}')}$  and  $\tilde{Z}_\nu^{(l, (\underline{n}', 0))}(r) = Z_\nu^{(l, \underline{n}')} (r)$ ,  $\nu = 0, 1$ . Finally, let us check that the coefficients in (50) stay unchanged. First assume that  $n$  is even. Then  $k_0 = 0$ ,  $n + 1$  is odd and  $\tilde{k}_0 = 1$  so  $\tilde{k}_1 = (k_0 - (n + 1) - a_\epsilon g + 2)/2 = k_1$ . If  $k \in \{k_1, k_1 + 1, \dots\}$ , then  $\tilde{k}' = 2k + (n + 1) + a_\epsilon g - 2 = k' + 1$ .

We note that  $\tilde{f}(z; l, (\underline{n}', 0), (\underline{\mu}', \mu'_{n+1})) = -\pi z f(z; l, \underline{n}', \underline{\mu}')$ , and that

$$\partial_z^{(k'+1)} z f(z; l, \underline{n}', \underline{\mu}') \Big|_{z=0} = (k' + 1) \partial_z^{(k')} f(z; l, \underline{n}', \underline{\mu}') \Big|_{z=0}.$$

This shows that  $\tilde{c}_k^{(l, (\underline{n}', 0))} = c_k^{(l, \underline{n}')}.$  Next assume that  $n$  is odd. Then  $k_0 = 1$ ,  $n + 1$  is even and  $\tilde{k}_0 = 0$ , so  $\tilde{k}_1 = k_1 - 1$ . Like before we find that  $\tilde{c}_k^{(l, (\underline{n}', 0))} = c_k^{(l, \underline{n}')} for  $k \in \{k_1, k_1 + 1, \dots\}$ . Finally  $\tilde{c}_{k_1}^{(l, (\underline{n}', 0))} = 0$  since  $\tilde{f}(0; l, (\underline{n}', 0), (\underline{\mu}', \mu'_{n+1})) = 0$ .$

Next, consider the case where  $\tilde{n} = n$ ,  $\tilde{\alpha}_j = \alpha_j$  and  $\tilde{\beta}_j = \beta_j + K_j \alpha_j$  for some integers  $K_j$  with  $\sum_{j=1}^n K_j = 0$ . Let  $\tilde{\rho}_j = \rho_j$  and  $\tilde{\sigma}_j = \sigma_j + K_j \rho_j$ ,  $j = 1, 2, \dots, n$ . We immediately get that  $\tilde{\mathcal{S}} = \mathcal{S}$  and that  $\tilde{z}_{\text{st}}(m, \underline{n}) = z_{\text{st}}(m, \underline{n})$  and  $\tilde{q}_{(m, \underline{n})} = q_{(m, \underline{n})}$  for  $(m, \underline{n}) \in \mathbb{Z} \times \mathcal{S}$ . Moreover,  $\tilde{\mathcal{I}}_1 = \mathcal{I}_1$  and the coefficients in (49) are unchanged. Next let us consider  $\tilde{\mathcal{I}}_2$  and the quantities related to that index set. First note that  $(0, \underline{n}') \in \tilde{\mathcal{I}}_2$  if and only if  $(0, \underline{n}') \in \mathcal{I}_2$  and that all quantities related to such a point  $(0, \underline{n}')$  are the same w.r.t. the two sets of non-normalized Seifert invariants  $\Delta$  and  $\tilde{\Delta}$ . Let us next consider a point  $(1, \underline{n}') \in \tilde{\mathcal{I}}_2$ . Thus  $n'_j \in \mathbb{Z} + \frac{1}{2}\beta_j$  if  $\alpha_j$  or  $K_j$  is even and  $n'_j \in \mathbb{Z} + \frac{1}{2}\beta_j + \frac{1}{2}$  if both  $\alpha_j$  and  $K_j$  are odd. But if  $\alpha_j$  is odd then  $\alpha_j/2 \in \frac{1}{2} + \mathbb{Z}$  so in all cases there is a one-one correspondence between the set of points  $(l, \underline{n}') \in \tilde{\mathcal{I}}_2$  with  $l = 1$  and the set of points  $(l, \underline{n}') \in \mathcal{I}_2$  with  $l = 1$ . A correspondence is given as follows: For  $(1, \underline{n}') \in \mathcal{I}_2$ , let  $n''_j = n'_j + \frac{1}{2}K_j \alpha_j \pmod{\alpha_j}$  such that  $0 \leq n''_j < \alpha_j$  and let  $\tilde{n}'_j = n''_j$  if  $0 \leq n''_j \leq \alpha_j/2$  and  $\tilde{n}'_j = -n''_j + \alpha_j$  otherwise. Then  $(1, \underline{\tilde{n}'}) \in \tilde{\mathcal{I}}_2$ . To see that  $\tilde{q}_{(1, \underline{\tilde{n}'})} = q_{(1, \underline{n}')}$  we observe that the  $q$ -function in (40) only depends on  $n'_j \pmod{\alpha_j}$  and is unchanged when changing the sign of  $n'_j$ . Moreover,

$$\frac{\rho_j}{\alpha_j} \left( n'_j + \frac{1}{2} K_j \alpha_j \right)^2 - (\sigma_j + \rho_j K_j) (\beta_j + \alpha_j K_j) = \frac{1}{4} K_j \pmod{\mathbb{Z}}.$$

We have  $\tilde{b}_{(1, \underline{\tilde{n}'})} = (-1)^{\sum_{j=1}^n K_j \rho_j} b_{(1, \underline{n}')}$  since  $\tilde{n}'_j/\alpha_j \in \frac{1}{2}\mathbb{Z}$  if and only if  $n'_j/\alpha_j \in \frac{1}{2}\mathbb{Z}$ . Assume that  $\sum_{j=1}^n \frac{\mu'_j \tilde{n}'_j}{\alpha_j} \in \mathbb{Z}$  and let  $\tilde{\mu}'_j = \mu'_j$  if  $0 \leq n''_j \leq \alpha_j/2$  and  $\tilde{\mu}'_j = -\mu'_j$  otherwise, where  $n''_j$  is as above. Then

$$\sum_{j=1}^n \frac{\tilde{\mu}'_j \tilde{n}'_j}{\alpha_j} = \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} + \frac{1}{2} \sum_j \mu'_j K_j \pmod{\mathbb{Z}}.$$

But  $\frac{1}{2} K_j - \frac{1}{2} \mu'_j K_j \in \mathbb{Z}$  so  $\frac{1}{2} \sum_j \mu'_j K_j \in \mathbb{Z}$ . We also have

$$\tilde{f}(z; 1, \underline{\tilde{n}'}, \underline{\tilde{\mu}'}) (-1)^{\sum_{j=1}^n K_j \rho_j} f(z; l, \underline{n}', \underline{\mu}').$$

These observations show that  $\tilde{c}_k^{(1, \underline{\tilde{n}'})} = c_k^{(1, \underline{n}')} in (50). Finally, let us notice that$

$$\tilde{b}_{(1, \underline{\tilde{n}'})} \left( \tilde{Z}_0^{(1, \underline{\tilde{n}'})}(r) + \tilde{Z}_1^{(1, \underline{\tilde{n}'})}(r) \right) = b_{(1, \underline{n}')} \left( Z_0^{(1, \underline{n}')} (r) + Z_1^{(1, \underline{n}')} (r) \right).$$

The easiest way to see this is to refer to the actual calculation of  $Z_{\text{polar}}(X; r)$ . Thus, for  $l \in \mathbb{Z}$  let  $J'_l$  be the set in (116) and let  $\tilde{J}'_l$  be the corresponding set w.r.t.  $\tilde{\Delta}$ . Then  $\tilde{J}'_l = J'_l$  for each even  $l$  and for  $l$  odd the map  $\underline{n}' \mapsto \underline{\tilde{n}'}$  described above gives a bijection from  $J'_l$  to  $\tilde{J}'_l$ . Moreover, if  $Z^l(\delta, \eta)$  is given by (117) and  $\tilde{Z}^l(\delta, \eta)$  is the corresponding function w.r.t.  $\tilde{\Delta}$  then  $\tilde{Z}^l(\delta, \eta) = Z^l(\delta, \eta)$  proving the claim. Note that we do not necessarily have  $\tilde{b}_{(1, \underline{\tilde{n}'})} \tilde{Z}_\nu^{(1, \underline{\tilde{n}'})}(r) = b_{(1, \underline{n}')} Z_\nu^{(1, \underline{n}')} (r)$  for  $\nu = 0, 1$ .

Let us next assume that  $E = 0$ . Exactly as above we get that there is a one-one correspondence between  $\mathcal{I}_2$  and  $\tilde{\mathcal{I}}_2$  and under this correspondence the function  $q$  in (40) is preserved. Moreover, we know that  $Z(X; r)$  is independent of the choice of non-normalized Seifert invariants. Assume  $\sum_{j=1}^n \frac{1}{\alpha_j} < n + a_\epsilon g - 2$ . Note that this condition is independent of the choice of non-normalized

Seifert invariants. In this case we have  $Z(X; r) = Z_{\text{polar}}(X; r)$ , where  $Z_{\text{polar}}(X; r)$  is given by (46) with  $E = 0$ . Thus in this case we get that  $Q^{(l, n')}(r) = b_{(1, \underline{n}')} \left( Z_0^{(1, \underline{n}')} (r) + Z_1^{(l, \underline{n}')} (r) \right)$  is independent of the choice of non-normalized Seifert invariants exactly as in the case  $E = 0$ .  $\square$

In case  $E = 0$  it follows from Theorem 4.4 that the large  $r$  asymptotics of  $Z(X; r)$  is given by  $Z_{\text{polar}}(X; r)$  in (46) (with  $E = 0$ ) except for the special cases  $\sum_{j=1}^n \frac{1}{\alpha_j} \geq n + a_\epsilon g - 2$ , where we also have the extra term  $Z_{\text{spec}}(X; r)$ .

Let us determine which of the Seifert manifolds with  $E = 0$  that satisfy the condition  $\sum_{j=1}^n \frac{1}{\alpha_j} \geq n + a_\epsilon g - 2$ . As noted above the condition is independent of the choice of non-normalized Seifert invariants if we work with such invariants. We can therefore safely restrict to consider (normalized) Seifert invariants. This also allows us to keep track of the types of Seifert fibrations. Therefore, let  $X = (\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$ . Recall that  $0 < \beta_j < \alpha_j$  so  $\alpha_j \geq 2$  for all  $j = 1, \dots, n$ . Thus  $\sum_{j=1}^n \frac{1}{\alpha_j} \leq \frac{n}{2}$  so  $\sum_{j=1}^n \frac{1}{\alpha_j} \geq n + a_\epsilon g - 2$  implies that  $2 - a_\epsilon g \geq \frac{n}{2}$ , and this is only satisfied in the cases  $g = 0, \epsilon = o, n \in \{0, 1, 2, 3, 4\}$  and  $g = 1, \epsilon = o, n = 0$  and  $g = 2, \epsilon = n, n = 0$  (recall that we only consider even genus for  $\epsilon = n$ ). In the last two cases the Seifert Euler number  $E = -b$ , so since this number is zero we have  $X = (o; 1 \mid 0) = T^2 \times S^1$  (the 3-torus) or  $X = (n; 2 \mid 0)$ , both being small Seifert manifolds. In case  $\epsilon = o, g = 0$ , and  $n \in \{0, 1, 2\}$ ,  $X$  is a lens space  $L(p, q)$ , see [Or, Sect. 5.4 (i) pp. 99–100]. We find in all these cases that the Seifert Euler number  $E = 0$  if and only if  $p = 0$ , hence  $X = S^2 \times S^1$  if  $E = 0$ . The remaining cases to consider are  $\epsilon = o, g = 0$  and  $n \in \{3, 4\}$ . Let us first consider the easy case  $n = 4$ . Here the  $\alpha_j$ 's have to satisfy  $\sum_{j=1}^4 \frac{1}{\alpha_j} \geq 2$ , hence we have  $\alpha_j = 2$  for all  $j \in \{1, 2, 3, 4\}$ . In this case  $E = -b - 4\frac{1}{2} = -b - 2$ , so  $b = -2$  when  $E = 0$ . Thus  $X = (o; 0 \mid -2; (2, 1), (2, 1), (2, 1), (2, 1))$ . This is the small Seifert manifold considered in [Or, Sec. 5.4 (iii) p. 101]. In fact, this manifold is homeomorphic by an orientation preserving homeomorphism to the Seifert manifold  $(n; 2 \mid 0)$ , cf. [JN, Theorem 5.1]. Let us finally consider the case  $\epsilon = o, g = 0, n = 3$ . Here we search for solutions to  $\sum_{j=1}^3 \frac{1}{\alpha_j} \geq 1$ . The subset of these satisfying the strict inequality  $\sum_{j=1}^3 \frac{1}{\alpha_j} > 1$  are the small Seifert manifolds considered in [Or, Sec. 5.4 (ii) pp. 100–101]. By a direct inspection one finds that none of these manifolds have Seifert Euler number  $E = 0$ . Finally we consider the manifolds with  $\sum_{j=1}^3 \frac{1}{\alpha_j} = 1$ . The only solutions to this equation with  $\alpha_1 \leq \alpha_2 \leq \alpha_3$  are  $(2, 3, 6)$ ,  $(2, 4, 4)$  and  $(3, 3, 3)$ . A direct inspection reveals that we have the following possibilities with  $E = 0$ :

$$\begin{aligned} X_1 &= (o; 0 \mid -1; (2, 1), (3, 1), (6, 1)), \\ X_2 &= (o; 0 \mid -2; (2, 1), (3, 2), (6, 5)) = -X_1, \\ X_3 &= (o; 0 \mid -1; (2, 1), (4, 1), (4, 1)), \\ X_4 &= (o; 0 \mid -2; (2, 1), (4, 3), (4, 3)) = -X_3, \\ X_5 &= (o; 0 \mid -1; (3, 1), (3, 1), (3, 1)), \\ X_6 &= (o; 0 \mid -2; (3, 2), (3, 2), (3, 2)) = -X_5, \end{aligned}$$

where, as usual,  $-X$  means  $X$  with the opposite orientation. Thus the Seifert fibrations with  $\sum_{j=1}^n \frac{1}{\alpha_j} \geq n + a_\epsilon g - 2$ ,  $a_\epsilon g$  even and  $E = 0$  are the small Seifert manifolds  $S^2 \times S^1$ ,  $(o; 1 \mid 0) = T^2 \times S^1$ ,  $(n; 2 \mid 0) \cong (o; 0 \mid -2; (2, 1), (2, 1), (2, 1), (2, 1))$  and the six 3-fibered large Seifert manifolds  $X_j$ ,  $j = 1, \dots, 6$ .

We saw above that all but 9 of the Seifert fibrations satisfying  $\sum_{j=1}^n \frac{1}{\alpha_j} \geq n + a_\epsilon g - 2$ ,  $a_\epsilon g$  even and  $E = 0$  are topologically the space  $S^2 \times S^1$ . There is an infinite number of such fibrations for  $S^2 \times S^1$ . If  $X = (o; 0 \mid b, (\alpha_1, \beta_1), (\alpha_2, \beta_2))$ , then  $E = -b - \beta_1/\alpha_1 - \beta_2/\alpha_2 = 0$  if and only

if  $b = -1$ ,  $\alpha_1 = \alpha_2$  and  $\beta_2 = \alpha_1 - \beta_1$ . Thus  $(o; 0 \mid 0)$  and  $(o; 0 \mid -1; (\alpha, \beta), (\alpha, \alpha - \beta))$ ,  $\alpha \geq 2$ , give the possible Seifert fibrations for  $S^2 \times S^1$ . (Note that there are no  $(o; 0 \mid b; (\alpha, \beta))$  with  $E = -b - \beta/\alpha = 0$  and  $0 < \beta < \alpha$ .) For all the Seifert fibrations satisfying  $\sum_{j=1}^n \frac{1}{\alpha_j} \geq n + a_\epsilon g - 2$ ,  $a_\epsilon g$  even and  $E = 0$  the term  $Z_{\text{spec}}(X; r) \neq 0$ .

**4.1. The moduli space of flat SU(2)–connections on Seifert manifolds.** It follows from the results in the previous section that the proof of Theorem 1.4 is finalized once we have shown that the values of the  $q$ –functions defined in (39) and (40) are in fact the Chern–Simons values of the flat SU(2)–connections on  $X$ . This is done by a simple comparison between the values of the  $q$ –functions and the values of the Chern–Simons functional of flat SU(2)–connections on the Seifert manifolds calculated by D. Auckly, cf. [Au1].

Let  $X$  be a Seifert manifold given by the Seifert invariants  $(\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  or the non-normalized Seifert invariants  $\{\epsilon; g; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\}$ . Let  $\beta_0$  be as in the previous section and let  $\mathcal{M}_X$  denote the moduli space of flat SU(2)–connections on  $X$ . There is a classical identification

$$\mathcal{M}_X = \text{Hom}(\pi_1(X), \text{SU}(2))/\text{SU}(2),$$

where  $/\text{SU}(2)$  means moduli conjugation. Let us recall Auckly's results. Assume first that  $\epsilon = o$  (so  $g \geq 0$ ). It is well-known that

$$\begin{aligned} \pi_1(X) &= \langle h, q_1, \dots, q_n, a_1, b_1, \dots, a_g, b_g \mid q_k^{\alpha_k} h^{\beta_k} = 1, h^{-\beta_0} q_1 \dots q_n \prod_{i=1}^g [a_i, b_i] = 1, \\ &\quad [a_j, h] = [b_j, h] = [q_k, h] = 1, j = 1, 2, \dots, g, k = 1, 2, \dots, n \rangle, \end{aligned}$$

cf. [Au1], [JN, Theorem 6.1]. Let  $g_1, \dots, g_n \in \text{SU}(2)$ ,  $\underline{n} \in Z^n$  and  $\varepsilon \in \{0, 1/2\}$ . Then Auckly defines representations

$$\omega = \omega(\varepsilon, \underline{n})[g_1, \dots, g_n] : \pi_1(X) \rightarrow \text{SU}(2)$$

in the following way, cf. [Au2, Definition p. 56], [Au1, Definition p. 231]. Let  $\tilde{\omega}$  be the homomorphism from the free group generated by  $h, q_1, \dots, q_n, a_1, b_1, \dots, a_{g-1}, b_{g-1}$  to  $\text{SU}(2)$  given by  $\tilde{\omega}(h) = e^{2\pi i \varepsilon}$ ,  $\tilde{\omega}(a_j) = \tilde{\omega}(b_j) = 1$ ,  $j = 1, \dots, g-1$ , and  $\tilde{\omega}(q_k) = g_k \exp(2\pi i(n_k - \varepsilon \beta_k)/\alpha_k) g_k^{-1}$ ,  $k = 1, \dots, n$ , where we identify  $\text{SU}(2)$  by the unit quaternions  $\text{Sp}_1$  in the usual way. Let

$$a + jb = \tilde{\omega}(q_n^{-1} \dots q_1^{-1} h^{\beta_0}), \quad a, b \in \mathbb{C}.$$

If  $a = 1$  (so  $b = 0$ ) we let  $x = y = 1$ . If not we let  $x \in S^1$  be a square root of  $\frac{a(2-2\text{Re}(a)) - |b|^2}{2-2\text{Re}(a) - |b|^2}$  and  $y = \frac{\bar{b}}{r(x^2-1)} + jr$ , where  $r \in [-1, 1]$  satisfies  $r^2 = 1 - \frac{|b|^2}{2-2\text{Re}(a)}$  (noting that  $1 - \frac{|b|^2}{2-2\text{Re}(a)} \geq 0$ ). We then define  $\omega$  by letting  $\omega(\gamma) = \tilde{\omega}(\gamma)$ ,  $\gamma \in \{h, q_1, \dots, q_n, a_1, b_1, \dots, a_{g-1}, b_{g-1}\}$ , and by letting  $\omega(a_g) = x$  and  $\omega(b_g) = y$ . (Thus, in case  $a \neq 1$ , there are four possible choices of  $\omega$  corresponding to the two possible choices of each of  $x$  and  $y$ . All four choices are needed in Proposition 4.6.) If  $g \geq 1$  then  $\omega$  extends to a  $\text{SU}(2)$  representation of  $\pi_1(X)$ . If  $g = 0$  then  $\omega$  extends to a representation  $\pi_1(X) \rightarrow \text{SU}(2)$  if and only if

$$e^{-2\pi i \beta_0 \varepsilon} \prod_{j=1}^n g_j \exp\left(2\pi i \left(\frac{n_j - \varepsilon \beta_j}{\alpha_j}\right)\right) g_j^{-1} = 1. \quad (53)$$

Above we can of course restrict to  $\underline{n} \in \mathcal{S}$ . If the Seifert Euler number  $E = -\beta_0 - \sum_{j=1}^n \frac{\beta_j}{\alpha_j} \neq 0$  then we define a representation

$$\rho = \rho(m, \underline{n}) : \pi_1(X) \rightarrow \text{SU}(2)$$

for each integer  $m$  and each  $\underline{n} \in \mathcal{S}$  by letting  $\rho(a_j) = \rho(b_j) = 1$ ,  $j = 1, \dots, g$ , and by putting

$$\begin{aligned}\rho(h) &= \exp\left(-2\pi i \frac{1}{E} \left(m + \sum_{k=1}^n \frac{n_k}{\alpha_k}\right)\right), \\ \rho(q_j) &= \exp\left(2\pi i \left(\frac{n_j}{\alpha_j} + \frac{1}{E} \frac{\beta_j}{\alpha_j} \left(m + \sum_{k=1}^n \frac{n_k}{\alpha_k}\right)\right)\right).\end{aligned}$$

Note that the image of  $\rho$  is contained in  $\mathbb{C} \cap \mathrm{Sp}_1 = \mathrm{U}(1) = S^1$ , so  $\rho$  is reducible. We have

**Proposition 4.6** ([Au2, Lemma p. 57]). *Let the situation be as above. If  $E = 0$  then any element of  $\mathcal{M}_X$  is in the same path component as one of the conjugacy classes of the  $\omega$ 's. If  $E \neq 0$  then any element of  $\mathcal{M}_X$  is in the same path component as one of the conjugacy classes of the  $\omega$ 's and  $\rho$ 's.*  $\square$

If  $l \in \{0, 1\}$ ,  $\underline{n}' \in \mathbb{Z}^n + \frac{1}{2}l\beta$  and  $g_1, \dots, g_n \in \mathrm{SU}(2)$  then we let  $\varepsilon = l/2$  and  $n_j = n'_j + \varepsilon\beta_j$  and let  $\omega'(l, \underline{n}')[g_1, \dots, g_n]$  be equal to  $\omega(\varepsilon, \underline{n})[g_1, \dots, g_n]$ . We note that  $\omega'(l, -\underline{n}')[g_1, \dots, g_n] = \omega'(l, \underline{n}')[g_1, \dots, g_n]$  thus we can restrict to  $(l, \underline{n}') \in \mathcal{I}_2$ .

Next we consider the Seifert manifolds with non-orientable base. Therefore, let  $X$  be as above with  $\epsilon = n$  (so  $g > 0$ ). We have

$$\begin{aligned}\pi_1(X) &= \langle h, q_1, \dots, q_n, a_1, \dots, a_g \mid q_k^{\alpha_k} h^{\beta_k} = 1, h^{-\beta_0} q_1 \dots q_n \prod_{i=1}^g a_i^2 = 1, \\ &\quad a_j^{-1} h a_j h = [q_k, h] = 1, j = 1, 2, \dots, g, k = 1, 2, \dots, n \rangle,\end{aligned}$$

cf. [Au1], [JN, Theorem 6.1]. Let  $g_1, \dots, g_n \in \mathrm{SU}(2)$ ,  $\underline{n} \in \mathbb{Z}^n$  and  $\varepsilon \in \{0, 1/2\}$ . Then Auckly defines representations

$$\nu = \nu(\varepsilon, \underline{n})[g_1, \dots, g_n] : \pi_1(X) \rightarrow \mathrm{SU}(2)$$

in the following way, cf. [Au2, Definition p. 56], [Au1, Definition p. 231]. First let  $\tilde{\nu}$  be the homomorphism from the free group generated by  $h, q_1, \dots, q_n, a_1, \dots, a_{g-1}$  to  $\mathrm{SU}(2)$  given by  $\tilde{\nu}(h) = \exp(2\pi i \varepsilon)$ ,  $\tilde{\nu}(q_k) = g_k \exp(2\pi i (n_k - \varepsilon\beta_k)/\alpha_k) g_k^{-1}$ , and  $\tilde{\nu}(a_j) = 1$ ,  $j = 1, \dots, g-1$ ,  $k = 1, \dots, n$ . We then define  $\nu$  by letting  $\nu(\gamma) = \tilde{\nu}(\gamma)$ ,  $\gamma \in \{h, q_1, \dots, q_n, a_1, \dots, a_{g-1}\}$ , and by putting  $\nu(a_g) = \mathrm{sqr}(\tilde{\nu}(q_n^{-1} \dots q_1^{-1} h^{\beta_0}))$ , where  $\mathrm{sqr}(x)$  is any element of  $\mathrm{SU}(2)$  such that  $(\mathrm{sqr}(x))^2 = x$ ,  $x \in \mathrm{SU}(2)$ . (We know that such an element exists because there is an element  $z \in \mathrm{SU}(2)$  such that  $zxz^{-1} \in S^1$ . Therefore we have an element  $y \in S^1$  such that  $y^2 = zxz^{-1}$ . But then  $(z^{-1}yz)^2 = x$ .) This defines a representation  $\nu : \pi_1(X) \rightarrow \mathrm{SU}(2)$ . (We will see below, that the Chern–Simons value of  $[\nu] \in \mathcal{M}_X$  is independent of the choice of this square root, so therefore we don't specify this choice here.) As in the oriented base case we can parametrize slightly differently by letting  $\nu'(l, \underline{n}')[g_1, \dots, g_n]$  be equal to  $\nu(\varepsilon, \underline{n})[g_1, \dots, g_n]$  for each  $l \in \{0, 1\}$  and  $\underline{n}' \in \mathbb{Z}^n + \frac{1}{2}l\beta$  with  $\varepsilon = l/2$  and  $\underline{n} = \underline{n}' + \varepsilon\beta$ . Also here we can restrict to  $(l, \underline{n}') \in \mathcal{I}_2$ .

If the Euler number  $E = -\beta_0 - \sum_{j=1}^n \frac{\beta_j}{\alpha_j} \neq 0$  then we define a representation

$$\sigma = \sigma(m, \underline{n}) : \pi_1(X) \rightarrow \mathrm{SU}(2)$$

for each integer  $m$  and each  $\underline{n} \in \mathcal{S}$  by letting  $\sigma(a_i) = j$ ,  $i = 1, \dots, g$ , and by putting

$$\begin{aligned}\sigma(h) &= \exp\left(-2\pi i \frac{1}{E} \left(\frac{g}{2} + m + \sum_{k=1}^n \frac{n_k}{\alpha_k}\right)\right), \\ \sigma(q_j) &= \exp\left(2\pi i \left(\frac{n_j}{\alpha_j} + \frac{1}{E} \frac{\beta_j}{\alpha_j} \left(\frac{g}{2} + m + \sum_{k=1}^n \frac{n_k}{\alpha_k}\right)\right)\right).\end{aligned}$$

We then have

**Proposition 4.7** ([Au2, Lemma p. 62]). *Let the situation be as above. If  $E = 0$  then any element of  $\mathcal{M}_X$  is in the same path component as one of the conjugacy classes of the  $\nu$ 's. If  $E \neq 0$  then any element of  $\mathcal{M}_X$  is in the same path component as one of the conjugacy classes of the  $\nu$ 's and  $\sigma$ 's.*  $\square$

Using [KK1, Theorem 4.2] Auckly proved the following result, where CS denotes the Chern–Simons functional defined in (3) (with  $G = \mathrm{SU}(2)$ ).

**Theorem 4.8** ([Au1, pp. 232–234], [Au2, pp. 63–70]). *Let the situation be as above. Let  $\gamma$  be a  $\mathrm{SU}(2)$ –representation of  $\pi_1(X)$  equal to  $\omega(\varepsilon, \underline{n})[g_1, \dots, g_n]$  or  $\nu(\varepsilon, \underline{n})[g_1, \dots, g_n]$ . Thinking of the conjugacy class  $[\gamma]$  as a gauge equivalence class of flat connections we have*

$$\mathrm{CS}([\gamma]) = -\varepsilon c - \varepsilon^2 E - \sum_{j=1}^n \left( \frac{\rho_j n_j^2 + 2n_j \varepsilon}{\alpha_j} \right) \pmod{\mathbb{Z}},$$

where the  $\rho_j$ 's are as before and  $c = 0$  if  $\varepsilon = \mathbf{o}$  and  $c = g$  if  $\varepsilon = \mathbf{n}$ . Next assume that  $E \neq 0$  and let  $\gamma$  be a  $\mathrm{SU}(2)$ –representation of  $\pi_1(X)$  equal to  $\rho(m, \underline{n})$  or  $\sigma(m, \underline{n})$ . Thinking of the conjugacy class  $[\gamma]$  as a gauge equivalence class of flat connections we have

$$\begin{aligned} \mathrm{CS}([\gamma]) &= \frac{1}{E} \left( m + \frac{c}{2} \right) \left( m + \frac{c}{2} + \sum_{k=1}^n \frac{n_k}{\alpha_k} \right) \\ &\quad - \sum_{j=1}^n \frac{\rho_j n_j^2 - \frac{n_j}{E} \left( m + \frac{c}{2} + \sum_{k=1}^n \frac{n_k}{\alpha_k} \right)}{\alpha_j} \pmod{\mathbb{Z}}, \end{aligned}$$

where  $c$  and  $\rho_j$  are as before.  $\square$

Actually Auckly works with (normalized) Seifert invariants all over. To see that all the above results holds when working with non-normalized Seifert invariants simply note that all the calculations of Auckly stay unchanged with the exception that one should put  $b = 0$  all over in Auckly's calculations as indicated above. It is instructive to give a more direct argument for Theorem 4.8. First simply note that since this theorem is true for (normalized) Seifert invariants it is also true for the non-normalized Seifert invariants  $\{\varepsilon; g \mid (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n), (1, b)\}$ , with  $0 < \beta_j < \alpha_j$ ,  $j = 1, 2, \dots, n$ , since  $n_{n+1}$  has to be zero. Secondly, we note that two sets of non-normalized Seifert invariants for the same Seifert fibration give the same set of Chern–Simons values. Of course this has to be the case since isomorphic Seifert fibrations are homeomorphic. However, it is in fact also easy to see directly from the formulas in Theorem 4.8. Thus adding or deleting an  $(\alpha, \beta)$ –pair  $(1, 0)$  preserves the set of Chern–Simons invariants since if the  $j$ th pair is such a pair then  $n_j$  has to be zero. Secondly, if we change  $(\alpha_j, \beta_j)$  to  $(\alpha_j, \beta_j + K_j \alpha_j)$  with  $\sum_{j=1}^n K_j = 0$ , then we can keep the  $\rho_j$ 's unchanged and thus the set of Chern–Simons invariants don't change.

The above Theorem 4.8 actually corrects a small error in Auckly's result. Thus the term  $-\varepsilon c$  is missing in the formula for  $\mathrm{CS}(\nu(\varepsilon, \underline{n})[g_1, \dots, g_n])$  given in [Au1], [Au2]. Let us give a short account for why this extra term has to be included, following Auckly's proof most of the way. As mentioned above Auckly bases his proof of Theorem 4.8 on a result of Kirk and Klassen, namely [KK1, Theorem 4.2]. Let us work with normalized invariants. Thus assume  $X = (\mathbf{n}; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  and let  $X_j$  be the space obtained from  $X$  by cutting out the interior of a tubular neighborhood of the  $j$ th exceptional fiber,  $j = 1, \dots, n$ , and let  $X_0$  be the space obtained from  $X$  by cutting out the interior of a tubular neighborhood of a regular fiber.



Thus

$$\begin{aligned}\pi_1(X_j) &= \langle h, q_1, \dots, q_n, a_1, \dots, a_g \mid h^{-b} q_1 \dots q_n \prod_{i=1}^g a_i^2 = 1, q_k^{\alpha_k} h^{\beta_k} = 1, k \neq j, \\ &\quad a_m^{-1} h a_m h = [q_k, h] = 1, m = 1, 2, \dots, g, k = 1, 2, \dots, n \rangle\end{aligned}$$

and

$$\begin{aligned}\pi_1(X_0) &= \langle h, q_1, \dots, q_n, a_1, \dots, a_g \mid q_k^{\alpha_k} h^{\beta_k} = [q_k, h] = 1, k = 1, 2, \dots, n, \\ &\quad a_m^{-1} h a_m h = 1, m = 1, 2, \dots, g, \rangle.\end{aligned}$$

Let  $\rho_j, \sigma_j \in \mathbb{Z}$  such that  $\alpha_j \sigma_j - \beta_j \rho_j = 1$  as usual. We note that  $\mu_j = q_j^{\alpha_j} h^{\beta_j}$  and  $\lambda_j = q_j^{\rho_j} h^{\sigma_j}$  are respectively a meridian and a longitude for the torus neighborhood in  $X$  around the  $j$ th exceptional fiber. Moreover,  $\mu_0 = (h^{-b} q_1 \dots q_n \prod_{k=1}^g a_k^2)^{-1}$  and  $\lambda_0 = h$  are a meridian and longitude respectively for the torus neighborhood in  $X$  around a regular fiber. Let  $\nu = \nu(\varepsilon, \underline{n})[g_1, \dots, g_n]$  and let us define curves  $\nu_j = \nu_j(\varepsilon, \underline{n})[g_1, \dots, g_n] : [0, 1] \rightarrow \text{Hom}(\pi_1(N_j), \text{SU}(2))$  by  $\nu_j(t)(x) = \nu(x)$  for  $t \in [0, 1]$  and  $x = h, x = a_1, \dots, a_{g-1}$  and  $x = q_1, \dots, q_{j-1}$  (a curve being a continuous curve here and in the following). Moreover, we let  $\nu_j(t)(q_i) = g_i \exp\left(-2\pi i \varepsilon \frac{\beta_i}{\alpha_i}\right) g_i^{-1}$  for  $i = j+1, \dots, n$  and  $\nu_j(t)(q_j) = g_j \exp\left(2\pi i \frac{n_j t - \varepsilon \beta_j}{\alpha_j}\right) g_j^{-1}$ . Finally we define  $\nu_j(t)(A_g)$  in the following way: First choose curves  $h_j : [0, 1] \rightarrow \text{SU}(2)$  such that  $h_j(t)^{-1} \nu_j(t)(q_n^{-1} \dots q_1^{-1} h^b) h_j(t) \in S^1$  and choose curves  $\varphi_j : [0, 1] \rightarrow \mathbb{R}$  such that  $h_j(t)^{-1} \nu_j(t)(q_n^{-1} \dots q_1^{-1} h^b) h_j(t) = e^{2i\varphi_j(t)}$ . Since  $\nu_j(1)(x) = \nu_{j+1}(0)(x)$  for  $j = 1, 2, \dots, n$  and  $x = h, q_1, \dots, q_n, a_1, \dots, a_{g-1}$ , where  $\nu_{n+1}(0) = \nu$ , we can choose the  $h_j$  and  $\varphi_j$  such that  $h_j(1) = h_{j+1}(0)$  and  $\varphi_j(1) = \varphi_{j+1}(0)$ ,  $j = 1, \dots, n-1$  and such that  $h_n(1) e^{i\varphi_n(1)} h_n(1)^{-1} = \nu(a_g)$ . Thus  $u_j : [0, 1] \rightarrow \text{SU}(2)$  given by  $u_j(t) = h_j(t) e^{i\varphi_j(t)} h_j(t)^{-1}$  define curves such that  $u_j(t)^2 = \left(\nu_j(t) \left(h^{-b} q_1 \dots q_n \prod_{k=1}^{g-1} a_k^2\right)\right)^{-1}$  and such that  $u_j(1) = u_{j+1}(0)$ ,  $j = 1, \dots, n-1$  and  $u_n(1) = \nu(a_g)$ . We let  $\nu_j(t)(a_g) = u_j(t)$ . Thus  $\nu_j(1) = \nu_{j+1}(0)$  and  $\nu_n(1) = \nu$ . Also we note that  $\nu_j(0) = \nu(\varepsilon, n_1, \dots, n_{j-1}, 0, \dots, 0)[g_1, \dots, g_n]$  so  $\nu_j(0)$  and  $\nu_j(1)$  both extend to  $\text{SU}(2)$ -representations of  $\pi_1(X)$ . We note that  $\nu_j(t)(\mu_j) = g_j e^{2\pi i a_j(t)} g_j^{-1}$  and  $\nu_j(t)(\lambda_j) = g_j e^{2\pi i b_j(t)} g_j^{-1}$ , where  $a_j(t) = n_j t$  and  $b_j(t) = (\rho_j n_j t + \varepsilon)/\alpha_j$ . By [KK1, Theorem 4.2] we thus find that

$$\text{CS}(\nu) - \text{CS}(\nu_1(0)) = -2 \sum_{j=1}^n \int_0^1 b_j(t) a_j'(t) dt = - \sum_{j=1}^n \frac{\rho_j n_j^2 + 2n_j \varepsilon}{\alpha_j} \pmod{\mathbb{Z}}.$$

To calculate  $\text{CS}(\nu_1(0))$  we construct a curve  $\gamma : [0, 1] \rightarrow \text{Hom}(\pi_1(X_0), \text{SU}(2))$  from the trivial representation to  $\nu_1(0)$  and apply [KK1, Theorem 4.2] once more. To be specific we let

$$\gamma(t)(h) = \begin{cases} 1 & , t \in [0, 1/4], \\ e^{\pi i \varepsilon (4t-1)} & , t \in [1/4, 3/4], \\ e^{2\pi i \varepsilon} & , t \in [3/4, 1] \end{cases}$$

and

$$\gamma(t)(q_j) = \begin{cases} 1 & , t \in [0, 1/4], \\ e^{-\pi i \varepsilon \frac{\beta_j}{\alpha_j} (4t-1)} & , t \in [1/4, 3/4], \\ h_j(4t-3) e^{-2\pi i \varepsilon \frac{\beta_j}{\alpha_j}} h_j(4t-3)^{-1} & , t \in [3/4, 1] \end{cases}$$

for  $j = 1, \dots, n$ , where  $h_j : [0, 1] \rightarrow \mathrm{SU}(2)$  is a curve from 1 to  $g_j$ . Moreover we let

$$\gamma(t)(a_k) = \begin{cases} u_1(4t) & , t \in [0, 1/4], \\ j & , t \in [1/4, 3/4], \\ u_1(4 - 4t) & , t \in [3/4, 1] \end{cases}$$

for  $k = 1, \dots, g - 1$ , where  $u_1 : [0, 1] \rightarrow \mathrm{SU}(2)$  is a curve from 1 to  $j$ . Finally we let

$$\gamma(t)(a_g) = \begin{cases} u_1(4t) & , t \in [0, 1/4], \\ j & , t \in [1/4, 3/4], \\ u_2(4t - 3) & , t \in [3/4, 1], \end{cases}$$

where  $u_2 : [0, 1] \rightarrow \mathrm{SU}(2)$  is a curve from  $j$  to  $\nu_1(0)(a_g)$ . Thus  $\gamma$  indeed defines a curve in  $\mathrm{Hom}(\pi_1(X_0), \mathrm{SU}(2))$  from the trivial representation to  $\nu_1(0)$ . We recall that  $\lambda_0 = h$  and thus have that  $\gamma(t)(\lambda_0) = e^{2\pi i b(t)}$ , where

$$b(t) = \begin{cases} 0 & , t \in [0, 1/4], \\ \frac{\varepsilon}{2}(4t - 1) & , t \in [1/4, 3/4], \\ \varepsilon & , t \in [3/4, 1]. \end{cases}$$

Choose piecewise smooth curves  $g : [0, 1] \rightarrow \mathrm{SU}(2)$  and  $a : [0, 1] \rightarrow \mathbb{R}$  such that  $\gamma(t)(\mu_0) = g(t)^{-1} e^{2\pi i a(t)} g(t)$ . By [KK1, Theorem 4.2] we find that

$$\mathrm{CS}(\gamma(1)) - \mathrm{CS}(\gamma(0)) = -2 \int_0^1 b(t) a'(t) dt \pmod{\mathbb{Z}}.$$

Here

$$-2 \int_0^1 b(t) a'(t) dt = -4\varepsilon \int_{1/4}^{3/4} t a'(t) dt + \varepsilon (3a(3/4) - a(1/4)) - 2\varepsilon a(1).$$

For  $t \in [1/4, 3/4]$  we can choose  $g(t) = 1$  and  $a(t) = -\frac{g}{2} - \frac{\varepsilon}{2} E(4t - 1)$ , where  $E$  is the Seifert Euler number as usual. This gives

$$-2 \int_0^1 b(t) a'(t) dt = -\varepsilon g - \varepsilon^2 E - 2\varepsilon a(1).$$

Finally we note that  $a(1) \in \mathbb{Z}$  since  $\gamma(1)(\mu_0) = 1$ , since  $\gamma(1) = \nu_1(0)$  extends to a representation of  $\pi_1(X)$ .

We note that the Chern–Simons invariants only depend on the genus of the base in the case of non-orientable base and that the set of Chern–Simons invariants in that case only depends on the parity of the genus. The Chern–Simons value of  $\sigma(m, \underline{n})$  for genus  $g$  is equal to the Chern–Simons value of  $\sigma(m + l, \underline{n})$  for genus  $g - 2l$ . We can of course replace the  $\omega$  and  $\nu$  above by the  $\omega'$  and  $\nu'$ .

We are now ready to show that the values of the  $q$ -functions in (39) and (40) are equal to the Chern–Simons invariants of the flat  $\mathrm{SU}(2)$ -connections on the Seifert manifold. In fact we find that the values differ by a sign. This sign discrepancy is either due to a general sign error in [KK1] or else the AEC is only true if we work with the complex conjugated invariants (which are the invariants associated to the mirror category of the modular category induced by the representation theory of  $U_q(\mathfrak{sl}_2(\mathbb{C}))$ ). A similar phenomenon was observed in [AH]. In any case this sign is not serious and is solely related to sign conventions.

Let  $X$  and  $\beta_0$  be as before and let  $\gamma$  be an equal to  $\omega'(l, \underline{n}')[g_1, \dots, g_n]$  or  $\nu'(l, \underline{n}')[g_1, \dots, g_n]$ , where  $(l, \underline{n}') \in \mathcal{I}_2$  and  $g_1, \dots, g_n \in \mathrm{SU}(2)$ . Let  $\varepsilon = l/2$  and  $\underline{n} = \underline{n}' + \varepsilon \underline{\beta}$ . We assume that (53) is satisfied if  $(\varepsilon, g) = (0, 0)$  i.e. we assume that

$$e^{-\pi i \beta_0 l} \prod_{j=1}^n g_j \exp \left( 2\pi i \frac{n'_j}{\alpha_j} \right) g_j^{-1} = 1 \tag{54}$$

in that case. The number  $q_{(l, \underline{n}')}$  in (40) is given by

$$\begin{aligned} q_{(l, \underline{n}')} &= \sum_{j=1}^n \frac{\rho_j}{\alpha_j} n_j'^2 - \varepsilon^2 \sum_{j=1}^n \sigma_j \beta_j \\ &= \sum_{j=1}^n \frac{\rho_j}{\alpha_j} n_j^2 + \varepsilon^2 \sum_{j=1}^n \left( \frac{\rho_j \beta_j^2}{\alpha_j} - \sigma_j \beta_j \right) - 2\varepsilon \sum_{j=1}^n \frac{\rho_j \beta_j}{\alpha_j} n_j \pmod{\mathbb{Z}}. \end{aligned}$$

By using that  $\rho_j \beta_j = \alpha_j \sigma_j - 1$  we get  $\sum_{j=1}^n \frac{\rho_j \beta_j^2}{\alpha_j} = E + \sum_{j=1}^n \beta_j \sigma_j$  so

$$\begin{aligned} q_{(l, \underline{n}')} &= \sum_{j=1}^n \frac{\rho_j}{\alpha_j} n_j^2 + \varepsilon^2 E - 2\varepsilon \sum_{j=1}^n \frac{\alpha_j \sigma_j - 1}{\alpha_j} n_j \\ &= \varepsilon^2 E + \sum_{j=1}^n \frac{\rho_j n_j^2 + 2n_j \varepsilon}{\alpha_j} \pmod{\mathbb{Z}}, \end{aligned}$$

where we use that  $2\varepsilon \sum_{j=1}^n \sigma_j n_j \in \mathbb{Z}$ . We have thus shown that

$$q_{(l, \underline{n}')} = -\text{CS}([\gamma]).$$

Let us next assume that the Seifert Euler number is nonzero and let  $\gamma$  be equal to  $\rho(m, \underline{n})$  or  $\sigma(m, \underline{n})$ . Then

$$\text{CS}([\gamma]) = - \sum_j \frac{\rho_j}{\alpha_j} n_j^2 + \frac{1}{4} E \left( \frac{2}{E} \left[ m + \frac{c}{2} + \sum_{k=1}^n \frac{n_k}{\alpha_k} \right] \right)^2 \pmod{\mathbb{Z}},$$

where  $c$  is the number of cross caps in the base of  $X$ . If  $c$  is even we see that

$$\text{CS}([\gamma]) = -q_{(m', \underline{n})},$$

where  $m' = -m - \frac{c}{2}$  and  $q_{(m', \underline{n})}$  is given by (39).

There is a small appropriate remark to make here. Namely, in our asymptotic formula in Theorem 4.1, only the values  $q_{(m, \underline{n})}$  for which  $z_{\text{st}}(m, \underline{n}) \in ]0, 1[$  are present. By the symmetry considerations above Corollary 5.15 we indeed have that the image of the  $q$ -function in (39) is given by the values  $q_{(m, \underline{n})}$  for which  $z_{\text{st}}(m, \underline{n}) \in [0, 1]$ . If  $z_{\text{st}}(m, \underline{n}) = l \in \{0, 1\}$ , put  $n_j' = n_j + \frac{1}{2} \beta_j l$  for all  $j$  and get that  $\sum_{j=1}^n \frac{n_j'}{\alpha_j} = m \in \mathbb{Z}$ . Thus  $(l, \underline{n}') \in \mathcal{I}_2^a$ . Moreover,  $q_{(m, \underline{n})}$  in (39) is equal to  $q_{(l, \underline{n}')}$  in (40).

**4.2. The genus 0 case.** In the case where the base is  $S^2$  it follows from the previous section that not all values  $q_{(l, \underline{n}')}$ ,  $(l, \underline{n}') \in \mathcal{I}_2$ , in (40) have to be Chern–Simons values. Namely, the point  $(l, \underline{n}')$  corresponds to a representation if and only if there exists  $g_1, \dots, g_n \in \text{SU}(2)$  such that (54) is satisfied. Of course it could happen that  $q_{(l, \underline{n}')}$  is still equal to a Chern–Simons value for a point  $(l, \underline{n}') \in \mathcal{I}_2$  that does not satisfy (54) but there are certainly cases where this is not the case. To give an example consider the manifold  $M_{-1}$  obtained by surgery on the 3-sphere along the figure 8 knot with framing  $-1$ . This manifold is equal to the Seifert manifold  $(o; 0 \mid -1; (2, 1), (3, 1), (7, 1))$ . It is easy to see that  $\mathcal{I}_1$  is empty in this case. One finds that  $\mathcal{I}_2^a = \{(l = 0, \underline{n}' = \underline{0})\}$  and that  $\mathcal{I}_2^b$  contains 15 points  $(l, \underline{n}')$  with  $l = 0$  and 8 points  $(l, \underline{n}')$  with  $l = 1$ . However, only the single point in  $\mathcal{I}_2^a$  and two points  $(l, \underline{n}') \in \{(1, (1/2, 1/2, 3/2)), (1, (1/2, 1/2, 5/2))\}$  in  $\mathcal{I}_2^b$  correspond to  $\text{SU}(2)$ -representations of  $\pi_1(M_{-1})$ , namely they are the only three points in  $\mathcal{I}_2$  satisfying (57) below. The point in  $\mathcal{I}_2^a$  corresponds to the trivial connection and has Chern–Simons invariant 0. For the two points in  $\mathcal{I}_2^b$  corresponding to connections we find  $q_{(1, (1/2, 1/2, 3/2))} = 121/168 \pmod{\mathbb{Z}}$  and  $q_{(1, (1/2, 1/2, 5/2))} = 25/168 \pmod{\mathbb{Z}}$ . The other 21 points

in  $\mathcal{I}_2^b$  have the following 21 different values for  $q_{(l,\underline{n}')} \pmod{\mathbb{Z}}$ :  $1/2, 2/3, 1/6, 3/7, 5/7, 6/7, 5/8, 3/14, 5/14, 13/14, 2/21, 8/21, 11/21, 7/24, 1/42, 25/42, 37/42, 3/56, 19/56, 27/56, 1/168$ , none of which are equal to one of our three Chern–Simons invariants.

Thus, to finalize the proof of the AEC for the Seifert manifolds  $X$  with base  $S^2$  we have to prove that the terms in our large  $r$  asymptotic formulas for  $\tau_r(X)$  in Theorems 4.1 and 4.4, corresponding to points  $(l, \underline{n}') \in \mathcal{I}_2$  such that  $q_{(l,\underline{n}')}$  is not a Chern–Simons value, are zero. In the genus zero case the Laurent polynomial  $Z_0^{(l,\underline{n}')} (r)$  is always zero so left is to prove that  $Z_1^{(l,\underline{n}')} (r) = 0$  and that the coefficients  $c_k^{(l,\underline{n}')} in (50) are zero for such a 'non-contributing' point  $(l, \underline{n}') \in \mathcal{I}_2$ .$

Since  $je^{i\theta}j^{-1} = e^{-i\theta}$  we have that there for all  $(l, \underline{n}') \in \mathcal{I}_2^a$  exists  $g_1, \dots, g_n \in \text{SU}(2)$  such that (54) is satisfied. Thus the part  $Z_{\text{int}}(X; r)$  in Theorem 4.1 and also the part  $Z_{\text{spec}}(X; r)$  in Theorem 4.4 have an asymptotic expansion in accordance with the AEC, so the AEC will follow if we can prove the following

**Conjecture 4.9.** *Let  $X$  be a Seifert manifold with base  $S^2$  and let  $(l, \underline{n}') \in \mathcal{I}_2^b$  be a point such that there do not exist  $g_1, \dots, g_n \in \text{SU}(2)$  such that (54) is satisfied. Then  $Z_1^{(l,\underline{n}')} (r) = 0$ .*

Let us prove this conjecture for the cases with 3 or less exceptional fibers.

**Theorem 4.10.** *The AEC is true for  $G = \text{SU}(2)$  and  $X$  any Seifert 3-manifold with base  $S^2$  and  $n \leq 3$  exceptional fibers.*

**Proof.** The case  $n \leq 2$  are the lens spaces (considering  $S^2 \times S^1$  a lens space) and for that case the AEC immediately follows from our formulas since  $Z_1^{(l,\underline{n}')} (r) = 0$  for all  $(l, \underline{n}') \in \mathcal{I}_2$  in that case (since we calculate the residue of an entire function). (As mentioned in the introduction Jeffrey has already proved that the AEC holds for the lens spaces.)

Let us next look at the case  $n = 3$ . Let us work with non-normalized Seifert invariants so let  $X = \{0; 0 \mid (\alpha_1, \beta_1), (\alpha_2, \beta_2), (\alpha_3, \beta_3)\}$ . Let us first determine which of the pairs  $(l, \underline{n}') \in \mathcal{I}_2^b$  that corresponds to a  $\text{SU}(2)$ -representation of  $\pi_1(X)$ . Any element  $q \in \text{SU}(2) = \text{Sp}_1$  can be written  $q = r + ai + bj + ck$ , where  $r, a, b, c \in \mathbb{R}$  with  $r^2 + a^2 + b^2 + c^2 = 1$ . The number  $r$  is called the real part of  $q$  and any two elements of  $\text{SU}(2)$  are conjugate if and only if they have the same real part. Thus the conjugacy class of any  $q \in \text{SU}(2) \setminus \{\pm 1\}$  is topologically a 2-sphere. Now, for any element  $q \in \text{SU}(2)$  we write  $q = \cos(\theta) + \sin(\theta)(ai + bj + ck)$  for a unique  $\theta \in [0, \pi]$  and  $a, b, c \in \mathbb{R}$  with  $a^2 + b^2 + c^2 = 1$ . (If  $q = \pm 1$ , then of course  $\sin(\theta) = 0$  and  $a, b, c$  are redundant.) Recall here that any element  $q \in \text{SU}(2)$  is conjugate to an element of  $S^1 = \{x + yi \mid x, y \in \mathbb{R}, x^2 + y^2 = 1\} \subseteq \text{SU}(2)$ . Thus,  $q = ge^{i\theta}g^{-1}$  for a  $g \in \text{SU}(2)$ ,  $\theta$  as above. Let  $S_\theta$  be the conjugacy class consisting of the elements of  $\text{SU}(2)$  with real part  $\cos(\theta)$ ,  $\theta \in [0, \pi]$ .

Let us find all possible  $(\theta_1, \theta_2, \theta_3) \in [0, \pi]^3$  such that

$$g_1 e^{i\theta_1} g_1^{-1} g_2 e^{i\theta_2} g_2^{-1} g_3 e^{i\theta_3} g_3^{-1} = 1 \quad (55)$$

for some  $g_1, g_2, g_3 \in \text{SU}(2)$ . By conjugation we can assume that  $g_1 = 1$  and the above equation then becomes

$$e^{i\theta_1} g e^{i\theta_2} g^{-1} = h e^{i\theta_3} h^{-1}$$

for some  $g, h \in \text{SU}(2)$ , that is, we search for triples  $(\theta_1, \theta_2, \theta_3) \in [0, \pi]^3$  such that there exists a  $u \in S_{\theta_2}$  with  $e^{i\theta_1} u \in S_{\theta_3}$ , i.e. such that the real part of  $e^{i\theta_1} u$  is  $\cos(\theta_3)$ . Writing  $u = \cos(\theta_2) + \sin(\theta_2)(ai + bj + ck)$  we find that the real part of  $e^{i\theta_1} u$  is  $\cos(\theta_1) \cos(\theta_2) - a \sin(\theta_1) \sin(\theta_2)$ . If  $\theta_2 = 0$  then  $\theta_3 = \theta_1$ . If  $\theta_2 = \pi$ , then  $\theta_3 = \pi - \theta_1$ . In the other cases  $S_{\theta_2}$  is a 2-sphere and

$a$  runs through  $[-1, 1]$  when  $u$  runs through  $S_\theta$ . Thus we get that (55) is satisfied for some  $g_1, g_2, g_3 \in \text{SU}(2)$  if and only if

$$|\theta_1 - \theta_2| \leq \theta_3 \leq \min\{\theta_1 + \theta_2, 2\pi - \theta_1 - \theta_2\}. \quad (56)$$

Note that the special cases  $\theta_2 \in \{0, \pi\}$  are covered by this condition. Thus  $\omega'(l, \underline{n}')[g_1, g_2, g_3]$  is a  $\text{SU}(2)$ -representation of  $\pi_1(X)$  if and only if  $(l, \underline{n}') \in \mathcal{I}_2$  satisfies the condition

$$\left| \frac{n'_1}{\alpha_1} - \frac{n'_2}{\alpha_2} \right| \leq \frac{n'_3}{\alpha_3} \leq \min \left\{ \frac{n'_1}{\alpha_1} + \frac{n'_2}{\alpha_2}, 1 - \frac{n'_1}{\alpha_1} - \frac{n'_2}{\alpha_2} \right\}. \quad (57)$$

The points  $(l, \underline{n}') \in \mathcal{I}_2^a$  are the points satisfying (57) with at least one of the two  $\leq$  being an equality. We have to prove that if  $(l, \underline{n}') \in \mathcal{I}_2^b$  does not satisfy (57), then  $Z_1^{(l, \underline{n}')} (r) = 0$ . It is not hard to see that

$$\begin{aligned} Z_1^{(l, \underline{n}')} (r) &= \frac{i}{4} \left( \prod_{j=1}^3 \sin \left( \pi \rho_j \frac{2n'_j}{\alpha_j} \right) \right) \sum_{\substack{\underline{\mu}' \in \{\pm 1\}^3 \\ a(\underline{\mu}', \underline{n}') > 0}} \prod_{j=1}^3 \mu'_j \\ &\quad \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq a(\underline{\mu}', \underline{n}')}} \frac{1}{\text{Sym}_\pm(m) \text{Sym}_\pm(m - a(\underline{\mu}', \underline{n}'))} \end{aligned} \quad (58)$$

for all  $(l, \underline{n}') \in \mathcal{I}_2$ , where  $a(\underline{\mu}', \underline{n}') = \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j}$ . Now let  $(l, \underline{n}') \in \mathcal{I}_2^b$ . Then

$$Z_1^{(l, \underline{n}')} (r) = \frac{i}{4} \left( \prod_{j=1}^3 \sin \left( \pi \rho_j \frac{2n'_j}{\alpha_j} \right) \right) \sum_{\substack{\underline{\mu}' \in \{\pm 1\}^3 \\ a(\underline{\mu}', \underline{n}') > 0}} \prod_{j=1}^3 \mu'_j \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m < a(\underline{\mu}', \underline{n}')}} \frac{1}{\text{Sym}_\pm(m)}.$$

The double inequality (57) is completely symmetric in  $n'_1, n'_2$  and  $n'_3$ . This should be so since we in the above argument have numbered the exceptional fibers arbitrary. To see the symmetry directly from (57), note that  $(\theta_1, \theta_2, \theta_3)$  satisfies (56) if and only if there exist  $u_j \in S_{\theta_j}$  such that  $u_1 u_2 u_3 = 1$ . But this is equivalent to  $u_2 u_3 u_1^{-1} = 1$  etc. (where of course  $u_1^{-1} \in S_{\theta_1}$ ). Thus it is enough to consider the case  $\frac{n'_1}{\alpha_1} \leq \frac{n'_2}{\alpha_2} \leq \frac{n'_3}{\alpha_3}$ . Assume this and assume that  $\underline{n}'$  does not satisfy (57). Since the first inequality in (57) is satisfied we have that the second is not satisfied. Assume first that  $\sum_{j=1}^3 \frac{n'_j}{\alpha_j} > 1$  (in which case we have that the right-hand side of (57) is  $1 - \frac{n'_1}{\alpha_1} - \frac{n'_2}{\alpha_2}$  and that the second inequality in (57) is not satisfied). In this case we have

$$Z_1^{(l, \underline{n}')} (r) = \frac{i}{4} \left( \prod_{j=1}^3 \sin \left( \pi \rho_j \frac{2n'_j}{\alpha_j} \right) \right) \left( \frac{3}{2} + \frac{1}{2} \sum_{\substack{\underline{\mu}' \in \{\pm 1\}^3 \\ 0 < a(\underline{\mu}', \underline{n}') < 1}} \prod_{j=1}^3 \mu'_j \right).$$

By going through all the 8 possibilities for  $\underline{\mu}' \in \{\pm 1\}^3$  one finds that  $a(\underline{\mu}', \underline{n}') \in ]0, 1[$  if and only if  $\underline{\mu}' \in \{(1, 1, -1), (1, -1, 1), (-1, 1, 1)\}$ , thus  $Z_1^{(l, \underline{n}')} (r) = 0$ .

Finally assume that  $\sum_{j=1}^3 \frac{n'_j}{\alpha_j} < 1$ . In that case

$$Z_1^{(l, \underline{n}')} (r) = \frac{i}{8} \left( \prod_{j=1}^3 \sin \left( \pi \rho_j \frac{2n'_j}{\alpha_j} \right) \right) \sum_{\substack{\underline{\mu}' \in \{\pm 1\}^3 \\ 0 < a(\underline{\mu}', \underline{n}') < 1}} \prod_{j=1}^3 \mu'_j.$$

We still restrict to the case  $\frac{n'_1}{\alpha_1} \leq \frac{n'_2}{\alpha_2} \leq \frac{n'_3}{\alpha_3}$  and have then that the second inequality in (57) is not satisfied if and only if  $\frac{n'_3}{\alpha_3} > \frac{n'_1}{\alpha_1} + \frac{n'_2}{\alpha_2}$ . Assuming this we find that  $a(\underline{\mu}', \underline{n}') \in ]0, 1[$  if and only if  $\underline{\mu}' \in \{(1, 1, 1), (1, -1, 1), (-1, 1, 1), (-1, -1, 1)\}$ , so  $Z_1^{(l, \underline{n}')} (r) = 0$ .  $\square$

Let us for completeness also calculate  $Z_1^{(l, \underline{n}')} (r)$  for  $(l, \underline{n}') \in \mathcal{I}_2^a$ . Note first that  $\rho_j \frac{2n'_j}{\alpha_j} \in \mathbb{Z}$  if and only if  $n'_j \in \{0, \alpha_j/2\}$  and in these cases  $Z_1^{(l, \underline{n}')} (r) = 0$  by (58). For  $\underline{\mu}' \in \{\pm 1\}^3$ , let  $a(\underline{\mu}, \underline{n}') = \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j}$  as above. By assumption there exists a  $\underline{\mu}' \in \{\pm 1\}^3$  such that  $a(\underline{\mu}', \underline{n}') \in \{0, 1\}$ . Assume first that  $a(\underline{\mu}', \underline{n}') = 1$  for a  $\underline{\mu}' \in \{\pm 1\}^3$ . Then either  $\mu'_j = 1$  for  $j = 1, 2, 3$  or there is a  $j \in \{1, 2, 3\}$  with  $n'_j = 0$ . In the last case  $Z_1^{(l, \underline{n}')} (r) = 0$ . In both cases  $\sum_{j=1}^3 \frac{n'_j}{\alpha_j} = 1$ . By (58) we get that

$$Z_1^{(l, \underline{n}')} (r) = \frac{i}{4} \left( \prod_{j=1}^3 \sin \left( \pi \rho_j \frac{2n'_j}{\alpha_j} \right) \right) \left( 1 + \frac{1}{2} \sum_{\substack{\underline{\mu}' \in \{\pm 1\}^3 \\ 0 < a(\underline{\mu}', \underline{n}') < 1}} \prod_{j=1}^3 \mu'_j \right).$$

By symmetry we can assume that  $\frac{n'_1}{\alpha_1} \leq \frac{n'_2}{\alpha_2} \leq \frac{n'_3}{\alpha_3}$  as usual. If  $\frac{n'_3}{\alpha_3} = \frac{1}{2}$ , then  $Z_1^{(l, \underline{n}')} = 0$ . If that is not the case then  $a(\underline{\mu}', \underline{n}') \in ]0, 1[$  if and only if  $\underline{\mu}' \in \{(1, 1, -1), (1, -1, 1), (-1, 1, 1)\}$  and

$$Z_1^{(l, \underline{n}')} (r) = -\frac{i}{8} \prod_{j=1}^3 \sin \left( \pi \rho_j \frac{2n'_j}{\alpha_j} \right) \quad (59)$$

which is also valid in case  $\frac{n'_3}{\alpha_3} = \frac{1}{2}$ .

Next assume that  $a(\underline{\mu}', \underline{n}') = 0$  for a  $\underline{\mu}' \in \{\pm 1\}^3$ . If  $\sum_{j=1}^3 \frac{n'_j}{\alpha_j} = 1$  we are in the above case, so assume that  $\sum_{j=1}^3 \frac{n'_j}{\alpha_j} < 1$ . Assume also that  $\frac{n'_1}{\alpha_1} \leq \frac{n'_2}{\alpha_2} \leq \frac{n'_3}{\alpha_3}$  as always. Then

$$Z_1^{(l, \underline{n}')} (r) = \frac{i}{8} \left( \prod_{j=1}^3 \sin \left( \pi \rho_j \frac{2n'_j}{\alpha_j} \right) \right) \sum_{\substack{\underline{\mu}' \in \{\pm 1\}^3 \\ 0 < a(\underline{\mu}', \underline{n}') < 1}} \prod_{j=1}^3 \mu'_j.$$

In this case we find that  $a(\underline{\mu}', \underline{n}') \in ]0, 1[$  if and only if  $\underline{\mu}' \in \{(1, 1, 1), (1, -1, 1), (-1, 1, 1)\}$  so again we find that  $Z_1^{(l, \underline{n}')} (r)$  is given by (59). Thus we have proved that  $Z_1^{(l, \underline{n}')} (r)$  is given by (59) for all  $(l, \underline{n}') \in \mathcal{I}_2^a$ .

To prove the AEC for Seifert manifolds with base  $S^2$  and  $n \geq 4$  exceptional fibers, one can proceed as in the above proof, but the combinatorics is of course harder. For a given  $(\theta_1, \dots, \theta_n) \in [0, \pi]^n$  there exist  $g_1, \dots, g_n \in \text{SU}(2)$  such that

$$\prod_{j=1}^n g_j e^{i\theta_j} g_j^{-1} = 1$$

if and only if there exist  $u_j \in S_{\theta_j}$ ,  $j = 2, 3, \dots, n-1$  such that the real part of  $e^{i\theta_1} u_2 \dots u_{n-1}$  is  $\cos(\theta_n)$ . Note that the function that maps  $(u_2, \dots, u_{n-1})$  to the real part of  $e^{i\theta_1} u_2 \dots u_{n-1}$  is a continuous map  $S_{\theta_2} \times \dots \times S_{\theta_{n-1}} \rightarrow [-1, 1]$ , so the image is an interval. Thus this analysis leads to a condition of the form

$$a \leq \theta_3 \leq b$$

where  $a, b \in [0, \pi]$  are continuous functions in  $(\theta_1, \theta_2, \dots, \theta_{n-1})$ . The condition (56) is the special case where  $n = 3$ .

Let  $X$  be an integral homology sphere with  $n \geq 3$  exceptional fibers and let  $\mathcal{M}_\rho$  be the component in the moduli space of irreducible flat  $\mathrm{SU}(2)$ -connections on  $X$  containing the class represented by the connection corresponding to a representation  $\rho : \pi_1(X) \rightarrow \mathrm{SU}(2)$ . Then Fintushel and Stern [FS] have proved that  $\mathcal{M}_\rho$  is a closed manifold of dimension  $2m - 6$  if  $\rho(q_j) \notin \{\pm 1\}$  for exactly  $m \geq 3$  of the  $j \in \{1, 2, \dots, n\}$ . (Note that the only reducible connection is the trivial connection in this case.) We also refer to [KK2] regarding results on the components of the moduli space of flat  $\mathrm{SU}(2)$ -connections on a Seifert fibered integral homology sphere.

**4.3. Small Seifert manifolds with base  $\mathbb{RP}^2$ .** Let  $X$  be a Seifert manifold with base  $\mathbb{RP}^2$  and zero or one exceptional fiber. Except for a few cases  $X$  is then homeomorphic to a prism manifold, i.e. a Seifert manifold of the form  $(o; 0 \mid b; (2, 1), (2, 1), (\alpha, \beta))$ . The few exceptions are homeomorphic to lens spaces except  $(n; 0 \mid 0)$  which is homeomorphic to  $\mathbb{RP}^3 \# \mathbb{RP}^3$ , see [JN, Theorem 5.1] and [Or, Sec. 5.4 (vii) p. 102] (there seem to be some small mistakes in [Or]. Thus  $(n; 1 \mid \pm 1)$  both have cyclic fundamental group, namely  $\mathbb{Z}_4$ , and  $(n; 1 \mid 0; (\alpha, 1))$  also has cyclic fundamental group, namely  $\mathbb{Z}_{4\alpha}$ . All three cases are homeomorphic to lens spaces according to [JN, Theorem 5.1].) Since  $\mathbb{RP}^3$  is homeomorphic to  $L(2, 1)$  and since the RT-invariants are multiplicative under connected sums and since the AEC behaves nicely w.r.t. products of RT-invariants we obtain by Theorem 4.10 the following

**Corollary 4.11.** *The AEC is true for  $G = \mathrm{SU}(2)$  and  $X$  any Seifert manifold with base  $\mathbb{RP}^2$  and zero or one exceptional fiber.*

The AEC can actually be proved directly for  $X = (n; 1 \mid 0)$  by a small calculation: By (36) we namely have

$$Z(X; r) = \sum_{\gamma=1}^{r-1} (-1)^\gamma \sin\left(\frac{\pi}{r}\gamma\right),$$

and by an elementary calculation we find that

$$Z(X; r) = -f\left(\frac{\pi}{r}\right) \left(1 + e^{2\pi i r \frac{1}{2}}\right),$$

where  $f(x) = \frac{\sin(x)}{2(1+\cos(x))}$ . Thus we find that

$$\tau_r(X) = \frac{1}{\sqrt{2r}} f\left(\frac{\pi}{r}\right) \left(1 + e^{2\pi i r \frac{1}{2}}\right). \quad (60)$$

By Theorem 4.8 the Chern–Simons invariants of flat  $\mathrm{SU}(2)$ -connections on  $X$  are 0 (mod  $\mathbb{Z}$ ) and  $1/2$  (mod  $\mathbb{Z}$ ).

#### 4.4. The contribution from the trivial connection and the Casson–Walker invariant.

From a topological point of view it is not necessarily a good idea to bring the asymptotics of the quantum invariants on minimal form; it is actually more natural to sum over the connected components of the moduli space instead of over the image set of the Chern–Simons functional. Let us consider the case of lens spaces. The lens spaces are identical with the Seifert manifolds with base  $S^2$  and zero, one or two exceptional fibers. Let  $(p, q)$  be a pair of coprime integers and let  $L(p, q)$  be the associated lens space given by surgery on  $S^3$  along the unknot with surgery coefficient  $-p/q$ . Thus we can and will assume in the following that  $p \geq 0$ . We have  $\pi_1(L(p, q)) = \mathbb{Z}/p\mathbb{Z}$  for  $p > 0$ , and  $L(0, 1) = S^2 \times S^1$  with fundamental group  $\mathbb{Z}$ . Hence  $L(p, q)$  is a rational homology sphere iff  $p \neq 0$  and an integral homology sphere iff  $p = 1$ , and  $L(p, q)$  and  $L(p', q')$  are nonhomeomorphic if  $p' \neq p$ . It is a classical result that  $L(p, q)$  and  $L(p, q')$  are homeomorphic if and only if  $q' = \pm q \bmod p$  or  $qq' = \pm 1 \bmod p$ . A homeomorphism is

orientation preserving if and only if the relevant sign is  $+$ . In particular, we can always assume that  $0 < q < p$  if  $|p| > 1$ .

Let us compare  $\tau_r(L(p, q))$  with the Casson–Walker invariant of  $L(p, q)$ ,  $p \neq 0$ . First we note that  $L(1, q) = S^3$  for all  $q \in \mathbb{Z}$ . In our normalization we have

$$\tau_r(S^3) = \sqrt{\frac{2}{r}} \sin\left(\frac{\pi}{r}\right) = \sqrt{\frac{2}{r}} \frac{\pi}{r} (1 + c_1 r^{-1} + c_2 r^{-2} + \dots),$$

where  $c_1 = 0$ . Note also that  $\lambda_C(S^3) = 0$ , where  $\lambda_C$  is the Casson–invariant of integral homology spheres, cf. [AM]. We note that there is only one flat  $SU(2)$ –connection on  $S^3$ , namely the trivial one.

Next let us investigate the lens spaces with  $|p| > 1$ . Assume that  $p > q > 0$  and let  $q^*$  denote the inverse of  $q \bmod p$ . By [HT, Theorem 5] we have

$$\tau_r(L(p, q)) = \sqrt{\frac{2}{pr}} \exp\left(\frac{\pi i}{2r} S\left(\frac{q}{p}\right)\right) \sum_{n=0}^{p-1} c_n(r) \exp\left(2\pi i r \frac{q^*}{p} n^2\right),$$

where

$$c_n(r) = i \cos\left(\frac{\pi}{pr}\right) \sin\left(2\pi \frac{q^*}{p} n\right) \sin\left(\frac{2\pi n}{p}\right) + \sin\left(\frac{\pi}{pr}\right) \cos\left(2\pi \frac{q^*}{p} n\right) \cos\left(\frac{2\pi n}{p}\right).$$

(Use that  $L(p, q) \cong L(p, q^*)$  and that  $S(q^*/p) = S(q/p)$ . The shift from  $q$  to  $q^*$  is done to facilitate a comparison to Chern–Simons invariants, see below.) The moduli space  $\mathcal{M}_{L(p, q)}$  of flat  $SU(2)$ –connections on  $L(p, q)$  can be identified with the set  $\mathcal{M} = \{0, 1, \dots, [p/2]\}$  via the identification of  $\mathcal{M}_{L(p, q)}$  with the set of representations  $\mathbb{Z}/p\mathbb{Z} \rightarrow SU(2)$  moduli conjugation by  $SU(2)$ . Here  $[x]$  is the integer part of  $x \in [0, \infty[$ . To be precise the integer  $n \in \{0, 1, \dots, [p/2]\}$  corresponds to the representation which maps the generator  $e^{2\pi i/p}$  to  $e^{2\pi i n/p}$ . By [KK1, Theorem 5.1] the Chern–Simons invariant of the flat  $SU(2)$ –connection corresponding to  $n \in \mathcal{M}$  is equal to  $q_n = -\frac{q^*}{p} n^2$ . (Note that Kirk and Klassen use another convention for the lens spaces. Thus  $L(p, q)$  in [KK1] is equal to  $L(p, -q)$  here.) Using a symmetry under the change of  $n$  to  $p - n$  we immediately get a formula of the form

$$\tau_r(L(p, q)) = \sqrt{\frac{2}{pr}} \exp\left(\frac{\pi i}{2r} S\left(\frac{q}{p}\right)\right) \sum_{n=0}^{[p/2]} a_n(r) \exp\left(2\pi i r \frac{q^*}{p} n^2\right).$$

Here, for  $p$  even we have  $a_n(r) = 2c_n(r)$  for  $n = 1, 2, \dots, p/2 - 1$  and  $a_0(r) = \sin(\pi/pr)$  and  $a_{p/2}(r) = (-1)^{1+q^*} \sin(\pi/pr) = \sin(\pi/pr)$ . For  $p$  odd we have  $a_n(r) = 2c_n(r)$  for  $n = 1, 2, \dots, (p-1)/2$  and  $a_0(r) = \sin(\pi/pr)$ . The trivial connection corresponds to  $n = 0$ . Let us write

$$\tau_r^n(p, q) = \sqrt{\frac{2}{pr}} \exp\left(\frac{\pi i}{2r} S\left(\frac{q}{p}\right)\right) a_n(r) \exp\left(2\pi i r \frac{q^*}{p} n^2\right)$$

for  $n \in \mathcal{M}$ . Then

$$\tau_r^0(p, q) = \sqrt{\frac{2}{pr}} \sin\left(\frac{\pi}{pr}\right) \exp\left(\frac{\pi i}{2r} S\left(\frac{q}{p}\right)\right) = \sqrt{\frac{2}{pr}} \frac{\pi}{pr} (1 + c_1 r^{-1} + c_2 r^{-2} + \dots),$$

where

$$c_1 = \frac{\pi i}{2} S\left(\frac{q}{p}\right) = 6\pi i \lambda_{CW}(L(p, q)).$$

Here  $\lambda_{CW}$  is Walker’s extension of the Casson–invariant to rational homology spheres, cf. [W].



Let us next investigate the sum of  $\tau_r^n(p, q)$  for which  $q_n = 0 \bmod \mathbb{Z}$ . We have  $q_n = 0 \bmod \mathbb{Z}$  if and only if  $p$  divides  $n^2$ . Let us consider  $L(9, q)$ ,  $q \in \{1, 2, 4, 5, 7, 8\}$ . We have  $q_n = 0 \bmod \mathbb{Z}$  if and only if  $n \in \{0, 3\}$ . Moreover

$$\tau_r^3(p, q) = 2\sqrt{\frac{2}{pr}} \exp\left(\frac{\pi i}{2r} S\left(\frac{q}{p}\right)\right) c_3(r),$$

where

$$\begin{aligned} c_3(r) &= i \cos\left(\frac{\pi}{pr}\right) \sin\left(2\pi \frac{q^*}{3}\right) \sin\left(\frac{2\pi}{3}\right) + \sin\left(\frac{\pi}{pr}\right) \cos\left(2\pi \frac{q^*}{3}\right) \cos\left(\frac{2\pi}{3}\right) \\ &= \frac{i\sqrt{3}}{2} \cos\left(\frac{\pi}{pr}\right) \sin\left(2\pi \frac{q^*}{3}\right) - \frac{1}{2} \sin\left(\frac{\pi}{pr}\right) \cos\left(2\pi \frac{q^*}{3}\right). \end{aligned}$$

Thus

$$\begin{aligned} &\tau_r^0(p, q) + \tau_r^3(p, q) \\ &= \sqrt{\frac{2}{pr}} \exp\left(\frac{\pi i}{2r} S\left(\frac{q}{p}\right)\right) \\ &\quad \times \left( \sin\left(\frac{\pi}{pr}\right) \left(1 - \cos\left(2\pi \frac{q^*}{3}\right)\right) + i\sqrt{3} \cos\left(\frac{\pi}{pr}\right) \sin\left(2\pi \frac{q^*}{3}\right) \right) \\ &= i\sqrt{\frac{6}{pr}} \sin\left(2\pi \frac{q^*}{3}\right) (1 + c_1 r^{-1} + c_2 r^{-2} + \dots), \end{aligned}$$

where

$$c_1 = \frac{\pi \left(1 - \cos\left(2\pi \frac{q^*}{3}\right)\right)}{ip\sqrt{3} \sin\left(2\pi \frac{q^*}{3}\right)} + \frac{\pi i}{2} S\left(\frac{q}{p}\right) = 6\pi i \left( s(q, p) - \frac{\left(1 - \cos\left(2\pi \frac{q^*}{3}\right)\right)}{6p\sqrt{3} \sin\left(2\pi \frac{q^*}{3}\right)} \right).$$

Note that  $s(q^*, p) = s(q, p)$  and put

$$a(q) = \frac{(1 - \cos(2\pi \frac{q}{3}))}{54\sqrt{3} \sin(2\pi \frac{q}{3})}$$

and  $b(q) = s(q, p) - a(q)$ . We find that  $s(1, 9) = 14/27$  and  $s(2, 9) = 4/27$ , while  $a(2) = -a(1) = -1/54$ , so  $b(1) = \frac{27}{28}s(1, 9)$  while  $b(2) = \frac{9}{8}s(2, 9)$ . We conclude that the Casson–Walker invariant of  $L(p, q)$  is naturally associated with the trivial connection's contribution to the asymptotic expansion of  $\tau_r(L(p, q))$  and not with the full contribution to this asymptotics coming from the zero set of the Chern–Simons functional.

Next let us turn to the general Seifert fibered rational homology spheres. A Seifert manifold  $X = (\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  is a rational homology sphere iff  $\epsilon = 0$ ,  $g = 0$  and  $E \neq 0$  or  $\epsilon = n$  and  $g = 1$ . The Seifert manifold  $X$  is an integral homology sphere if and only if  $\epsilon = 0$ ,  $g = 0$  and  $|E|\mathcal{A} = 1$ . Here  $E = -b - \sum_{j=1}^n \frac{\beta_j}{\alpha_j}$  is the Seifert Euler number and  $\mathcal{A} = \prod_{j=1}^n \alpha_j$  as usual. In this paper we have not calculated the asymptotics for the cases where  $\epsilon = n$  and  $g = 1$ , except for the ones with zero or one exceptional fiber. Let us first consider the cases with base  $S^2$  and nonzero Seifert Euler number. For these cases we have by [Les, Proposition 6.1.1] that

$$\lambda_{\text{CW}}(X) = \frac{1}{12} \left( 3\text{sign}(E) - E - \sum_{i=1}^n S\left(\frac{\beta_i}{\alpha_i}\right) - \frac{1}{E} \left( 2 - n + \sum_{i=1}^n \frac{1}{\alpha_i^2} \right) \right), \quad (61)$$

where we have used the facts that  $|H_1(X; \mathbb{Z})| = |E|\mathcal{A}$  and that we for any rational homology sphere  $N$  have  $\lambda_{\text{CLW}}(N) = \frac{1}{2}|H_1(N; \mathbb{Z})|\lambda_{\text{CW}}(N)$ , see [Les, p. 13].

Under the identification of the moduli space  $\mathcal{M}_X$  of flat  $\mathrm{SU}(2)$ -connections on  $X$  by the set of  $\mathrm{SU}(2)$ -representations of  $\pi_1(X)$  moduli conjugation by  $\mathrm{SU}(2)$ , the trivial connection is identified by the trivial representation. Using notation from the previous section we note that  $\omega'(l, \underline{n}')[g_1, \dots, g_n]$  is the trivial representation if and only if  $l = 0$  and  $\underline{n}' = \underline{0}$  and in this case  $\omega'$  is independent of the  $g_j$ 's. Next assume that  $E \neq 0$  and let  $\rho = \rho(m, \underline{n})$ . We note that  $\rho(h) = \exp(\pi i z_{\mathrm{st}}(-m, \underline{n}))$ , where  $z_{\mathrm{st}}$  is given by (38). Hence  $\rho$  is trivial only if  $z_{\mathrm{st}}(-m, \underline{n}) \in 2\mathbb{Z}$  and in this case  $(-m, \underline{n}) \notin \mathcal{I}_1$ . It should be noted that for  $n \in \{0, 1, 2\}$ ,  $X$  is a lens space, so in that case  $\mathcal{M}_X$  is a finite discrete space. This is also the case for  $n = 3$ , but not for  $n \geq 4$ , see also our remarks at the end of Sec. 4.2. For  $n \geq 4$  it is not known whether the connected components of  $\mathcal{M}_X$  are parametrized by  $\mathcal{I}_1 \cup \mathcal{I}_2^a \cup \mathcal{I}_2^c$ ,  $\mathcal{I}_2^c$  being the set of points in  $\mathcal{I}_2^b$  satisfying (54).

In any case, let us calculate the contribution to the asymptotics of  $\tau_r(X)$  coming from the point  $(l, \underline{n}') = (0, \underline{0})$ . First note that there is no polar contribution, that is,  $Z_0^{(0, \underline{0})}(r)$  and  $Z_1^{(0, \underline{0})}(r)$  are both zero. The part of the large  $r$  asymptotics of  $\tau_r(X)$  associated to the zero point in  $\mathcal{I}_2$ , denoted  $\tau_r^0(X)$  in the following, is therefore given by

$$\frac{i^{n+1}}{2^n \sqrt{r}} \frac{e^{-i \frac{3\pi}{4} \mathrm{sign}(E)}}{\sqrt{A}} \exp \left( \frac{i\pi}{2r} \left[ 3\mathrm{sign}(E) - E - \sum_{j=1}^n S \left( \frac{\beta_j}{\alpha_j} \right) \right] \right) Z^0(X; r),$$

where  $Z^0(X; r)$  is the (not necessarily convergent) power series given as follows: Let  $k_0 = 0$  if  $n$  is odd and  $k_0 = 1$  otherwise. Moreover, let  $k_1 = (k_0 - n + 1)/2$ . Then  $Z^0(X; r) = \sum_{k=k_1}^{\infty} c_k r^{-k}$ , where

$$c_k = \frac{(-2i)^{n+1}}{\pi^{n-1}} \frac{e^{i \frac{\pi}{4} \mathrm{sign}(E)}}{\sqrt{2\pi|E|}} \frac{\Gamma(k + \frac{1}{2})}{k'!} \left( \frac{2i}{\pi E} \right)^k \partial_z^{(k')} \left\{ \left( \frac{\pi z}{\sin(\pi z)} \right)^{n-1} \prod_{j=1}^{n+1} \sin \left( \frac{\pi z}{\alpha_j} \right) \right\} \Big|_{z=0},$$

where  $k' = 2k + n - 1$  and  $\alpha_{n+1} = 1$ . Therefore

$$c_k = \frac{(-2i)^{n+1}}{\pi^{n-2}} \frac{e^{i \frac{\pi}{4} \mathrm{sign}(E)}}{\sqrt{2\pi|E|}} \frac{\Gamma(k + \frac{1}{2})}{k'!} \left( \frac{2i}{\pi E} \right)^k \partial_z^{(k'-1)} \left\{ \left( \frac{\pi z}{\sin(\pi z)} \right)^{n-2} \prod_{j=1}^n \sin \left( \frac{\pi z}{\alpha_j} \right) \right\} \Big|_{z=0}.$$

In general we have for  $m \in \{0, 1, \dots, n\}$  that

$$\begin{aligned} & \partial_z^{(l)} \left\{ \left( \frac{\pi z}{\sin(\pi z)} \right)^m \prod_{j=1}^n \sin \left( \frac{\pi z}{\alpha_j} \right) \right\} \Big|_{z=0} \\ &= \begin{cases} 0 & , l = 0, 1, \dots, m-1 \\ \frac{l! \pi^m}{(l-m)!} \partial_z^{(l-m)} \left\{ \frac{\prod_{j=1}^n \sin \left( \frac{\pi z}{\alpha_j} \right)}{\sin^m(\pi z)} \right\} \Big|_{z=0} & , l = m, m+1, \dots \end{cases} \end{aligned}$$

Since we have already treated the lens spaces, we can assume that  $n > 2$ . In that case  $k_1 < 0$  and  $k' - 1 = n - 2$  if and only if  $k = 0$ , so  $c_k = 0$  for  $k = k_1, \dots, -1$  and

$$\begin{aligned} c_k &= (-2i)^{n+1} \frac{e^{i \frac{\pi}{4} \mathrm{sign}(E)}}{\sqrt{2\pi|E|}} \frac{\Gamma(k + \frac{1}{2})}{(2k)!} \left( \frac{2i}{\pi E} \right)^k \partial_z^{(2k)} \left\{ \frac{\prod_{j=1}^n \sin \left( \frac{\pi z}{\alpha_j} \right)}{\sin^{n-2}(\pi z)} \right\} \Big|_{z=0} \\ &= (-2i)^{n+1} \frac{e^{i \frac{\pi}{4} \mathrm{sign}(E)}}{\sqrt{2|E|}} \frac{1}{k!} \left( \frac{i}{2\pi E} \right)^k \partial_z^{(2k)} \left\{ \frac{\prod_{j=1}^n \sin \left( \frac{\pi z}{\alpha_j} \right)}{\sin^{n-2}(\pi z)} \right\} \Big|_{z=0} \end{aligned}$$

for  $k = 0, 1, \dots$ . Therefore

$$\tau_r^0(X) = -i \operatorname{sign}(E) \sqrt{\frac{2}{r|E|\mathcal{A}}} \exp \left( \frac{i\pi}{2r} \left[ 3 \operatorname{sign}(E) - E - \sum_{j=1}^n S \left( \frac{\beta_j}{\alpha_j} \right) \right] \right) \tilde{Z}^0(X; r),$$

where  $\tilde{Z}^0(X; r) = \sum_{k=0}^{\infty} d_k r^{-k}$  with

$$d_k = \frac{1}{k!} \left( \frac{i}{2\pi E} \right)^k \partial_z^{(2k)} \left\{ \frac{\prod_{j=1}^n \sin \left( \frac{\pi z}{\alpha_j} \right)}{\sin^{n-2}(\pi z)} \right\} \Big|_{z=0}.$$

We have

$$f(z) := \frac{\prod_{j=1}^n \sin \left( \frac{\pi z}{\alpha_j} \right)}{\sin^{n-2}(\pi z)} = \frac{(\pi z)^2}{\mathcal{A}} \left( 1 + \frac{1}{6} \left( n - 2 - \sum_{j=1}^n \frac{1}{\alpha_j^2} \right) (\pi z)^2 + c(\pi z)^4 + \dots \right).$$

Thus  $f(0) = 0$ ,  $f^{(2)}(0) = 2\frac{\pi^2}{\mathcal{A}}$  and

$$f^{(4)}(0) = \frac{24\pi^4}{\mathcal{A}} \frac{1}{6} \left( n - 2 - \sum_{j=1}^n \frac{1}{\alpha_j^2} \right) = 12\pi^2 \frac{1}{6} \left( n - 2 - \sum_{j=1}^n \frac{1}{\alpha_j^2} \right) f^{(2)}(0).$$

Hence

$$\sum_{k=0}^{\infty} d_k r^{-k} = r d_1 \left( 1 + \frac{\pi i}{2E} \left( n - 2 - \sum_{j=1}^n \frac{1}{\alpha_j^2} \right) r^{-1} + a_2 r^{-2} + \dots \right)$$

and finally by (61)

$$\tau_r^0(X) = \frac{\pi}{|E|\mathcal{A}} \sqrt{\frac{2r}{|E|\mathcal{A}}} (1 + 6\pi i \lambda_{\text{CW}}(X) r^{-1} + b_2 r^{-2} + \dots).$$

Let us next consider the Seifert manifolds with base  $\mathbb{RP}^2$  and zero or one exceptional fiber. Except for the case  $(n; 1 \mid 0)$  all these manifolds are covered by the above calculation, since they are all homeomorphic to Seifert manifolds with base  $S^2$  and nonzero Seifert Euler number, see Sec. 4.3. Left is to consider the case  $X = (n; 1 \mid 0)$ . By (60) we have to find the first terms in the Taylor expansion of  $f(x) = \frac{\sin(x)}{2(1+\cos(x))}$ . But  $f$  is an odd function so we have  $f(x) = cx \sum_{j=0}^{\infty} c_{2j} x^{2j}$  for  $x \in ]-\pi, \pi[$ . Thus  $f\left(\frac{\pi}{r}\right) = c \frac{\pi}{r} (1 + c_1 r^{-1} + \pi^2 c_2 r^{-2} + \dots)$  for  $r \geq 4$ , where  $c_1 = 0$ . But  $\lambda_{\text{CW}}(\mathbb{RP}^3) = 0$ , thus  $\lambda_{\text{CW}}(X) = 0$ , so again we find that the part of the expansion of  $\tau_r(X)$  related to the trivial connection has a form  $ar^d (1 + 6\pi i \lambda_{\text{CW}}(X) r^{-1} + b_2 r^{-2} + \dots)$ .

The Casson–Walker invariant  $\lambda_{\text{CW}}$  has been extended by Lescop to an invariant  $\lambda_{\text{CLW}}$  of all closed oriented 3-manifolds, cf. [Les]. We note that  $\tau_r(S^2 \times S^1) = 1 = 1 + c_1 r^{-1} + c_2 r^{-2} + \dots$ , where  $c_l = 0$  for all  $l = 1, 2, \dots$ . We have previously seen that  $S^2 \times S^1$  has an infinite number of Seifert fibered structures. Thus  $S^2 \times S^1 = (o; 0 \mid -1; (\alpha, \beta), (\alpha, \alpha - \beta))$  for any  $\alpha \in \mathbb{Z}_{\geq 2}$  and any  $\beta \in \{1, 2, \dots, \alpha - 1\}$  coprime to  $\alpha$ . Using this together with [Les, Proposition 6.1.1], we find that  $\lambda_{\text{CLW}}(S^2 \times S^1) = -1/12$ . The moduli space  $\mathcal{M}_{S^2 \times S^1}$  is topologically a closed interval since  $\pi_1(S^2 \times S^1) = \mathbb{Z}$ . (An element  $\rho \in \operatorname{Hom}(\mathbb{Z}, \operatorname{SU}(2))$  is determined by  $\rho(1)$ . By conjugation we can assume that  $\rho(1) = e^{i\theta}$ ,  $\theta \in [0, \pi]$ .) In particular, the moduli space  $\mathcal{M}_{S^2 \times S^1}$  is connected. Thus it seems that Lescop’s extension of the Casson–Walker invariant to 3-manifolds not being rational homology spheres is not part of the quantum  $\operatorname{SU}(2)$ -invariants in the same way as the Casson–Walker invariants of rational homology 3-spheres. At least this extension does not seem

to be proportional to any of the coefficients in the asymptotic expansion of the quantum invariant as the simple example  $S^2 \times S^1$  reveals.

## 5. PROOF OF THEOREM 4.1

The proof of Theorem 4.1 is rather long and technical but the single steps in the proof use elementary analysis. To streamline the proof we will in this section emphasize the main ideas in the arguments and defer technical details to a number of appendices. The proof of Theorem 4.4, which uses the same ideas as the proof of Theorem 4.1, is much shorter and is given in Sect. 6.

In the introduction we gave an outline of the proof. Let us recall the basic ideas. Let  $X$  be a Seifert manifold described by (normalized) Seifert invariants  $(\epsilon; g \mid b; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n))$  or non-normalized Seifert invariants  $\{\epsilon; g; (\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)\}$ . The first problem we encounter when calculating the large  $r$  asymptotics of  $r \mapsto Z(X; r)$  is that this function is given by a sum in which both the terms and the summation range depend on  $r$ , cf. Theorem 3.1. This problem is solved by using certain symmetries to replace  $\sum_{\gamma=1}^{r-1}$  by another sum of the form  $\sum_{\gamma \in \mathbb{Z}}$ . After this manoeuvre we use Poisson's summation formula to change the expression for  $Z(X; r)$  to an infinite sum of integrals. Apparently all this seems to complicate things. However, we obtain the big advantage that the large  $r$  asymptotics of the integrals in the thus obtained sum can be calculated with a strong method called the steepest descent method. Roughly speaking this will express  $Z(X; r)$  as an infinite sum of main contributions plus an infinite sum of remainder terms. The final step will be to rewrite the infinite sum of main contributions as a finite sum in which the summation range does not depend on  $r$  and to show that the infinite sum of remainder terms is small compared to this finite sum in the large  $r$  limit.

We want to generalize the situation. Therefore, let us first consider the function  $Z(X; r)$  in (36) in greater detail. As mentioned above the first step is to rewrite this expression so that  $r$  only appears in the summands and not in the summation range. To this end we need to extend the function  $h$  in (37) to  $\mathbb{Z}$ . There is of course no problem in doing this by using the expression (37). However, we will need that the extended function  $h$  satisfies the following symmetries

$$\begin{aligned} h(r\gamma) &= 0, \\ h(-\gamma) &= (-1)^n h(\gamma), \\ h(\gamma + 2r) &= h(\gamma) \end{aligned} \tag{62}$$

valid for all  $\gamma \in \mathbb{Z}$ . To establish these symmetries we need to examine the proof of [Ha2, Theorem 8.4]. Let us note here that the first symmetry  $h(r\gamma) = 0$ ,  $\gamma \in \mathbb{Z}$ , is not satisfied if  $n = 0$ . When this identity is needed we will force it to be true also in case  $n = 0$ , see below (69). The two other symmetries are trivially satisfied when  $n = 0$ .

Assume  $n > 0$ . The Rademacher Phi function  $\Phi$  is defined on  $\text{PSL}(2, \mathbb{Z}) = \text{SL}(2, \mathbb{Z})/\{\pm 1\}$  by

$$\Phi \begin{bmatrix} a & b \\ c & d \end{bmatrix} = \begin{cases} \frac{a+d}{c} - S\left(\frac{d}{c}\right) & , c \neq 0, \\ \frac{b}{d} & , c = 0, \end{cases} \tag{63}$$

where the Dedekind symbol  $S(d/c)$  is given by (34). By the proof of [Ha2, Theorem 8.4] the function  $h$  is given by

$$h(\gamma) = \kappa \prod_{i=1}^n (\tilde{M}_i)_{\gamma,1} \tag{64}$$

with

$$\kappa = \sqrt{\mathcal{A}} (2r)^{n/2} i^{-n} \exp \left( \frac{i\pi}{4} \sum_{j=1}^n \Phi(M_j) \right) \exp \left( -\frac{i\pi}{2r} \sum_{j=1}^n \frac{\rho_j}{\alpha_j} \right),$$

where  $M_i = \begin{pmatrix} -\beta_i & -\sigma_i \\ \alpha_i & \rho_i \end{pmatrix}$ ,  $i = 1, \dots, n$ . Here  $A \mapsto \tilde{A}$  is a unitary  $(r-1)$ -dimensional representation of  $PSL(2, \mathbb{Z})$ . By unitarity we have  $\tilde{A}_{j,k} = \overline{(\tilde{A}^{-1})_{k,j}}$  for  $j, k \in \{1, 2, \dots, r-1\}$ , where  $\bar{\cdot}$  is complex conjugation. By using this with  $A = M_i$  together with the formula for  $(\tilde{M}_i^{-1})_{1,\gamma}$  given in [J1, Propositions 2.7 (a) and 2.8] we get

$$\begin{aligned} (\tilde{M}_i)_{\gamma,1} &= i \frac{1}{\sqrt{2r\alpha_i}} e^{-\frac{i\pi}{4}\Phi(M_i)} \\ &\times \sum_{\mu=\pm 1} \sum_{m=0}^{\alpha_i-1} \mu \exp \left( \frac{i\pi}{2r\alpha_i} [-\beta_i \gamma^2 - 2\gamma(2rm + \mu) + \rho_i(2rm + \mu)^2] \right), \end{aligned} \quad (65)$$

and inserting this expression for  $(\tilde{M}_i)_{\gamma,1}$  into (64) leads to (37).

Now, to prove (62), we only need to establish the same symmetries for the functions  $\gamma \mapsto (\tilde{M}_i)_{\gamma,1}$  (extended to  $\mathbb{Z}$ ). To do this we make a little simple observation. The group  $SL(2, \mathbb{Z})$  is generated by the two matrices

$$\Xi = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \Theta = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \quad (66)$$

Any  $A \in SL(2, \mathbb{Z})$  can be written  $A = B\Xi$ , where  $B = A\Xi^{-1} = -A\Xi$ . In particular

$$\tilde{A}_{j,k} = \sum_{l=1}^{r-1} \tilde{B}_{j,l} \tilde{\Xi}_{l,k}, \quad (67)$$

where

$$\tilde{\Xi}_{l,k} = \sqrt{\frac{2}{r}} \sin \left( \frac{\pi l k}{r} \right), \quad l, k = 1, \dots, r-1.$$

This expression is valid for all  $k \in \mathbb{Z}$ , so we can use (67) to extend the function  $k \mapsto \tilde{A}_{j,k}$  from  $\{1, \dots, r-1\}$  to  $\mathbb{Z}$ . We can also use the explicit expression for  $k \mapsto \tilde{A}_{j,k}$  given by [J1, Propositions 2.7 (a) and 2.8] to extend this function to  $\mathbb{Z}$ , but it follows by the proof of these propositions in [J1] that these two extensions coincide. By (67) we immediately get that

$$\begin{aligned} \tilde{A}_{j,rk} &= 0, \\ \tilde{A}_{j,-k} &= -\tilde{A}_{j,k}, \\ \tilde{A}_{j,k+2r} &= \tilde{A}_{j,k} \end{aligned} \quad (68)$$

for all  $k \in \mathbb{Z}$  and  $j \in \{1, \dots, r-1\}$ , giving the needed symmetries for the entries  $(\tilde{M}_i)_{\gamma,1}$ . In the above kind of argument we do of course not need to use unitarity of the representation  $A \mapsto \tilde{A}$ . The reason for using this unitarity is that it leads to an expression for  $h(\gamma)$  which is slightly better to work with in the proof of Theorem 1.4.

Assume still that  $n > 0$ . By introducing a parameter  $\xi > 0$  we get by (36) and (62) that

$$Z(X; r) = \frac{1}{2} \left( \frac{i}{2} \right)^n \lim_{\xi \rightarrow 0^+} \sum_{\gamma=-r+1}^r P(\gamma) H(\gamma, \xi), \quad (69)$$

where

$$H(\gamma, \xi) = \frac{e^{\pi i a_\epsilon g \gamma} h(\gamma)}{\sin^{n+a_\epsilon g-2} \left( \frac{\pi}{r} (\gamma - i\xi) \right)}$$

and where  $P(\gamma) = (-1)^{a_\epsilon g}$  for  $\gamma = -r + 1, \dots, -1$ ,  $P(\gamma) = 1$  for  $\gamma = 1, \dots, r$ . Moreover we let  $P(x) = (P(x-) + P(x+))/2$ ,  $x = 0, r$ , where  $P(x\pm) = \lim_{t \rightarrow x\pm} P(t)$ . If we multiply  $H(\gamma, \xi)$  by

$$\frac{\sin\left(\frac{\pi}{r}\gamma\right)}{\sin\left(\frac{\pi}{r}(\gamma - i\xi)\right)}$$

in case  $n = 0$ , (69) also holds in that case. For later use we extend  $P$  not only to  $\mathbb{Z}$  but to all of  $\mathbb{R}$  by letting  $P(\gamma) = (-1)^{a_\epsilon g}$  for  $-r < \gamma < 0$  and  $P(\gamma) = 1$  for  $0 < \gamma < r$  and letting  $P$  be periodic with a period of  $2r$ . Note that for  $a_\epsilon g$  even (that is for  $\epsilon = 0$  or  $g$  even)  $P$  is identically one. The parameter  $\xi$  can of course be ignored in cases where  $n + a_\epsilon g - 2 \leq 0$ . To continue we use the following formula which is a special case of Lemma 7.1. If  $f : \mathbb{Z} \rightarrow \mathbb{C}$  is a function with a period of  $N$  then

$$\sum_{k=0}^{N-1} f(k) = N \lim_{\varepsilon \rightarrow 0+} \sqrt{\varepsilon} \sum_{k \in \mathbb{Z}} e^{-\pi \varepsilon k^2} f(k). \quad (70)$$

By applying this to the sum (69) we obtain

$$Z(X; r) = r \left(\frac{i}{2}\right)^n \lim_{\xi \rightarrow 0+} \lim_{\varepsilon \rightarrow 0+} \sqrt{\varepsilon} \sum_{\gamma \in \mathbb{Z}} e^{-\pi \varepsilon \gamma^2} P(\gamma) H(\gamma, \xi).$$

We can exchange the order of the two limits. To see this, simply use (70) with  $f(k) = P(k) \lim_{\xi \rightarrow 0+} H(k, \xi)$  to get

$$Z(X; r) = r \left(\frac{i}{2}\right)^n \lim_{\varepsilon \rightarrow 0+} \sqrt{\varepsilon} \sum_{\gamma \in \mathbb{Z}} e^{-\pi \varepsilon \gamma^2} P(\gamma) \lim_{\xi \rightarrow 0+} H(\gamma, \xi).$$

By uniform convergence of the sum  $\sum_{\gamma \in \mathbb{Z}} e^{-\pi \varepsilon \gamma^2} P(\gamma) H(\gamma, \xi)$  with respect to  $\xi$  on an interval of the form  $]0, \xi_0]$ ,  $\xi_0 > 0$ , we then have that

$$Z(X; r) = r \left(\frac{i}{2}\right)^n \lim_{\varepsilon \rightarrow 0+} \sqrt{\varepsilon} \lim_{\xi \rightarrow 0+} \sum_{\gamma \in \mathbb{Z}} e^{-\pi \varepsilon \gamma^2} P(\gamma) H(\gamma, \xi). \quad (71)$$

The expression (37) allows us to consider  $h$  as an entire function. In particular we can consider  $\gamma \mapsto H(\gamma, \xi)$  as a meromorphic function on  $\mathbb{C}$  with a countable number of isolated poles. To proceed we change (71) to a sum of integrals by using the Poisson summation formula

$$\sum_{k \in \mathbb{Z}} \varphi(k) = \sum_{m \in \mathbb{Z}} \int_{-\infty}^{\infty} e^{2\pi i m x} \varphi(x) dx. \quad (72)$$

By a result of Zygmund [Z, p. 68] this formula is valid for every function  $\varphi$  which is absolutely integrable over  $\mathbb{R}$ , of bounded variation, and satisfies  $2\varphi(x) = \varphi(x+) + \varphi(x-)$  for all  $x \in \mathbb{R}$ . We use Poisson's formula with  $\varphi(\gamma) = e^{-\pi \varepsilon \gamma^2} P(\gamma) H(\gamma, \xi)$  (with fixed  $\xi$ ). By (37) and the above we get

$$Z(X; r) = r \left(\frac{i}{2}\right)^n \lim_{\delta \rightarrow 0+} \sqrt{\delta} \lim_{\eta \rightarrow 0+} \sum_{m \in \mathbb{Z}} \sum_{\underline{\mu} \in \{\pm 1\}^n} \sum_{\underline{n} = \underline{0}}^{\underline{\alpha}-1} g(\underline{\mu}, \underline{n}) \int_{-\infty}^{\infty} \kappa(z) e^{r Q_m(z)} dz, \quad (73)$$

where

$$\begin{aligned}
g(\underline{\mu}, \underline{n}) &= \left( \prod_{j=1}^n \mu_j \right) \exp \left( 2\pi i \sum_{j=1}^n \frac{\rho_j}{\alpha_j} [r n_j^2 + \mu_j n_j] \right), \\
\kappa(z) &= \kappa_\eta(z; \underline{\mu}, \delta) = e^{-\pi \delta z^2} P(rz) \frac{\exp \left[ -i\pi \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right]}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))}, \\
Q_m(z) &= Q_m(z; \underline{n}) = 2\pi i \left( \frac{a_\epsilon g}{2} + m - \sum_{j=1}^n \frac{n_j}{\alpha_j} \right) z + \frac{i\pi}{2} E z^2.
\end{aligned}$$

(Put  $z = \gamma/r$ ,  $\delta = \varepsilon r^2$  and  $\eta = \xi/r$ .) In case  $n = 0$  the function  $\kappa(z)$  contains an extra factor  $\sin(\pi z)/\sin(\pi(z-i\eta))$ . But

$$\sin(\pi z) = \frac{i}{2} \sum_{\mu \in \{\pm 1\}} \mu \exp(-i\pi \mu z),$$

so formally we can treat the case  $n = 0$  as the case  $n = 1$  with  $(\alpha_1, \beta_1) = (1, 0)$

We note that it was necessary in the process of changing the expression (36) to (73) to involve the sum  $\sum_{\mu \in \{\pm 1\}^n} \sum_{\underline{n}=\underline{0}}^{\underline{\alpha}-1}$ . This was needed for establishing (some of) the properties (62). Note that  $z \mapsto P(rz)$  is independent of  $r$ .

**Remark 5.1.** A remark is appropriate here. It is well-known that the Poisson formula (72) is valid for functions in the Schwartz space  $\mathcal{S}(\mathbb{R})$  of smooth functions, that together with their derivatives are rapidly decreasing at infinity, see e.g. [Ho]. If  $a_\epsilon g$  is even then  $P$  is identically 1 and  $\varphi(\gamma) = e^{-\pi \varepsilon \gamma^2} H(\gamma, \xi)$  is in  $\mathcal{S}(\mathbb{R})$  since  $\gamma \mapsto H(\gamma, \xi)$  is periodic by (62). If  $a_\epsilon g$  is odd it is not clear how many times  $\gamma \mapsto P(\gamma)H(\gamma, \xi)$  is differentiable in a point  $\gamma \in r\mathbb{Z}$ . We can, however, choose  $P$  to be smooth also for  $a_\epsilon g$  odd. In this case we can simply let  $P$  be a smooth function equal to 1 on  $[3/4, r - 3/4]$ , equal to 0 on  $[0, 1/4]$  and on  $[r - 1/4, r]$ , and equal to  $-P(-\gamma)$  on  $[-r, 0]$ . Finally we extend  $P$  to all of  $\mathbb{R}$  by letting it have the period  $2r$ . The problem with this approach is that  $z \mapsto P(rz)$  so defined depends on  $r$ .

For the method used in the following we will need to extend  $P$  to an entire (or at least meromorphic) function with certain properties, see above Remark 5.2. It is not clear at the moment for the author if such a  $P$  exists, and therefore we restrict to the case  $a_\epsilon g$  even, i.e.  $\epsilon = 0$  or  $g$  even. The calculations presented in this paper were first done for the case  $\epsilon = 0$  where  $a_\epsilon g$  is always even, and first later the author started to consider the case  $\epsilon = n$ . It was then realized that the calculations could only be carried through as they stand for half of the Seifert manifolds with nonorientable base, namely the ones with a base with even genus. We note that one can probably calculate the asymptotics of  $Z(X; r)$  by other means than applied in this paper, see e.g. [Ro1] and [Ro3, Appendix] for some suggestions. It should be possible to use one of these methods to handle the case of  $a_\epsilon g$  odd, but we will defer that to another paper.

We approximate the integrals  $\int_{-\infty}^{\infty} \kappa(z) e^{rQ_m(z)} dz$  in (73) by the steepest descent method. This is a general method for obtaining asymptotic expansions of contour integrals of the form

$$I(t) = \int_C g(z) e^{tf(z)} dz$$

in the limit  $t \rightarrow \infty$ , where  $f(z)$  and  $g(z)$  are analytic functions on some domain  $\Omega$  in the complex plane, both independent of the real positive parameter  $t$ , and  $C$  is a contour inside  $\Omega$ . Actually we allow  $g(z)$  to have poles in  $\Omega$  away from the contours involved. The basic idea is to deform

$C$  inside  $\Omega$  into a new path of integration  $C'$  so that the following conditions hold:

- i) The path  $C'$  passes through one or more of the stationary points of  $f$  (that is the zeroes of  $f'(z)$ );
- ii) The imaginary part of  $f(z)$  is constant on  $C'$ .

We will only need to consider the case of non-degenerate stationary points of  $f$ , i.e. stationary points  $p$  such that  $f''(p) \neq 0$ . In that case one finds that there is a unique contour  $C'$  through  $p$  satisfying ii) above and satisfying that  $\operatorname{Re}(f(p) - f(z)) < 0$  for all  $z$  on  $C' \setminus \{p\}$  near  $p$ . This contour is called the steepest descent contour through  $p$ . It's argument is  $(\pi - \arg(f''(p)))/2$ , see e.g. [BH, Theorem 7.1] or [B, Sec. 5.4]. To determine the asymptotic expansion of  $I(t)$  in the limit  $t \rightarrow \infty$  we have to calculate the large  $t$  asymptotics of the new contour integral

$$J(t) = \int_{C'} g(z) e^{tf(z)} dz.$$

The difference between  $I(t)$  and  $J(t)$  can be zero or contain a sum of residue contributions, coming from the poles of  $g(z)$  crossed in the process of deforming the contour from  $C$  to  $C'$ . Moreover, this difference can contain contributions given by integrals along contours connecting the contours  $C$  and  $C'$ , but these contributions should be  $o(t)$  compared to  $J(t)$  in the limit  $t \rightarrow \infty$ . The integral  $J(t)$  will for quite general  $f(z)$  and  $g(z)$  be of a form for which the Laplace method can be used to calculate the large  $t$  asymptotics. In our case these integrals will however be so simple that a general description of the Laplace method is not relevant. Moreover, we will not go into details about why the steepest descent contour  $C'$  is a particularly 'good' contour when it comes to finding the large  $t$  asymptotics of  $I(t)$ . From a technical point of view a main reason is, as already pointed out, that the Laplace method is applicable to the new integral  $J(t)$  along  $C'$ . We refer to [Wo, pp. 84-90], [BH, pp. 262-268] and [B, Chap. 5] for further details.

In case  $f$  has more than one stationary point one has normally to determine the contribution to the large  $t$  asymptotics of  $I(t)$  coming from each stationary point by using a 'short' steepest descent contour through each such point. One should of course also secure that it is possible to deform the original contour  $C$  into the steepest descent contour within the domain  $\Omega$ .

In our situation  $t = r$ ,  $f(z) = Q_m(z)$ ,  $g(z) = \kappa(z)$ ,  $\Omega = \mathbb{C}$ , and the contour  $C$  is the real axes. If the Seifert Euler number  $E = 0$  the functions  $Q_m$  do not have stationary points so in this case the steepest descent method is not applicable. The case  $E = 0$  will be taken care of in Sec. 6 as already pointed out earlier. If  $E \neq 0$  the phase function  $Q_m$  has exactly one stationary point

$$z_{\text{st}} = -\frac{2}{E} \left( \frac{a_e g}{2} + m - \sum_{j=1}^n \frac{n_j}{\alpha_j} \right)$$

which is non-degenerate.

**5.1. Calculation of asymptotics. A general case.** To make the calculations useable for other situations and for shortening notation we will generalize the situation above without making things more difficult. To be more concrete we will study functions of the form

$$Z(r) = \sum_{\gamma=1}^{r-1} \frac{h(\gamma)}{\sin^k\left(\frac{\pi}{r}\gamma\right)}, \quad (74)$$

where  $k \in \mathbb{Z}$  and

$$h(\gamma) = \sum_{\lambda \in \Lambda} g(\lambda) p(\gamma) \exp\left(i \left[ \frac{A}{r} \gamma^2 + B\gamma \right]\right). \quad (75)$$



We assume that  $A$  and  $B$  are real and independent of  $r$ . Moreover, we assume that  $g$  is independent of  $\gamma$  and that  $p$  is an entire function. The index set  $\Lambda$  is assumed to be a finite subset of  $\mathbb{R}^e$  for some  $e \in \mathbb{Z}_{>0}$ . We also assume that  $h$  satisfies (62) with  $n = k$  for all  $\gamma \in \mathbb{Z}$ . We will allow  $g$  to depend on  $r$  and  $A, B$ , and  $p$  to depend on  $\lambda$ . By (70) we get

$$Z(r) = r \lim_{\varepsilon \rightarrow 0_+} \sqrt{\varepsilon} \lim_{\xi \rightarrow 0_+} \sum_{\gamma \in \mathbb{Z}} e^{-\pi \varepsilon \gamma^2} \frac{h(\gamma)}{\sin^k\left(\frac{\pi}{r}(\gamma - i\xi)\right)}.$$

(We can exchange the order of the limits  $\lim_{\varepsilon \rightarrow 0_+}$  and  $\lim_{\xi \rightarrow 0_+}$  in (76). This follows exactly as in the special case where  $Z(r) = Z(X; r)$ ). The function  $x \mapsto e^{-\pi \varepsilon x^2} \frac{h(x)}{\sin^k\left(\frac{\pi}{r}(x - i\xi)\right)}$  belongs to  $\mathcal{S}(\mathbb{R})$  since  $x \mapsto \frac{h(x)}{\sin^k\left(\frac{\pi}{r}(x - i\xi)\right)}$  is periodic. Therefore

$$Z(r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} \sum_{m \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) \int_{-\infty}^{\infty} f(z; m, \lambda) dz \quad (76)$$

by the Poisson formula (72), where

$$f(z; m, \lambda) = e^{-\pi \delta z^2} q(z) \frac{e^{rQ_m(z)}}{\sin^k(\pi(z - i\eta))}, \quad (77)$$

with  $q(z) = q(z; \lambda) = p(rz)$  and  $Q_m(z) = Q(z) + 2\pi imz$ , where  $Q(z) = Q(z; \lambda) = i(Az^2 + Bz)$ . In case  $k \leq 0$  we can and will replace  $1/\sin^k(\pi(z - i\eta))$  in  $f(z; m, \lambda)$  by  $\sin^{|k|}(\pi z)$  and remove  $\lim_{\eta \rightarrow 0_+}$ . (Note that we do not lose any generality by not allowing the polynomial  $Q$  to have a constant term. Such a constant term  $c$  would just produce an extra factor  $e^{irc}$  in  $g$ .) From the first identity in (62) we have

$$\sum_{\lambda \in \Lambda} g(\lambda) q(z) e^{rQ(z)} = 0 \quad (78)$$

for all  $z \in \mathbb{Z}$ .

In case  $Z(r) = Z(X; r)$  (with  $a_\epsilon g$  even) we have by (73) that

$$\begin{aligned} \Lambda &= \{\pm 1\}^n \times \mathcal{S}, \\ A &= \frac{\pi}{2} E = -\frac{\pi}{2} \left( \beta_0 + \sum_{j=1}^n \frac{\beta_j}{\alpha_j} \right), \\ B &= B(\underline{n}) = -2\pi \sum_{j=1}^n \frac{n_j}{\alpha_j}, \\ k &= n + a_\epsilon g - 2, \\ g(\underline{\mu}, \underline{n}) &= \left( \frac{i}{2} \right)^n \left( \prod_{j=1}^n \mu_j \right) \exp \left( 2\pi i \sum_{j=1}^n \frac{\rho_j}{\alpha_j} [rn_j^2 + \mu_j n_j] \right), \\ q(z) &= q(z; \underline{\mu}) = \exp \left( -i\pi \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right) \end{aligned} \quad (79)$$

for  $(\underline{\mu}, \underline{n}) \in \Lambda$ , where  $\mathcal{S}$  is given by (41).

Let us return to the general case. We will need to make a series of assumptions on  $q$ . Firstly, we will assume that  $q$  is independent of  $r$  (or equivalently, that  $z \mapsto p(rz)$  is independent of  $r$ ). This assumption is made to make the steepest descent method applicable to the asymptotic analysis of integrals  $\int_{-\infty}^{\infty} f(z; m, \lambda) dz$ . Secondly, we need to make some assumptions of a technical nature.

The need for these will only be transparent within the proofs of results to follow. In general, they are made to secure convergency of certain infinite sums. First of all we assume that there exist real positive numbers  $a_n, b_n$  such that

$$\left| \frac{d^n q(z; \lambda)}{dz^n} \right| \leq b_n e^{a_n |z|} \quad (80)$$

for all  $z \in \mathbb{C}$ , all  $\lambda \in \Lambda$ , and  $n = 0, 1, \dots$ . Moreover, we assume that  $q$  is periodic along each line parallel to the real line. In fact we assume that there exists an integer  $m_q > 0$  such that

$$q(z + m_q; \lambda) = q(z; \lambda) \quad (81)$$

for all  $z \in \mathbb{C}$  and all  $\lambda \in \Lambda$ . It will follow that we can relax the assumption (80) slightly by replacing  $e^{a_n |z|}$  by  $e^{a_n |z|^\tau}$  with  $\tau \in [1, 2[$ , but since the above suffices for our purpose we will continue by using (80). In the case  $Z(r) = Z(X; r)$  the above conditions are satisfied, see (79).

**Remark 5.2.** This remark is a continuation of Remark 5.1. In the case where  $Z(r) = Z(X; r)$  and  $a_\epsilon g$  is odd  $q$  contains an extra factor  $P(rz)$ , where  $P$  is the function in (69). To include this case in the following calculations one will need to extend  $P$  to an entire function in such a way that the above assumptions on  $q$  are not violated. It is not clear to the author if this can be done.

By the formulae  $\sin(z) = (e^{iz} - e^{-iz})/(2i)$  and  $\cos(z) = (e^{iz} + e^{-iz})/2$  we see that both  $|\sin(z)|$  and  $|\cos(z)|$  are less than or equal to  $e^{|z|}$  for all  $z \in \mathbb{C}$ . For  $k \leq 0$  we can therefore let  $1/\sin^k(\pi z)$  be included in  $q(z)$  without violating the above assumptions on  $q$ .

In the following we will suppress  $\lambda$  in many of the expressions. One should keep in mind that all functions depending on  $A, B$  and/or  $q$  may depend on  $\lambda$ . The calculation of the large  $r$  asymptotics of  $Z(r)$  is divided naturally into several parts. To make it possible to follow the rather long computation we have tried to stress this by dividing the calculation into several subsections.

**5.1.1. A partition of  $Z(r)$  using the steepest descent method.** In the remaining part of Sec. 5 we assume that  $A \neq 0$ . The case  $A = 0$  will be handled in Sec. 6. We use the method of steepest descent to calculate the large  $r$  asymptotics of the integrals  $\int_{-\infty}^{\infty} f(z; m) dz$  in (76). The polynomial  $Q_m(z; \lambda)$  has only one critical point

$$z_{\text{st}}(m, \lambda) = -\frac{2\pi m + B}{2A} \quad (82)$$

which is non-degenerate. The steepest descent contour, denoted  $C_{\text{sd}}(m, \lambda)$ , is the contour

$$\text{Im}(z) = \text{sign}(A)(\text{Re}(z) - z_{\text{st}}(m, \lambda)). \quad (83)$$

We orient  $C_{\text{sd}}$  by starting at  $-\infty e^{\frac{i\pi}{4} \text{sign}(A)} + z_{\text{st}}$ . In the process of deforming the real axes into  $C_{\text{sd}}$  the integration contour crosses (for  $k > 0$ ) those poles

$$z_l(\eta) = l + i\eta, \quad l \in \mathbb{Z}, \quad (84)$$

for which

$$\text{sign}(A)(l - z_{\text{st}}) > \eta. \quad (85)$$

**Remark 5.3.** It is possible in parts of the calculation of the large  $r$  asymptotics of  $Z(r)$  in (76) to let  $Q$  be of a more general form, in particular if no singularities are present, i.e. if  $k \leq 0$ . One can e.g. assume that  $Q$  is an arbitrary analytic function satisfying that  $Q_m(z) = 2\pi i m z + Q(z)$  has exactly one stationary point  $z_m$  for each  $m$ . Moreover, we assume that  $z_m$  is nondegenerate.

For the stationary point  $z_m$  we deform the integration contour in  $\int_{-\infty}^{\infty} f(z; m) dz$  from the real axes to the line  $\gamma_m(t) = z_m + \alpha_m t$ ,  $t \in \mathbb{R}$ , where

$$\alpha_m = \exp\left(\frac{i\pi}{2} - \frac{i}{2}\text{Arg}(Q_m''(z_m))\right).$$

We then have

$$Q_m(\gamma_m(t)) = Q_m(z_m) + \frac{1}{2}Q_m''(z_m)\alpha_m^2 t^2 + \dots,$$

where  $Q_m''(z_m)\alpha_m^2 = -|Q_m''(z_m)| < 0$ . In case of polar contributions, i.e. in case  $k > 0$  we will also assume that  $\gamma_m$  is not parallel with the real axes, i.e. we assume that  $Q_m''(z_m)$  is not real and negative. This assumption is imposed to avoid problems with the sum of residue contributions (see Lemma 5.5 below). The integrals  $\int_{\gamma_m} f(z; m) dz$  can now be handled by using the Laplace method as illustrated in [B, Sec. 4.4]. There are several reasons why we will not do the calculations in this generality. First of all it will put us too far away from the example of main interest in this paper, the case  $Z(r) = Z(X; r)$ . A more serious reason is however that to be able to carry out the analysis we will have to impose too many assumptions. Some of these assumptions will then in concrete examples be hard to check (or at least these checks will demand long computations), so one does not gain very much. We have found that the expression (77) is a good compromise between generality and ability to carry through the analysis without too many assumptions.

We will need to impose a periodicity assumption on the function  $z_{\text{st}} : \mathbb{Z} \times \Lambda \rightarrow \mathbb{R}$ . To be specific we assume that there exists an integer  $H > 0$  such that

$$z_{\text{st}}(m + H, \lambda) - z_{\text{st}}(m, \lambda) = -\frac{\pi}{A}H \in \mathbb{Z} \quad (86)$$

for all  $(m, \lambda) \in \mathbb{Z} \times \Lambda$ . This implies that the set  $\{z_{\text{st}}(m, \lambda) \pmod{\mathbb{Z}} \mid (m, \lambda) \in \mathbb{Z} \times \Lambda\}$  is finite. Note that (86) is satisfied if and only if

$$A = A(\lambda) = \pi \frac{H}{P} \quad (87)$$

with  $P = P(\lambda) \in \mathbb{Z} \setminus \{0\}$  for all  $\lambda \in \Lambda$ . In the following we assume that  $A$  is given by (87) and reserve the symbols  $H$  and  $P$  for this. By (86) there exists an  $\eta_0 \in ]0, 1/2[$  such that

$$z_{\text{st}}(m, \lambda) \notin \mathbb{Z} + ([-\eta_0, \eta_0] \setminus \{0\}) \quad (88)$$

for all  $(m, \lambda) \in \mathbb{Z} \times \Lambda$ . For many of the next results we only have to assume (88) but later on we will have to assume the stronger (86). In parts of the calculation we will also need to assume that

$$B = B(\lambda) \in \pi\mathbb{Q} \quad (89)$$

for all  $\lambda \in \Lambda$ . When  $Z(r) = Z(X; r)$ , (86) and (89) are satisfied by (79).

Let us introduce some notation used in the remaining part of this paper. The steepest descent contour  $C_{\text{sd}}(m, \lambda)$ ,  $(m, \lambda) \in \mathbb{Z} \times \Lambda$ , is parametrized by

$$\gamma_{(m, \lambda)}(t) = te^{\frac{i\pi}{4}\text{sign}(A)} + z_{\text{st}}(m, \lambda). \quad (90)$$

We let

$$\kappa_{\eta}(z; \lambda, \delta) = e^{-\pi\delta z^2} \frac{q(z; \lambda)}{\sin^k(\pi(z - i\eta))}, \quad (91)$$

where  $\delta$  and  $\eta$  are non-negative parameters. We note that

$$e^{rQ_m(\gamma_{(m, \lambda)}(t))} = e^{-irAz_{\text{st}}^2(m, \lambda)} e^{-r|A|t^2}. \quad (92)$$

Let

$$W = \{ (m, \lambda) \in \mathbb{Z} \times \Lambda \mid z_{\text{st}}(m, \lambda) \notin \mathbb{Z} \}, \quad (93)$$

and put

$$A_0 = \min\{ |A(\lambda)| \mid \lambda \in \Lambda \}. \quad (94)$$

The open disk in the complex plane with centre  $z$  and radius  $\rho > 0$  is denoted  $D(z, \rho)$ , and the punctured disk  $D(z, \rho) \setminus \{z\}$  is denoted  $D'(z, \rho)$ .

The next lemma, Lemma 5.4, is necessary because we deal with infinite contours so we have to check that the integrals along the steepest descent contours are convergent. Since the real axes and the steepest descent contour are not attached to each other in the ‘end points’ we also have to check that there are no contributions from integrals along contours connecting the real axes to  $C_{\text{sd}}(m)$  at respectively minus and plus infinity.

**Lemma 5.4.** *Let  $\eta_0 > 0$  be as in (88) and let  $(m, \lambda) \in \mathbb{Z} \times \Lambda$  be arbitrary but fixed. Then*

$$\int_{-\infty}^{\infty} f(z; m, \lambda) dz = \int_{C_{\text{sd}}(m, \lambda)} f(z; m, \lambda) dz + 2\pi i \sum_{\substack{l \in \mathbb{Z} \\ \text{sign}(A)(l - z_{\text{st}}(m, \lambda)) > \eta}} \text{Res}_{z=z_l(\eta)} \{f(z; m, \lambda)\} \quad (95)$$

for all  $\eta \in ]0, \eta_0]$  and all  $\delta \in ]0, \frac{r|A|}{\pi}]$ . The sum of residues is zero if  $k \leq 0$ .

The proof is given in Appendix B. By (76) and Lemma 5.4 we conclude that

$$\begin{aligned} Z(r) &= r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} \sum_{m \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) \\ &\quad \times \left( \int_{C_{\text{sd}}(m, \lambda)} f(z; m, \lambda) dz + 2\pi i \sum_{\substack{l \in \mathbb{Z} \\ \text{sign}(A)(l - z_{\text{st}}(m, \lambda)) > \eta}} \text{Res}_{z=z_l(\eta)} \{f(z; m, \lambda)\} \right). \end{aligned} \quad (96)$$

Next we are going to separate the contributions coming from respectively poles and the integrals along the steepest descent contours.

**Lemma 5.5.** *Let  $\lambda \in \Lambda$  be arbitrary but fixed. Then the infinite series*

$$\Sigma_1(\lambda) = \sum_{m \in \mathbb{Z}} \int_{C_{\text{sd}}(m, \lambda)} f(z; m, \lambda) dz$$

is absolutely convergent for all  $\eta \in ]0, \eta_0]$  and all  $\delta \in ]0, 2r|A|/\pi[$ , and the infinite series

$$\Sigma_2(\lambda) = \sum_{m \in \mathbb{Z}} \sum_{\substack{l \in \mathbb{Z} \\ \text{sign}(A)(l - z_{\text{st}}(m, \lambda)) > \eta}} \text{Res}_{z=z_l(\eta)} \{f(z; m, \lambda)\}$$

is absolutely convergent for all  $\eta > 0$  and all  $\delta > 0$ . For all  $\eta, \delta \in ]0, \infty[$  we have

$$\Sigma_2(\lambda) = \sum_{l \in \mathbb{Z}} \beta(l, \lambda) \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi} l > \frac{|A|}{\pi} \eta}} \text{Res}_{z=i\eta} \{\phi(z; l, m, \lambda)\},$$

where

$$\begin{aligned} \beta(l, \lambda) &= (-1)^{kl} e^{ir(A l^2 + B l)}, \\ \phi(z; l, m, \lambda) &= e^{-\pi \delta (z+l)^2} q(z+l; \lambda) e^{ir A z^2} \frac{e^{2\pi i r (m + \frac{B}{2\pi} + \frac{A}{\pi} l) z}}{\sin^k(\pi(z - i\eta))}. \end{aligned} \quad (97)$$

Note that  $\Sigma_2(\lambda) = 0$  for  $k \leq 0$ .

For the proof, see Appendix B. Note that the sum over  $m$  in the expression (76) is absolutely convergent by Lemma 5.5. By (78)

$$\sum_{\lambda \in \Lambda} g(\lambda) \beta(l, \lambda) q(z + l) e^{ir(Az^2 + Bz + 2Alz)} = (-1)^{kl} \sum_{\lambda \in \Lambda} g(\lambda) q(z + l) e^{rQ(z+l)} = 0 \quad (98)$$

for all  $z, l \in \mathbb{Z}$ . By (96) and Lemma 5.5 we have

$$Z(r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} (Z_{\text{int},1}(\delta, \eta) + Z_{\text{int,polar}}(\delta, \eta) + Z_{\text{polar},1}(\delta, \eta)), \quad (99)$$

where

$$\begin{aligned} Z_{\text{int},1}(\delta, \eta) &= \sum_{(m,\lambda) \in W} g(\lambda) \int_{C_{\text{sd}}(m,\lambda)} f(z; m, \lambda) dz, \\ Z_{\text{int,polar}}(\delta, \eta) &= \sum_{\lambda \in \Lambda} g(\lambda) \sum_{\substack{m \in \mathbb{Z} \\ z_{\text{st}}(m,\lambda) \in \mathbb{Z}}} \int_{C_{\text{sd}}(m,\lambda)} f(z; m, \lambda) dz, \\ Z_{\text{polar},1}(\delta, \eta) &= 2\pi i \sum_{\lambda \in \Lambda} g(\lambda) \sum_{l \in \mathbb{Z}} \beta(l, \lambda) \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi} l > \frac{|A|}{\pi} \eta}} \text{Res}_{z=i\eta} \{ \phi(z; l, m, \lambda) \}. \end{aligned} \quad (100)$$

Here we suppress the dependency on  $r$  and also to some extent the dependency on  $\lambda$  in our notation for not making it too clumsy. We have split the sum  $\Sigma_1$  in Lemma 5.5 into two parts, since this will be convenient for the following calculations. The reason for the names  $Z_{\text{int,polar}}$  and  $Z_{\text{polar},1}$  instead of  $Z_{\text{int},2}$  and  $Z_{\text{polar}}$  will become clear below.

Let us take a closer look at  $Z_{\text{int,polar}}(\delta, \eta)$ . Assume that  $z_{\text{st}}(m, \lambda) = l \in \mathbb{Z}$ . By changing variable to  $w = z - l$ , as in the proof of Lemma 5.5, the contour  $C_{\text{sd}}(m, \lambda)$  changes to  $C(\lambda, 0)$ , where, for  $\rho \in \mathbb{R}$ , the contour  $C(\lambda, \rho)$  is given by

$$\text{Im}(w) = \text{sign}(A) \text{Re}(w) - \rho$$

oriented so that we start at  $-\infty e^{\frac{i\pi}{4} \text{sign}(A)} - i\rho$ . It follows that

$$Z_{\text{int,polar}}(\delta, \eta) = \sum_{\lambda \in \Lambda} g(\lambda) \sum_{\substack{l \in \mathbb{Z} \\ \frac{B}{2\pi} + \frac{A}{\pi} l \in \mathbb{Z}}} \beta(l, \lambda) I(l; \lambda, \delta, \eta),$$

where  $I(l; \lambda, \delta, \eta) = \int_{C(\lambda, 0)} \phi(w; \lambda, m, l) dw$ , where

$$\phi(w; \lambda, m, l) = e^{-\pi\delta(w+l)^2} q(w+l) \frac{e^{irAw^2}}{\sin^k(\pi(w-i\eta))}$$

by (97) since  $m + \frac{B}{2\pi} + \frac{A}{\pi} l = 0$ . By changing variable once more to  $y = w - i\eta$  we get  $I(l; \lambda, \delta, \eta) = \int_{C(\lambda, \eta)} F(y) dy$  with

$$F(y) = F(y; l, \lambda, \delta, \eta) = e^{-\pi\delta(y+l+i\eta)^2} q(y+l+i\eta) \frac{e^{irA(y+i\eta)^2}}{\sin^k(\pi y)}. \quad (101)$$

We split the function  $F$  into its even and odd parts, i.e.  $F(y) = F^+(y) + F^-(y)$ , where

$$F^\pm(y) = \frac{1}{2}(F(y) \pm F(-y)).$$

**Lemma 5.6.** *Let  $\lambda \in \Lambda$  and  $l \in \mathbb{Z}$  be arbitrary but fixed. Then*

$$\int_{C(\lambda, \eta)} F^-(y; l, \lambda, \delta, \eta) dy = \pi i \operatorname{Res}_{y=0} F(y; \lambda, l, \delta, \eta)$$

for any  $\eta \in ]0, 1[$  and any  $\delta \in ]0, 2r|A|/\pi[$ . Moreover, the series

$$Z_{\text{polar}, 2}(\delta, \eta) := \sum_{\lambda \in \Lambda} g(\lambda) \sum_{\substack{l \in \mathbb{Z} \\ \frac{B}{2\pi} + \frac{A}{\pi} l \in \mathbb{Z}}} \beta(l, \lambda) \pi i \operatorname{Res}_{y=0} F(y; l, \lambda, \delta, \eta)$$

is absolutely convergent for all  $\delta, \eta \in ]0, \infty[$ . For  $k \leq 0$  this sum is zero.

The proof is given in Appendix B. By Lemma 5.5 the sum  $Z_{\text{int}, \text{polar}}(\delta, \eta)$  is absolutely convergent for  $\eta \in ]0, \eta_0[$  and  $\delta \in ]0, 2rA_0/\pi[$ , and by Lemma 5.6 we then get that

$$Z_{\text{int}, 2}(\delta, \eta) := \sum_{\lambda \in \Lambda} g(\lambda) \sum_{\substack{l \in \mathbb{Z} \\ \frac{B}{2\pi} + \frac{A}{\pi} l \in \mathbb{Z}}} \beta(l, \lambda) \int_{C(\lambda, \eta)} F^+(y; l, \lambda, \delta, \eta) dy \quad (102)$$

is absolutely convergent and that

$$Z_{\text{int}, \text{polar}}(\delta, \eta) = Z_{\text{polar}, 2}(\delta, \eta) + Z_{\text{int}, 2}(\delta, \eta)$$

for these  $(\delta, \eta)$ . We put

$$Z_{\text{polar}}(\delta, \eta) = Z_{\text{polar}, 1}(\delta, \eta) + Z_{\text{polar}, 2}(\delta, \eta).$$

To conclude we have so far proven that

$$Z(r) = r \lim_{\delta \rightarrow 0+} \sqrt{\delta} \lim_{\eta \rightarrow 0+} (Z_{\text{int}, 1}(\delta, \eta) + Z_{\text{int}, 2}(\delta, \eta) + Z_{\text{polar}}(\delta, \eta)), \quad (103)$$

where  $Z_{\text{int}, 1}(\delta, \eta)$  and  $Z_{\text{int}, 2}(\delta, \eta)$  are given by respectively (100) and (102), and

$$Z_{\text{polar}}(\delta, \eta) = \sum_{l \in \mathbb{Z}} Z^l(\delta, \eta), \quad (104)$$

where  $Z^l(\delta, \eta) = Z_1^l(\delta, \eta) + Z_2^l(\delta, \eta)$  with

$$\begin{aligned} Z_1^l(\delta, \eta) &= 2\pi i \sum_{\lambda \in \Lambda} g(\lambda) \beta(l, \lambda) \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi} l > \frac{|A|}{\pi} \eta}} \operatorname{Res}_{z=i\eta} \{\phi(z; l, m, \lambda)\}, \\ Z_2^l(\delta, \eta) &= \pi i \sum_{\substack{\lambda \in \Lambda \\ \frac{B}{2\pi} + \frac{A}{\pi} l \in \mathbb{Z}}} g(\lambda) \beta(l, \lambda) \operatorname{Res}_{z=i\eta} \left\{ \phi \left( z; \lambda, - \left( \frac{B}{2\pi} + \frac{A}{\pi} l \right), l \right) \right\} \end{aligned}$$

(where, as usual, an empty sum is put equal to zero). By (82) we have that  $m + \frac{B}{2\pi} + \frac{A}{\pi} l > \frac{|A|}{\pi} \eta$  is equivalent to  $\operatorname{sign}(A)(l - z_{\text{st}}(m, \lambda)) > \eta$ . By (88) this is also equivalent to  $\operatorname{sign}(A)(l - z_{\text{st}}(m, \lambda)) > 0$  for  $\eta \in ]0, \eta_0[$ . For  $\eta \in ]0, \eta_0[$  we therefore have

$$Z^l(\delta, \eta) = 2\pi i \sum_{\lambda \in \Lambda} g(\lambda) \beta(l, \lambda) \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi} l \geq 0}} \operatorname{Res}_{z=i\eta} \left\{ \frac{\phi(z; l, m, \lambda)}{\operatorname{Sym}_{\pm} \left( m + \frac{B}{2\pi} + \frac{A}{\pi} l \right)} \right\}, \quad (105)$$

where  $\operatorname{Sym}_{\pm}$  is given by (44).

Let us make some comments on the strategy we will follow from here. Idealistically we can calculate the large  $r$  asymptotics of  $Z(r)$  in (103) by calculating the large  $r$  asymptotics of the limit  $\lim_{\delta \rightarrow 0+} \sqrt{\delta} \lim_{\eta \rightarrow 0+} f(\delta, \eta)$  for  $f$  equal to each of the functions  $Z_{\text{int}, 1}$ ,  $Z_{\text{int}, 2}$  and  $Z_{\text{polar}}$

(assuming that these limits exist). This is actually what we will attempt to do. In fact we start by showing that

$$Z_{\text{polar}}(r) := r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}(\delta, \eta) \quad (106)$$

exists, and at the same time we calculate this limit. Thus we not only give an asymptotic description of this limit but we actually find an exact expression for it. To do this we need to assume that certain “symmetry” conditions are satisfied.

To handle the two other limits  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{int}, \nu}(\delta, \eta)$ ,  $\nu = 1, 2$ , more care need be taken. Since the limits on the right-hand sides of (103) and (106) exist we have that

$$Z_{\text{int}}(r) := r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{int}}(\delta, \eta) \quad (107)$$

exists, where  $Z_{\text{int}}(\delta, \eta) = Z_{\text{int},1}(\delta, \eta) + Z_{\text{int},2}(\delta, \eta)$ . We will, however, not calculate this limit explicitly, but only give it an asymptotic description. This is of course sufficient for the proof of Theorem 4.1. Thus we will show that we for each  $N \in \mathbb{Z}_{\geq 0}$  have a decomposition

$$Z_{\text{int}, \nu}(\delta, \eta) = Z_{\text{int}, \nu}(N; \delta, \eta) + R_{\nu}(N; \delta, \eta)$$

such that

$$Z_{\text{int}, \nu}(r; N) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{int}, \nu}(N; \delta, \eta)$$

exists,  $\nu = 1, 2$ . This implies then that

$$R(r; N) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} (R_1(N; \delta, \eta) + R_2(N; \delta, \eta))$$

exists. We will not show that  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} R_{\nu}(N; \delta, \eta)$  exists for each  $\nu = 1, 2$  separately. Instead we show that  $R_{\nu}(N; \delta, \eta)$  is bounded from above by a certain function  $A_{\nu}(N; \delta, \eta)$  for which

$$A_{\nu}(r; N) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} A_{\nu}(N; \delta, \eta)$$

can easily be calculated and shown to be small compared to  $Z_{\text{int}, \nu}(r; N)$  in the limit of large  $r$ ,  $\nu = 1, 2$ . This will prove that  $Z_{\text{int},1}(r; N) + Z_{\text{int},2}(r; N)$  is the large  $r$  asymptotics of  $Z_{\text{int}}(r)$  to an order depending on  $N$ . To carry out this program we have to assume that certain functions admit certain periodicities.

One final small remark is appropriate. When dealing with  $Z_{\text{int},1}(\delta, \eta)$  we will start by getting rid of the parameter  $\eta$  by showing that

$$Z_{\text{int},1}(\delta) := \lim_{\eta \rightarrow 0_+} Z_{\text{int},1}(\delta, \eta)$$

exists and then make a decomposition

$$Z_{\text{int},1}(\delta) = Z_{\text{int},1}(N; \delta) + R_1(N; \delta)$$

for each  $N \in \mathbb{Z}_{\geq 0}$ . This makes things a little easier. When dealing with  $Z_{\text{int},2}(\delta, \eta)$  this is not possible. We will see that the positive parameter  $\eta$  plays a crucial role in the calculations and can only be removed in the final step.

**5.1.2. A calculation of the polar contribution  $Z_{\text{polar}}(r)$ .** In this section we calculate the limit  $Z_{\text{polar}}(r)$  in (106). If  $k \leq 0$  this is zero so here we assume that  $k > 0$ . By (104)  $Z_{\text{polar}}(\delta, \eta)$  is given by two infinite sums, the  $l$ -sum and the  $m$ -sum. We will change the  $l$ -sum by using (70) ‘backwards’. The  $m$ -sum will be calculated explicitly. Before we can do this we need to write  $Z^l(\delta, \eta)$  in (105) in a different way. A main problem is that the lower bound in the  $m$ -sum depends on  $l$  in the present expression for  $Z^l(\delta, \eta)$ .

We begin by a list of assumptions. Recall that the index set  $\Lambda$  is a subset of  $\mathbb{R}^e$ . We assume that  $A, B, g$ , and  $q(z)$  (for a fixed but arbitrary  $z \in \mathbb{C}$ ) as functions of  $\lambda$  are defined on all of  $\mathbb{R}^e$ , and that  $A$  and  $B$  are real on all of  $\mathbb{R}^e$ .

Let  $V_l : \mathbb{R}^e \rightarrow \mathbb{R}$  and  $U_l : \mathbb{C} \times \mathbb{R}^e \rightarrow \mathbb{C}$  be given by

$$\begin{aligned} V_l(x) &= -\frac{B(x)}{2\pi} - \frac{A(x)}{\pi}l, \\ U_l(z; x) &= g(x)\beta(l, x)q(z+l; x)e^{irA(x)z^2}. \end{aligned}$$

Assume that there for each  $l \in \mathbb{Z}$  exists a bijection  $T_l : \mathbb{R}^e \rightarrow \mathbb{R}^e$  such that

$$V(x) := V_l(T_l^{-1}(x))$$

only depends on  $x$  (and not on  $l$ ). We will assume that  $T_0 = \text{id}_{\mathbb{R}^e}$ . By (105) and (97) we then have

$$Z^l(\delta, \eta) = 2\pi i \sum_{x \in \Lambda(l)} \sum_{\substack{m \in \mathbb{Z} \\ m - V(x) \geq 0}} \frac{\text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{K_l(z; x)}{\sin^k(\pi(z-i\eta))} e^{2\pi ir(m-V(x))z} \right\}}{\text{Sym}_{\pm}(m - V(x))}, \quad (108)$$

where  $\Lambda(l) = T_l(\Lambda)$  and  $K_l(z; x) = U_l(z; T_l^{-1}(x))$ . In the above expression the lower bound in the  $m$ -sum still depends on  $l$  since  $\Lambda(l)$  depends on  $l$ . Moreover, in order to be able to use (70) on the  $l$ -sum, the expression needs to be ‘sufficiently’ periodic in  $l$ . We will see below what is meant by that. Let us just remark here that a main part consists of changing the finite index set  $\Lambda(l)$  to a new index set  $\Gamma(l)$  which is periodic in  $l$ . To be able to do this we need to assume that there are some additional symmetries present. Firstly, we will assume that there exists a positive integer  $L$  and a subset  $\mathcal{R}$  of  $\mathbb{R}^e$  containing  $\cup_{l \in \mathbb{Z}} \Lambda(l)$  such that  $K_{l+L}(z; x) = K_l(z; x)$  for each  $l \in \mathbb{Z}$ ,  $z \in \mathbb{C}$  and  $x \in \mathcal{R}$ . Secondly, we assume that there exists a family of bijections  $\{S_j : \mathcal{R} \rightarrow \mathcal{R}\}_{j \in I}$  all independent of  $l$  such that the maps  $x \mapsto K_l(z; x)$  and  $x \mapsto V(x) \pmod{\mathbb{Z}}$  restricted to  $\mathcal{R}$  are invariant under these transformations, i.e.  $V(S_j(x)) \equiv V(x) \pmod{\mathbb{Z}}$  for all  $x \in \mathcal{R}$  and  $K_l(z; S_j(x)) = K_l(z; x)$  for all  $x \in \mathcal{R}$  and all  $z \in \mathbb{C}$  and  $l \in \mathbb{Z}$ .

For a tuple of integers  $\mathcal{C} = (j_1, \dots, j_m) \in I^m$  and a sequence of signs  $\underline{\epsilon} \in \{\pm 1\}^m$  we let

$$S_{\mathcal{C}}^{\underline{\epsilon}} = S_{j_m}^{\epsilon_m} \circ \dots \circ S_{j_1}^{\epsilon_1},$$

where  $S_j^{-1}$  is the inverse of  $S_j$ . We then finally make the following ‘symmetry’ assumption: There exists a family of subsets  $\Gamma(l) \subseteq \mathcal{R}$ ,  $l \in \mathbb{Z}$ , periodic in  $l$  such that there for any  $l \in \mathbb{Z}$  and any  $x \in \Lambda(l)$  exists a tuple of integers  $\mathcal{C}_x^l$  and a sequence of signs  $\underline{\epsilon}_x^l$  satisfying  $S_{\mathcal{C}_x^l}^{\underline{\epsilon}_x^l}(x) \in \Gamma(l)$ . For simplicity we will assume that the map  $\Lambda(l) \rightarrow \Gamma(l)$ ,  $x \mapsto S_{\mathcal{C}_x^l}^{\underline{\epsilon}_x^l}(x)$ , is a bijection, but the following analysis can actually be carried out with only minor changes in case this map is only surjective.

Since the product of the periods of two periodic functions is a period for both of the functions we can assume that the family of sets  $\Gamma(l)$  is periodic in  $l$  with a period of  $L$ , i.e.  $\Gamma(l+L) = \Gamma(l)$  for all  $l \in \mathbb{Z}$ . Since the expression (108) only depends on  $m$  and  $V(x)$  through their difference  $m - V(x)$  we can always modify the different values of  $V(x)$  by appropriate integers. In particular we have

$$Z^l(\delta, \eta) = 2\pi i \sum_{\nu \in \Gamma(l)} \sum_{\substack{m \in \mathbb{Z} \\ m - V(\nu) \geq 0}} \frac{\text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{K_l(z; \nu)}{\sin^k(\pi(z-i\eta))} e^{2\pi ir(m-V(\nu))z} \right\}}{\text{Sym}_{\pm}(m - V(\nu))}. \quad (109)$$



Note that

$$\sum_{\nu \in \Gamma(l)} K_l(z; \nu) e^{-2\pi i r V(\nu)z} \quad (110)$$

is periodic in  $l$  with a period of  $L$  for any fixed  $z \in \mathbb{C}$ . By (98) we have

$$\sum_{\nu \in \Gamma(l)} K_l(z; \nu) e^{-2\pi i r V(\nu)z} = (-1)^{kl} \sum_{\lambda \in \Lambda} g(\lambda) q(z+l) e^{rQ(z+l)} = 0 \quad (111)$$

for all  $z \in \mathbb{Z}$ .

Let us consider the case  $Z(r) = Z(X; r)$ . Here we have  $\Lambda \subseteq \mathbb{R}^{2n}$  by (79), so  $e = 2n$  here. Let  $T_l(x, y) = (x, y + l\beta/2)$  for  $x, y \in \mathbb{R}^n$ . Then  $\Lambda(l) = T_l(\Lambda) \subseteq \{\pm 1\}^n \times (\frac{1}{2}\mathbb{Z})^n$  for all  $l \in \mathbb{Z}$ . In particular, we can let  $\mathcal{R} = \{\pm 1\}^n \times (\frac{1}{2}\mathbb{Z})^n$ . We have

$$V(x, y) = V_l(T_l^{-1}(x, y)) = \sum_{j=1}^n \frac{y_j - l\beta_j/2}{\alpha_j} + \frac{l}{2} \sum_{j=1}^n \frac{\beta_j}{\alpha_j} = \sum_{j=1}^n \frac{y_j}{\alpha_j}$$

which is independent of  $l$  (and  $x$ ). For  $j = 1, 2, \dots, n$  we let  $S_j : \mathcal{R} \rightarrow \mathcal{R}$  be the transformation  $S_j(\underline{\mu}, y) = (\underline{\mu}, y + \alpha_j e_j)$ , where  $e_j$  is the standard  $j$ th unit vector of  $\mathbb{R}^n$ . Then  $V(S_j(\underline{\mu}, y)) = V(\underline{\mu}, y) + 1$  for all  $j$  and all  $(\underline{\mu}, y) \in \mathcal{R}$ . By (79) and (97) we have

$$\begin{aligned} U_l(z; (x, y)) &= g(x, y) \beta(l, y) q(z+l; y) e^{irAz^2} \\ &= (-1)^{nl} \left(\frac{i}{2}\right)^n \left(\prod_{j=1}^n x_j\right) \exp\left(2\pi i \sum_{j=1}^n \frac{\rho_j}{\alpha_j} [ry_j^2 + x_j y_j]\right) \exp\left(\frac{\pi i r}{2} E z^2\right) \\ &\quad \times \exp\left(\frac{\pi i r}{2} E l^2\right) \exp\left(-2\pi i r \left(\sum_{j=1}^n \frac{y_j}{\alpha_j}\right) l\right) \exp\left(-\pi i \left(\sum_{j=1}^n \frac{x_j}{\alpha_j}\right) (z+l)\right) \end{aligned}$$

for  $z \in \mathbb{C}$  and  $x, y \in \mathbb{R}^n$ . But then

$$\begin{aligned} K_l(z; (x, y')) &= (-1)^{nl} \left(\frac{i}{2}\right)^n \left(\prod_{j=1}^n x_j\right) \exp\left(2\pi i \sum_{j=1}^n x_j \left(\frac{\rho_j}{\alpha_j} y'_j - \frac{1}{2} \sigma_j l\right)\right) \\ &\quad \times \exp\left(2\pi i r \left(-\frac{1}{4} \beta_0 l^2 + \sum_{j=1}^n \left[\frac{\rho_j}{\alpha_j} y_j'^2 - \frac{1}{4} \sigma_j \beta_j l^2\right]\right)\right) \\ &\quad \times \exp\left(\frac{\pi i r}{2} E z^2\right) \exp\left(-\pi i \left(\sum_{j=1}^n \frac{x_j}{\alpha_j}\right) z\right) \\ &\quad \times \exp\left(-2\pi i r \sum_{j=1}^n \left(\sigma_j \left(y'_j - \frac{1}{2} \beta_j l\right) l\right)\right) \end{aligned}$$

for  $z \in \mathbb{C}$  and  $x, y' \in \mathbb{R}^n$ . Thus, for fixed  $(x, y') \in \mathcal{R}$  and  $z \in \mathbb{C}$ ,  $K_l(z; (x, y'))$  is periodic in  $l$  with a period of 2. Moreover,  $(x, y') \mapsto K_l(z, (x, y'))$ ,  $\mathcal{R} \rightarrow \mathbb{C}$ , is invariant under the transformations  $S_j$  for all  $l \in \mathbb{Z}$  and all  $z \in \mathbb{C}$ . Finally, it is easy to see that we can put

$$\Gamma(l) = \{\pm 1\}^n \times \{\underline{n}' \in \mathbb{Z}^n + \frac{1}{2}\underline{\beta}l \mid \underline{0} \leq \underline{n}' < \underline{\alpha}\}.$$

In particular, the period  $L = 2$  in this case. Note that the last factor in the above expression for  $K_l(z; (x, y'))$ , namely  $\exp\left(-2\pi i r \sum_{j=1}^n \left(\sigma_j \left(y'_j - \frac{1}{2} \beta_j l\right) l\right)\right)$ , is 1 for  $(x, y') \in \Gamma(l)$ .

Let us return to the general case. The next step is to get rid of the infinite sum over  $m$  in (109). A problem here is that the lower bound in this sum is  $V(\nu)$ . To overcome this difficulty we use the following small lemma which proof is left to the reader. The function  $\text{Sym}_\pm$  is given by (44).

**Lemma 5.7.** *If  $a \in \mathbb{R}$  and  $f : \mathbb{Z} \rightarrow \mathbb{C}$  is a function such that*

$$\sum_{\substack{m \in \mathbb{Z} \\ m \geq a}} \frac{f(m)}{\text{Sym}_\pm(m-a)}$$

*is convergent, then*

$$\sum_{\substack{m \in \mathbb{Z} \\ m \geq a}} \frac{f(m)}{\text{Sym}_\pm(m-a)} = \sum_{m=0}^{\infty} \frac{f(m)}{\text{Sym}_\pm(m)} - \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |a|}} \frac{\text{sign}(a)f(\text{sign}(a)m)}{\text{Sym}_\pm(m)\text{Sym}_\pm(m-|a|)}.$$

□

By using this result on (109) we immediately get

$$\begin{aligned} Z^l(\delta, \eta) &= 2\pi i \sum_{\nu \in \Gamma(l)} \left\{ \sum_{m=0}^{\infty} \frac{\text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{K_l(z;\nu)}{\sin^k(\pi(z-i\eta))} e^{2\pi i r(m-V)z} \right\}}{\text{Sym}_\pm(m)} \right. \\ &\quad \left. - \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |V|}} \frac{\text{sign}(V) \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{K_l(z;\nu)}{\sin^k(\pi(z-i\eta))} e^{2\pi i r \text{sign}(V)(m-|V|)z} \right\}}{\text{Sym}_\pm(m)\text{Sym}_\pm(m-|V|)} \right\}, \end{aligned} \quad (112)$$

where we write  $V$  for  $V(\nu)$ . We could of course have used Lemma 5.7 directly on (105). A problem with this is however that the resulting expression contains a sum of the form  $\sum_{0 \leq m \leq |\frac{B}{2\pi} + \frac{A}{\pi}l|}$  depending on  $l$  in a bad (=nonperiodic) way. If we use Lemma 5.7 on (108) the resulting expression contains a sum of the form  $\sum_{0 \leq m \leq |V(x)|}$ . The problem here is that  $V(x)$  depends on  $x \in \Lambda(l)$  and thereby on  $l$  in a bad way.

Let  $\nu \in \Gamma(l)$  be fixed and let

$$\Psi(z; m) = \Psi(z; l, m, \nu, \delta) = e^{-\pi\delta(z+l)^2} K_l(z; \nu) e^{2\pi i r(m-V)z}.$$

By using the identity

$$\sum_{m=0}^{\infty} \frac{e^{2\pi i r m z}}{\text{Sym}_\pm(m)} = \frac{i}{2} \cot(\pi r z),$$

valid for all  $z \in D(i\eta, \eta)$ , we get

$$\sum_{m=0}^{\infty} \frac{\text{Res}_{z=i\eta} \left\{ \frac{\Psi(z; m)}{\sin^k(\pi(z-i\eta))} \right\}}{\text{Sym}_\pm(m)} = \frac{i}{2} \text{Res}_{z=i\eta} \left\{ \frac{\Psi(z; 0)}{\sin^k(\pi(z-i\eta))} \cot(\pi r z) \right\}. \quad (113)$$

To see this, assume that  $q(z+l)$  has a zero of order  $k_l \geq 0$  in  $z = i\eta$ . If  $k_l \geq k$  then both sides of (113) are zero. If  $k_l < k$  we let  $k' = k - k_l$  and put

$$u(z) = \Psi(z; 0) \frac{(z-i\eta)^{k'}}{\sin^k(\pi(z-i\eta))}.$$

Then

$$\sum_{m=0}^{\infty} \frac{\text{Res}_{z=i\eta} \left\{ \frac{\Psi(z;m)}{\sin^k(\pi(z-i\eta))} \right\}}{\text{Sym}_{\pm}(m)} = \sum_{m=0}^{\infty} \frac{1}{(k'-1)!} \left( \frac{d}{dz} \right)^{k'-1} u_m(z) \Big|_{z=i\eta},$$

where  $u_m(z) = u(z) \frac{e^{2\pi i r m z}}{\text{Sym}_{\pm}(m)}$ . Note that  $u_m$  is analytic on  $D(i\eta, \eta)$ , where we assume that  $\eta \leq 1$ . Now let  $\rho \in ]0, \eta[$ . For  $z \in D(i\eta, \rho)$  we have

$$\sum_{m=0}^{\infty} |u_m(z)| \leq |u(z)| \sum_{m=0}^{\infty} \frac{e^{-2\pi m r (\eta - \rho)}}{\text{Sym}_{\pm}(m)}$$

where the infinite sum on the right-hand side is convergent, so by Weierstrass' M-test  $U(z) = \sum_{m=0}^{\infty} u_m(z)$  is uniformly convergent on  $D(i\eta, \rho)$ . It follows that  $U$  is analytic on  $D(i\eta, \eta)$  and that all its derivatives may be calculated by term-by-term differentiation, see e.g. [T, p. 95]. Therefore

$$\begin{aligned} \text{Res}_{z=i\eta} \frac{\Psi(z;0)}{\sin^k(\pi(z-i\eta))} \frac{i}{2} \cot(\pi r z) &= \frac{1}{(k'-1)!} \left( \frac{d}{dz} \right)^{k'-1} u(z) \frac{i}{2} \cot(\pi r z) \Big|_{z=i\eta} \\ &= \frac{1}{(k'-1)!} \left( \frac{d}{dz} \right)^{k'-1} U(z) \Big|_{z=i\eta} = \frac{1}{(k'-1)!} \sum_{m=0}^{\infty} \left( \frac{d}{dz} \right)^{k'-1} u_m(z) \Big|_{z=i\eta}. \end{aligned}$$

By (112) and (113) we have

$$\begin{aligned} Z^l(\delta, \eta) &= 2\pi i \sum_{\nu \in \Gamma(l)} \left\{ \frac{i}{2} \text{Res}_{z=i\eta} \left\{ \frac{\Psi(z;l,0,\nu,\delta)}{\sin^k(\pi(z-i\eta))} \cot(\pi r z) \right\} \right. \\ &\quad \left. - \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |V|}} \frac{\text{sign}(V) \text{Res}_{z=i\eta} \left\{ \frac{\Psi(z;l,\text{sign}(V)m,\nu,\delta)}{\sin^k(\pi(z-i\eta))} \right\}}{\text{Sym}_{\pm}(m) \text{Sym}_{\pm}(m-|V|)} \right\}. \end{aligned} \quad (114)$$

**Lemma 5.8.** *Let  $\eta_2 \in ]0, 1/r[$ . Then the infinite series  $\sum_{l \in \mathbb{Z}} Z^l(\delta, \eta)$  is uniformly convergent with respect to  $\eta$  on  $]0, \eta_2]$ . We therefore have*

$$Z_{\text{polar}}(\delta) := \lim_{\eta \rightarrow 0+} \sum_{l \in \mathbb{Z}} Z^l(\delta, \eta) = \sum_{l \in \mathbb{Z}} Z^l(\delta),$$

where

$$\begin{aligned} Z^l(\delta) &= \lim_{\eta \rightarrow 0+} Z^l(\delta, \eta) \\ &= 2\pi i \sum_{\nu \in \Gamma(l)} \left\{ \frac{i}{2} \text{Res}_{z=0} \left\{ \frac{\Psi(z;l,0,\nu,\delta)}{\sin^k(\pi z)} \cot(\pi r z) \right\} \right. \\ &\quad \left. - \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |V|}} \frac{\text{sign}(V) \text{Res}_{z=0} \left\{ \frac{\Psi(z;l,\text{sign}(V)m,\nu,\delta)}{\sin^k(\pi z)} \right\}}{\text{Sym}_{\pm}(m) \text{Sym}_{\pm}(m-|V|)} \right\}. \end{aligned}$$

The proofs of this lemma and the next proposition are given in Appendix C. By the next proposition we end the calculation of the contribution to  $Z(r)$  coming from the polar term  $Z_{\text{polar}}(\delta, \eta)$ . Note that this contribution is given an exact (non-asymptotic) description.

**Proposition 5.9.** *We have*

$$Z_{\text{polar}}(r) := r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} Z_{\text{polar}}(\delta) = \sum_{l=0}^{L-1} Z^l,$$

where

$$\begin{aligned} Z^l &= \frac{2\pi r}{L} \sum_{\nu \in \Gamma(l)} \left[ \frac{i}{2} \text{Res}_{z=0} \left\{ \frac{K_l(z; \nu) e^{-2\pi i r V z}}{\sin^k(\pi z)} \cot(\pi r z) \right\} \right. \\ &\quad \left. - \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |V|}} \frac{\text{sign}(V)}{\text{Sym}_{\pm}(m) \text{Sym}_{\pm}(m - |V|)} \text{Res}_{z=0} \left\{ \frac{K_l(z; \nu) e^{2\pi i r \text{sign}(V)(m - |V|)z}}{\sin^k(\pi z)} \right\} \right], \end{aligned}$$

where  $V = V(\nu)$ .

Note that this expression does not in general give a function as in (30). The factors  $e^{2\pi i r q_j}$  can, however, be hidden in the expression. Let us see that this is indeed the case when  $Z(r) = Z(X; r)$ .

**Corollary 5.10.** *Let  $Z(r) = Z(X; r)$ . Then*

$$Z_{\text{polar}}(r) = \sum_{(l, \underline{n}') \in J} b_{(l, \underline{n}')} \exp(2\pi i r q_{(l, \underline{n}')} r) \left( Z_a^{(l, \underline{n}')} (r) + Z_2^{(l, \underline{n}')} (r) \right),$$

where  $q_{(l, \underline{n}')}$  is given by (40),

$$J = \left\{ (l, \underline{n}') \in \{0, 1\} \times \left( \frac{1}{2} \mathbb{Z} \right)^2 \left| \begin{array}{l} \underline{n}' \in \mathbb{Z}^n + \frac{1}{2} l \underline{\beta}, \\ \underline{0} \leq \underline{n}' < \underline{\alpha} \end{array} \right. \right\},$$

and  $b_{(l, \underline{n}')} = \pi i (-1)^{nl} (-1)^n$ , and where

$$\begin{aligned} Z_a^{(l, \underline{n}')} (r) &= \frac{i}{2} \text{Res}_{z=0} \left\{ \frac{\exp \left( \frac{\pi i r}{2} E z^2 - 2\pi i r \left( \sum_{j=1}^n \frac{n'_j}{\alpha_j} \right) z \right)}{\sin^{n+a_\epsilon g-2}(\pi z)} \cot(\pi r z) \right. \\ &\quad \left. \times \prod_{j=1}^n \sin \left( 2\pi \left[ \frac{\rho_j}{\alpha_j} n'_j - \frac{z}{2\alpha_j} - \frac{1}{2} \sigma_j l \right] \right) \right\}, \\ Z_b^{(l, \underline{n}')} (r) &= - \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq \left\lfloor \sum_{j=1}^n \frac{n'_j}{\alpha_j} \right\rfloor}} \frac{\text{sign} \left( \sum_{j=1}^n \frac{n'_j}{\alpha_j} \right)}{\text{Sym}_{\pm}(m) \text{Sym}_{\pm} \left( m - \left\lfloor \sum_{j=1}^n \frac{n'_j}{\alpha_j} \right\rfloor \right)} \\ &\quad \times \text{Res}_{z=0} \left\{ \frac{\exp \left( \frac{\pi i r}{2} E z^2 + 2\pi i r \text{sign} \left( \sum_{j=1}^n \frac{n'_j}{\alpha_j} \right) \left( m - \left\lfloor \sum_{j=1}^n \frac{n'_j}{\alpha_j} \right\rfloor \right) z \right)}{\sin^{n+a_\epsilon g-2}(\pi z)} \right. \\ &\quad \left. \times \prod_{j=1}^n \sin \left( 2\pi \left[ \frac{\rho_j}{\alpha_j} n'_j - \frac{z}{2\alpha_j} - \frac{1}{2} \sigma_j l \right] \right) \right\}. \end{aligned}$$

**Proof.** The corollary is an immediate consequence of Proposition 5.9 and the remarks above Lemma 5.7 about the Seifert case. We use that

$$\begin{aligned} & \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \exp \left( -\pi i \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} n'_j - \frac{1}{2} \sigma_j l \right) \right] \\ &= (2i)^n \prod_{j=1}^n \sin \left( 2\pi \left[ \frac{\rho_j}{\alpha_j} n'_j - \frac{z}{2\alpha_j} - \frac{1}{2} \sigma_j l \right] \right). \end{aligned}$$

□

We observe that this expression is not identical with the one stated in Theorem 4.1. It has the correct form, but the set  $J$  contains twice as many elements as the set  $\mathcal{I}_2$  in (43). By (40) it follows that if  $(l, \underline{n}') = (l, n'_1, \dots, n'_j) \in J$  then  $(l, \underline{n}' + (\alpha_j - 2n'_j)e_j) \in J$  and

$$q(l, \underline{n}' + (\alpha_j - 2n'_j)e_j) = q(l, \underline{n}').$$

What is even more important is that we expect the connected components of the moduli space of flat  $\text{SU}(2)$ -connections on  $X$  to be parametrized by  $\mathcal{I}_1 \cup \mathcal{I}_2$ . Therefore it is desirable to obtain an expression for  $Z_{\text{polar}}(r)$  similar to the one in the above corollary but with the sum over  $J$  replaced by a sum over  $\mathcal{I}_2$ . To obtain the result stated in Theorem 4.1 we have to use some extra symmetries before using Lemma 5.7. Let  $Z(r) = Z(X; r)$  and let

$$J_l = \left\{ \underline{n}' \in \mathbb{Z}^n + \frac{1}{2}\underline{\beta}l \mid \underline{0} \leq \underline{n}' < \underline{\alpha} \right\}. \quad (115)$$

From (109) and the remarks above Lemma 5.7 about the Seifert case we get

$$\begin{aligned} Z^l(\delta, \eta) &= 2\pi i (-1)^{nl} \left( \frac{i}{2} \right)^n \sum_{\underline{n}' \in J_l} \exp(2\pi i r q(l, \underline{n}')) \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \\ &\quad \times \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} n'_j - \frac{1}{2} \sigma_j l \right) \right] \sum_{\substack{m \in \mathbb{Z} \\ m \geq \sum_{j=1}^n \frac{n'_j}{\alpha_j}}} \frac{1}{\text{Sym}_{\pm} \left( m - \sum_{j=1}^n \frac{n'_j}{\alpha_j} \right)} \\ &\quad \times \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \frac{\exp \left( \frac{\pi i r}{2} E z^2 \right)}{\sin^{n+a_{\epsilon} g-2}(\pi(z-i\eta))} \right. \\ &\quad \times \exp \left( -\pi i \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right) \exp \left( 2\pi i r \left( m - \sum_{j=1}^n \frac{n'_j}{\alpha_j} \right) z \right) \left. \right\}. \end{aligned}$$

Let

$$J'_l = \left\{ \underline{n}' \in \mathbb{Z}^n + \frac{1}{2}\underline{\beta}l \mid \underline{0} \leq \underline{n}' \leq \frac{1}{2}\underline{\alpha} \right\}. \quad (116)$$

In the expression for  $Z^l(\delta, \eta)$  we can change the sum  $\sum_{\underline{n}' \in J_l}$  to

$$\sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\underline{n}' \in J'_l} \frac{1}{\prod_{j=1}^n \text{Sym}_{\mathbb{Z}_{\pm}} \left( \frac{n'_j}{\alpha_j} \right)}$$

if we at the same time substitute  $\mu'_j n'_j$  for  $n'_j$  everywhere. The function  $\text{Sym}_{\mathbb{Z}_{\pm}}$  is given by (45). This leads to the identity

$$\begin{aligned}
Z^l(\delta, \eta) &= 2\pi i (-1)^{nl} \left(\frac{i}{2}\right)^n \sum_{\underline{n}' \in J'_l} \frac{1}{\prod_{j=1}^n \text{Sym}_{\mathbb{Z}_{\pm}}\left(\frac{n'_j}{\alpha_j}\right)} \exp(2\pi i r q_{(l, \underline{n}')} ) \\
&\quad \times \sum_{\substack{\underline{\mu}' \in \{\pm 1\}^n \\ m \geq \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j}}} \sum_{\substack{m \in \mathbb{Z} \\ m \geq \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j}}} \frac{1}{\text{Sym}_{\pm}\left(m - \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j}\right)} \sum_{\underline{\mu} \in \{\pm 1\}^n} \left(\prod_{j=1}^n \mu_j\right) \\
&\quad \times \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \\
&\quad \times \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \frac{\exp\left(\frac{\pi i r}{2} E z^2\right)}{\sin^{n+a_{\epsilon} g-2}(\pi(z-i\eta))} \right. \\
&\quad \left. \times \exp \left( -\pi i \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right) \exp \left( 2\pi i r \left( m - \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \right) z \right) \right\}. \tag{117}
\end{aligned}$$

By Lemma 5.7 and the remarks following that lemma we get

$$Z^l(\delta, \eta) = 2\pi i (-1)^{nl} \left(\frac{i}{2}\right)^n \sum_{\underline{n}' \in J'_l} \frac{1}{\prod_{j=1}^n \text{Sym}_{\mathbb{Z}_{\pm}}\left(\frac{n'_j}{\alpha_j}\right)} \exp(2\pi i r q_{(l, \underline{n}')} ) \left( Z_0^{(l, \underline{n}')}(\delta, \eta) + Z_1^{(l, \underline{n}')}(\delta, \eta) \right),$$

where

$$\begin{aligned}
Z_0^{(l, \underline{n}')}(\delta, \eta) &= \sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\underline{\mu} \in \{\pm 1\}^n} \left(\prod_{j=1}^n \mu_j\right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \\
&\quad \times \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \frac{\exp\left(\frac{\pi i r}{2} E z^2 - 2\pi i r \left( \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \right) z\right)}{\sin^{n+a_{\epsilon} g-2}(\pi(z-i\eta))} \right. \\
&\quad \left. \times \frac{i}{2} \cot(\pi r z) \exp \left( -i\pi \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right) \right\} \tag{118}
\end{aligned}$$

and

$$\begin{aligned}
Z_1^{(l, \underline{n}')}(\delta, \eta) &= - \sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \\
&\times \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq \left\lfloor \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \right\rfloor}} \frac{\text{sign} \left( \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \right)}{\text{Sym}_{\pm}(m) \text{Sym}_{\pm} \left( m - \left\lfloor \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \right\rfloor \right)} \\
&\times \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \exp \left( -i\pi \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right) \right. \\
&\times \left. \frac{\exp \left( \frac{\pi i r}{2} E z^2 + 2\pi i r \text{sign} \left( \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \right) \left( m - \left\lfloor \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \right\rfloor \right) z \right)}{\sin^{n+a_{\epsilon} g-2}(\pi(z-i\eta))} \right\}. \quad (119)
\end{aligned}$$

Here

$$\begin{aligned}
&\sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \exp \left( -i\pi \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right) \\
&= (2i)^n \prod_{j=1}^n \mu'_j \sin \left( 2\pi \left[ \frac{\rho_j}{\alpha_j} n'_j - \frac{z}{2} \frac{\mu'_j}{\alpha_j} - \frac{1}{2} \sigma_j l \right] \right),
\end{aligned}$$

where we use that  $\sin(x - \pi \sigma_j l) = \sin(x + \pi \sigma_j l)$  for  $x \in \mathbb{R}$  since  $\sigma_j l \in \mathbb{Z}$ . Furthermore, we have

$$\begin{aligned}
&\sum_{\underline{\mu}' \in \{\pm 1\}^n} \left[ \prod_{j=1}^n \mu'_j \sin \left( 2\pi \left[ \frac{\rho_j}{\alpha_j} n'_j - \frac{z}{2} \frac{\mu'_j}{\alpha_j} - \frac{1}{2} \sigma_j l \right] \right) \right] \exp \left( -2\pi i r \left( \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \right) z \right) \\
&= (-2)^n \prod_{j=1}^n \left[ i \sin \left[ 2\pi \left( \frac{\rho_j}{\alpha_j} n'_j - \frac{1}{2} \sigma_j l \right) \right] \cos \left( \pi \frac{z}{\alpha_j} \right) \sin \left( 2\pi r z \frac{n'_j}{\alpha_j} \right) \right. \\
&\quad \left. + \cos \left[ 2\pi \left( \frac{\rho_j}{\alpha_j} n'_j - \frac{1}{2} \sigma_j l \right) \right] \sin \left( \pi \frac{z}{\alpha_j} \right) \cos \left( 2\pi r z \frac{n'_j}{\alpha_j} \right) \right].
\end{aligned}$$

By inserting these expressions in the expressions for  $Z_0^{(l, \underline{n}')}(\delta, \eta)$  and  $Z_1^{(l, \underline{n}')}(\delta, \eta)$  and by proceeding like in the proofs of Lemma 5.8 and Proposition 5.9 we finally obtain that  $Z_{\text{polar}}(r) = Z_{\text{polar}}(X; r)$ , where  $Z_{\text{polar}}(X; r)$  is given by (46).

**5.1.3. An asymptotic description of  $Z_{\text{int},1}(\delta, \eta)$ 's contribution to  $Z_{\text{int}}(r)$ .** We proceed by analyzing the contribution to  $Z_{\text{int}}(r)$  in (107) coming from

$$Z_{\text{int},1}(\delta, \eta) = \sum_{(m, \lambda) \in W} g(\lambda) \int_{C_{\text{sd}}(m, \lambda)} f(z; m, \lambda) dz.$$

We saw in Lemma 5.5 that this infinite series is absolutely convergent for  $\delta \in ]0, 2rA_0/\pi[$  and  $\eta \in ]0, \eta_0]$ , where  $\eta_0$  is the positive constant from (88) and  $A_0$  is given by (94).

**Lemma 5.11.** *For every fixed  $\delta \in ]0, 2rA_0/\pi[$  the infinite series  $Z_{\text{int},1}(\delta, \eta)$  is uniformly convergent w.r.t.  $\eta$  on an interval of the form  $[0, \eta_1]$ , where  $\eta_1 > 0$ .*

The proof is given in Appendix D. By this lemma we conclude that

$$Z_{\text{int},1}(\delta) := \lim_{\eta \rightarrow 0_+} Z_{\text{int},1}(\delta, \eta) = \sum_{(m,\lambda) \in W} g(\lambda) \lim_{\eta \rightarrow 0_+} \int_{C_{\text{sd}}(m,\lambda)} f(z; m, \lambda) dz.$$

Recall here that  $f(z; m, \lambda)$  but not  $g(\lambda)$  and  $C_{\text{sd}}(m, \lambda)$  depend on  $\eta$ . Let in the following  $(m, \lambda) \in W$  be arbitrary but fixed and let  $z_{\text{st}} = z_{\text{st}}(m, \lambda)$ . By (90)–(92) we have

$$\int_{C_{\text{sd}}(m,\lambda)} f(z; m, \lambda) dz = e^{\frac{i\pi}{4}\text{sign}(A)} e^{-irAz_{\text{st}}^2} \int_{-\infty}^{\infty} \psi_{\eta}(t; \delta) \exp(-r|A|t^2) dt,$$

where  $\psi_{\eta}(t; \delta) = \kappa_{\eta}(\gamma_{(m,\lambda)}(t); \lambda, \delta)$ . By (80) and (135) we have

$$|\psi_{\eta}(t; \delta)| \leq K e^{Ct+a_0|t|}$$

for all  $t \in \mathbb{R}$  and all  $\eta \in [0, \eta_1]$ , where  $C = -\pi\sqrt{2}z_{\text{st}}\delta$  and  $K = \frac{b_0}{M^k} e^{-\pi\delta z_{\text{st}}^2} e^{a_0|z_{\text{st}}|}$ . Since  $t \mapsto K e^{Ct+a_0|t|} e^{-r|A|t^2} \in L^1(\mathbb{R})$  and  $\lim_{\eta \rightarrow 0_+} \psi_{\eta}(t; \delta) = \psi_0(t; \delta)$  for all  $t$  we conclude by Lebesgue's dominated convergence theorem that

$$\lim_{\eta \rightarrow 0_+} \int_{C_{\text{sd}}(m,\lambda)} f(z; m, \lambda) dz = \int_{C_{\text{sd}}(m,\lambda)} f_0(z; m, \lambda) dz,$$

where  $f_0(z; m, \lambda)$  is equal to  $f(z; m, \lambda)$  in (77) with  $\eta = 0$ , i.e.

$$f_0(z; m, \lambda) = e^{-\pi\delta z^2} q(z; \lambda) \frac{e^{rQ(z)} e^{2\pi i r m z}}{\sin^k(\pi z)}. \quad (120)$$

We have shown

**Lemma 5.12.** *For every fixed  $\delta \in ]0, 2rA_0/\pi[$  we have*

$$Z_{\text{int},1}(\delta) = \lim_{\eta \rightarrow 0_+} Z_{\text{int},1}(\delta, \eta) = \sum_{(m,\lambda) \in W} g(\lambda) \int_{C_{\text{sd}}(m,\lambda)} f_0(z; m, \lambda) dz,$$

where the functions  $f_0(z; m, \lambda)$  are given by (120).  $\square$

Next we give an asymptotic description of the integrals  $I(m, \lambda) = \int_{C_{\text{sd}}(m,\lambda)} f_0(z; m, \lambda) dz$ ,  $(m, \lambda) \in W$ . We follow a standard and completely elementary procedure for calculating such asymptotics. The idea is to obtain a partition of  $I(m, \lambda)$  into parts in such a way that one part constitutes the “main contribution” and the other parts are remainder terms, which can be evaluated easily. The starting point is to expand the preexponential factor  $\kappa_0$  of the integrand  $f_0(z) = \kappa_0(z) e^{rQ_m(z)}$  around the stationary point of  $Q_m$ .

Let us first give some preliminary remarks. By (88) we can choose a  $\rho > 0$  such that  $\kappa_0$  is analytic on  $D(z_{\text{st}}(m, \lambda), 4\rho)$  for all  $(m, \lambda) \in W$ . In fact we can use any  $\rho \leq \eta_0/4$ . By (81) there exists a constant  $C_{\rho}$  such that

$$|q(z + z_{\text{st}}(m, \lambda); \lambda)| \leq C_{\rho} \quad (121)$$

for all  $(m, \lambda) \in \mathbb{Z} \times \Lambda$  and all  $z \in D(0, 4\rho)$ . Let in the following  $(m, \lambda) \in W$  be fixed and write  $z_{\text{st}}$  for  $z_{\text{st}}(m, \lambda)$ . Expand  $\kappa_0$  as a power series around  $z_{\text{st}}$

$$\kappa_0(z; \delta) = \sum_{j=0}^{\infty} a_j(\delta) (z - z_{\text{st}})^j \quad (122)$$

for  $z \in D(z_{\text{st}}, 4\rho)$ . Moreover, let  $N \in \mathbb{Z}_{\geq 0}$  be arbitrary but fixed in the following and write

$$\kappa_0(z; \delta) = \sum_{j=0}^N a_j(\delta) (z - z_{\text{st}})^j + R_N(z; \delta). \quad (123)$$



We claim that the main contribution to an order of  $N$  of the integral  $\int_{C_{\text{sd}}(m,\lambda)} f_0(z; m, \lambda) dz$  is given by

$$I(N; \delta) = I(N; m, \lambda, \delta) = \int_{C_{\text{sd}}(m,\lambda)} \kappa_0(z; N; \delta) e^{rQ_m(z)} dz,$$

where  $\kappa_0(z; N; \delta) = \sum_{j=0}^N a_j(\delta)(z - z_{\text{st}})^j$ . Accordingly, the remainder term is given by

$$\epsilon(N; \delta) = \epsilon(N; m, \lambda, \delta) = I(m, \lambda) - I(N; m, \lambda, \delta).$$

At a first glance this can seem a little peculiar since the power series of  $\kappa_0$  is only defined in a small neighborhood around  $z_{\text{st}}$ . The point is that the main contribution comes from integrating  $\kappa_0(z; N; \delta)$  along the part of  $C_{\text{sd}}(m, \lambda)$  lying inside  $D(z_{\text{st}}, 4\rho)$ . However, by integrating along all of  $C_{\text{sd}}(m, \lambda)$  we obtain an explicit and nice expression for the main contribution. By (92) we have

$$\begin{aligned} I(N; \delta) &= \sum_{j=0}^N a_j(\delta) \int_{C_{\text{sd}}(m,\lambda)} (z - z_{\text{st}})^j e^{rQ_m(z)} dz \\ &= \sum_{j=0}^N a_j(\delta) \left( e^{\frac{i\pi}{4} \text{sign}(A)} \right)^{j+1} e^{-irAz_{\text{st}}^2} \int_{-\infty}^{\infty} t^j e^{-r|A|t^2} dt \\ &= 2 \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} a_j(\delta) \left( e^{\frac{i\pi}{4} \text{sign}(A)} \right)^{j+1} e^{-irAz_{\text{st}}^2} \int_0^{\infty} t^j e^{-r|A|t^2} dt \\ &= \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} a_j(\delta) \left( e^{\frac{i\pi}{4} \text{sign}(A)} \right)^{j+1} e^{-irAz_{\text{st}}^2} \Gamma\left(\frac{j+1}{2}\right) \left(\frac{1}{r|A|}\right)^{\frac{j+1}{2}}. \end{aligned}$$

Let us next describe  $\epsilon(N; \delta)$ . Note that for any nice function  $g : \mathbb{R} \rightarrow \mathbb{C}$  and any  $0 \leq a < b \leq \infty$  we have  $\int_{-b}^{-a} g(t) e^{-r|A|t^2} dt = \int_a^b g(-t) e^{-r|A|t^2} dt$ . Thus

$$\epsilon(N; \delta) = e^{\frac{i\pi}{4} \text{sign}(A)} e^{-irAz_{\text{st}}^2} \sum_{\nu \in \{0,1\}} \left( E_{\nu}(N; \delta) + J_{\nu}(\delta) - \sum_{j=0}^N a_j(\delta) F_{\nu}^j \right),$$

where

$$\begin{aligned} J_{\nu}(\delta) &= \int_{2\rho}^{\infty} \psi((-1)^{\nu} t; \delta) \exp(-r|A|t^2) dt, \\ E_{\nu}(N; \delta) &= \int_0^{2\rho} R_N(\gamma_{(m,\lambda)}((-1)^{\nu} t); \delta) \exp(-r|A|t^2) dt, \end{aligned}$$

for  $\nu = 0, 1$ , and  $F_{\nu}^j = \int_{2\rho}^{\infty} (\gamma_{(m,\lambda)}((-1)^{\nu} t) - z_{\text{st}})^j \exp(-r|A|t^2) dt$  for  $\nu = 0, 1$  and  $j = 0, 1, \dots, N$ . By using the incomplete gamma function  $\Gamma(\alpha, u_0) = \int_{u_0}^{\infty} u^{\alpha-1} e^{-u} du$  we get

$$F_{\nu}^j = \frac{1}{2} \left( (-1)^{\nu} e^{\frac{i\pi}{4} \text{sign}(A)} \right)^j \left( \frac{1}{r|A|} \right)^{\frac{j+1}{2}} \Gamma\left(\frac{j+1}{2}, 4|A|\rho^2 r\right).$$

Using that  $F_0^j + F_1^j = (1 + (-1)^j) F_0^j$  and  $e^{\frac{i\pi}{4} \text{sign}(A)} \left( e^{\frac{i\pi}{4} \text{sign}(A)} \right)^{2j} |A|^{-\frac{2j+1}{2}} = (i/A)^{\frac{2j+1}{2}}$  we get

$$I(N; \delta) = e^{-irAz_{\text{st}}^2} \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} a_j(\delta) \Gamma\left(\frac{j+1}{2}\right) \left(\frac{i}{rA}\right)^{\frac{j+1}{2}},$$

$$\begin{aligned} \epsilon(N; \delta) = e^{-irAz_{\text{st}}^2} & \left\{ e^{\frac{i\pi}{4}\text{sign}(A)} \sum_{\nu=0,1} (E_\nu(N; \delta) + J_\nu(\delta)) \right. \\ & \left. - \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} a_j(\delta) \left( \frac{i}{rA} \right)^{\frac{j+1}{2}} \Gamma\left(\frac{j+1}{2}, 4|A|\rho^2 r\right) \right\}. \end{aligned}$$

In accordance with the above we put

$$\begin{aligned} Z_{\text{int},1}(N; \delta) &= \sum_{(m,\lambda) \in W} g(\lambda) I(N; m, \lambda, \delta), \\ R_1(N; \delta) &= Z_{\text{int},1}(\delta) - Z_{\text{int},1}(N; \delta) = \sum_{(m,\lambda) \in W} g(\lambda) \epsilon(N; m, \lambda, \delta). \end{aligned}$$

This decomposition of  $Z_{\text{int},1}(\delta)$  is justified by the following lemma.

**Lemma 5.13.** *For any  $N \in \mathbb{Z}_{\geq 0}$  and  $\delta > 0$  the infinite series*

$$\begin{aligned} Z_{\text{int},1}(N; \delta) &= \sum_{(m,\lambda) \in W} g(\lambda) \exp(-irAz_{\text{st}}^2) \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} a_j(\delta) \Gamma\left(\frac{j+1}{2}\right) \left(\frac{i}{rA}\right)^{\frac{j+1}{2}} \\ \Sigma^1(N; \delta) &= \sum_{(m,\lambda) \in W} g(\lambda) \exp(-irAz_{\text{st}}^2) \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} a_j(\delta) \left(\frac{i}{rA}\right)^{\frac{j+1}{2}} \Gamma\left(\frac{j+1}{2}, 4|A|\rho^2 r\right), \\ \Sigma_\nu^2(N; \delta) &= \sum_{(m,\lambda) \in W} g(\lambda) \exp(-irAz_{\text{st}}^2) E_\nu(N; \delta), \quad \nu = 0, 1, \end{aligned}$$

are absolutely convergent. Moreover, there exists a  $r_0 \in \mathbb{Z}_{\geq 2}$  and a  $\delta_0 > 0$  such that the infinite series

$$\Sigma_\nu^3(\delta) = \sum_{(m,\lambda) \in W} g(\lambda) \exp(-irAz_{\text{st}}^2) J_\nu(\delta), \quad \nu = 0, 1,$$

are absolutely convergent for all  $\delta \in ]0, \delta_0]$  and  $r \in \mathbb{Z}_{\geq r_0}$ . For all such  $\delta$  and  $r$  and all  $N \in \mathbb{Z}_{\geq 0}$  we therefore have  $Z_{\text{int},1}(\delta) = Z_{\text{int},1}(N; \delta) + R_1(N; \delta)$ , where

$$R_1(N; \delta) = -\Sigma^1(N; \delta) + e^{\frac{i\pi}{4}\text{sign}(A)} \sum_{\nu=0,1} (\Sigma_\nu^2(N; \delta) + \Sigma_\nu^3(\delta)).$$

The proof is given in Appendix D. By the following lemma we calculate one of the two main contribution to  $Z_{\text{int}}(r)$  in (107) to arbitrary high order  $N \in \mathbb{Z}_{\geq 0}$ . Recall here that in general we allow  $g(\lambda)$  to depend on  $r$ . By the above lemma we can write

$$R_1(N; \delta) = \sum_{\lambda \in \Lambda} g(\lambda) \sum_{m \in \mathbb{Z}: (m,\lambda) \in W} e^{-irAz_{\text{st}}^2} D(N, \lambda; m, \delta)$$

for  $r \in \mathbb{Z}_{\geq r_0}$  and  $\delta \in ]0, \delta_0]$ , where

$$\begin{aligned} D(N, \lambda; m, \delta) &= - \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} a_j(\delta) \left(\frac{i}{rA}\right)^{\frac{j+1}{2}} \Gamma\left(\frac{j+1}{2}, 4|A|\rho^2 r\right) \\ &\quad + e^{\frac{i\pi}{4}\text{sign}(A)} \sum_{\nu \in \{0,1\}} (E_\nu(N; \delta) + J_\nu(\delta)). \end{aligned}$$

**Proposition 5.14.** *For each fixed  $\lambda \in \Lambda$  and  $j \in \mathbb{Z}_{\geq 0}$  the functions*

$$G_j(m, \lambda) = \exp(-irAz_{\text{st}}(m, \lambda)^2) \partial_z^{(j)} \frac{q(z)}{\sin^k(\pi z)} \Big|_{z=z_{\text{st}}(m, \lambda)} \quad (124)$$

*are periodic in  $m$  with a period  $M_\lambda$  independent of  $j \in \mathbb{Z}_{\geq 0}$ . Moreover, for each such set of periods  $\{M_\lambda\}_{\lambda \in \Lambda}$  and each  $N = 0, 1, 2, \dots$  we have*

$$\begin{aligned} Z_{\text{int},1}(r; N) &:= r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} Z_{\text{int},1}(N; \delta) \\ &= \sqrt{\frac{ri}{\pi A}} \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} \frac{1}{(j/2)!} \left(\frac{i}{4r}\right)^{\frac{j}{2}} \sum_{\lambda \in \Lambda} \frac{|A|}{M_\lambda} A^{-\frac{j}{2}} g(\lambda) \\ &\quad \times \sum_{\substack{m \in \mathbb{Z} \bmod M_\lambda \\ z_{\text{st}}(m, \lambda) \in \mathbb{R} \setminus \mathbb{Z}}} \exp(-irAz_{\text{st}}^2(m, \lambda)) \partial_z^{(j)} \frac{q(z)}{\sin^k(\pi z)} \Big|_{z=z_{\text{st}}(m, \lambda)}. \end{aligned} \quad (125)$$

*Finally, let  $\delta_0 > 0$  and  $r_0 \in \mathbb{Z}_{\geq 2}$  be as in Lemma 5.13. For each  $N = 0, 1, 2, \dots$  and  $\lambda \in \Lambda$  there exists a positive function  $\delta \mapsto \varepsilon_1(N, \lambda; \delta)$  on  $]0, \delta_0]$  (also depending on  $r$ ) such that*

$$\sum_{m \in \mathbb{Z}: (m, \lambda) \in W} |D(N, \lambda; m, \delta)| \leq \varepsilon_1(N, \lambda; \delta),$$

*for all  $\delta \in ]0, \delta_0]$  and all  $r \in \mathbb{Z}_{\geq r_0}$  and such that  $r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \varepsilon_1(N, \lambda; \delta)$  exists and is  $O(r^{-N/2})$  on  $\mathbb{Z}_{\geq r_0}$ . If  $|g(\lambda)|$  is independent of  $r$  it follows that there for each  $N = 0, 1, 2, \dots$  exists a positive function  $\delta \mapsto \varepsilon_1(N; \delta)$  on  $]0, \delta_0]$  (also depending on  $r$ ) such that*

$$|R_1(N; \delta)| \leq \varepsilon_1(N; \delta)$$

*for  $\delta \in ]0, \delta_0]$  and  $r \in \mathbb{Z}_{\geq r_0}$  and such that  $r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \varepsilon_1(N; \delta)$  exists and is  $O(r^{-N/2})$  on  $\mathbb{Z}_{\geq r_0}$ .*

The proof is given in Appendix D. By this we have finalized the description of  $Z_{\text{int},1}(\delta, \eta)$ 's contribution to the large  $r$  asymptotics of  $Z(r)$ .

Before leaving this section, let us take a closer look at the Seifert case, i.e. the case where  $Z(r) = Z(X; r)$ . Let us find periods  $M_\lambda$  as in Proposition 5.14. Recall that  $\mathcal{A} = \prod_{j=1}^n \alpha_j > 0$ . Put  $P = 2\mathcal{A}$  and

$$H = \mathcal{A}E = -\beta_0 \mathcal{A} - \sum_{j=1}^n \frac{\mathcal{A}}{\alpha_j} \beta_j = -\beta_0 \alpha_1 \dots \alpha_n - \sum_{j=1}^n \alpha_1 \dots \alpha_{j-1} \beta_j \alpha_{j+1} \dots \alpha_n.$$

Then  $A = \pi H/P$ . Note that  $|H|$  is the order of the torsion part of  $H_1(X; \mathbb{Z})$  in case of orientable base, i.e  $\epsilon = 0$ , cf. [JN, Corollary 6.2]. For the following we refer to (79). By (82) we have that

$$z_{\text{st}}(m, \underline{n}) = -\frac{P}{H} \left( m - \sum_{j=1}^n \frac{n_j}{\alpha_j} \right)$$

for  $(m, \underline{n}) \in \mathbb{Z} \times \mathbb{Z}^n$ . (Thus this is independent of  $\underline{\mu} \in \{\pm 1\}^n$ .) For each fixed  $\underline{n} \in \mathbb{Z}^n$  the function  $m \mapsto \exp(-irAz_{\text{st}}(m, \underline{n})^2)$  is periodic with a period of  $|H|$ . This follows by the facts that  $z_{\text{st}}(m + H, \underline{n}) = z_{\text{st}}(m, \underline{n}) - P$  and  $P \in 2\mathbb{Z}$ . The functions

$$h_{\underline{\mu}}(z) = \frac{q(z; \underline{\mu})}{\sin^{n+a_\epsilon g-2}(\pi z)}$$

are periodic with a period of  $P$  for any  $\underline{\mu} \in \{\pm 1\}^n$ , and therefore the derivatives of  $h_{\underline{\mu}}$  in  $z_{\text{st}}(m, \underline{n})$  are periodic in  $m$  with a period of  $|H|$ . We can therefore put  $M_{(\underline{\mu}, \underline{n})} = |H|$  for each  $(\underline{\mu}, \underline{n}) \in \Lambda$  and get from (125) that

$$Z_{\text{int},1}(r; N) = \frac{1}{2\mathcal{A}} \left(\frac{i}{2}\right)^n \sqrt{\frac{2ir}{E}} \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} \frac{1}{(j/2)!} \left(\frac{i}{2\pi E}\right)^{\frac{j}{2}} r^{-\frac{j}{2}} \sum_{\substack{\underline{n} \in \mathcal{S} \\ z_{\text{st}} \in \mathbb{R} \setminus \mathbb{Z}}} \sum_{m \in \mathbb{Z} \bmod |H|} \exp(2\pi i r q_{(m, \underline{n})}) \\ \times \sum_{\underline{\mu} \in \{\pm 1\}^n} \left(\prod_{j=1}^n \mu_j\right) \exp\left(2\pi i \sum_{j=1}^n \mu_j \frac{\rho_j n_j}{\alpha_j}\right) \partial_z^{(j)} \frac{\exp\left(-i\pi \left(\sum_{j=1}^n \frac{\mu_j}{\alpha_j}\right) z\right)}{\sin^{n+a_\epsilon g-2}(\pi z)} \Big|_{z=z_{\text{st}}},$$

where  $q_{(m, \underline{n})}$  is given by (39). By calculating the sum over  $\underline{\mu}$  we get

$$Z_{\text{int},1}(r; N) = \frac{(-1)^n}{2\mathcal{A}} \sqrt{\frac{2ir}{E}} \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} \frac{1}{(j/2)!} \left(\frac{i}{2\pi E}\right)^{\frac{j}{2}} r^{-\frac{j}{2}} \\ \times \sum_{\substack{\underline{n} \in \mathcal{S} \\ z_{\text{st}} \in \mathbb{R} \setminus \mathbb{Z}}} \sum_{m \in \mathbb{Z} \bmod |H|} \exp(2\pi i r q_{(m, \underline{n})}) c_{(m, \underline{n})}^{(j)},$$

where we for  $j \in \{0, 1, 2, \dots\}$  and  $(m, \underline{n}) \in \mathbb{Z} \times \mathbb{Z}^n$  have put

$$c_{(m, \underline{n})}^{(j)} = \partial_z^{(j)} \left\{ \frac{\prod_{j=1}^n \sin\left(\frac{\pi}{\alpha_j} [2\rho_j n_j - z]\right)}{\sin^{n+a_\epsilon g-2}(\pi z)} \right\} \Big|_{z=z_{\text{st}}(m, \underline{n})}.$$

In particular,

$$Z_{\text{int},1}(r; N) = \frac{(-1)^n}{2\mathcal{A}} \sqrt{\frac{2ir}{E}} \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} \frac{1}{(j/2)!} \left(\frac{i}{2\pi E}\right)^{\frac{j}{2}} r^{-\frac{j}{2}} \\ \times \sum_{\substack{\underline{n} \in \mathcal{S} \\ z_{\text{st}} \in ]0, 2\mathcal{A}[\cap(\mathbb{R} \setminus \mathbb{Z})}}} \sum_{m \in \mathbb{Z}} \exp(2\pi i r q_{(m, \underline{n})}) c_{(m, \underline{n})}^{(j)}.$$

In this expression certain symmetries are present. Let  $\underline{e}_1, \dots, \underline{e}_n$  be the standard basis in  $\mathbb{R}^n$  and let

$$\begin{aligned} h_j(m, \underline{n}) &= (m+1, \underline{n} + \alpha_j \underline{e}_j), \quad j = 1, 2, \dots, n \\ f(m, \underline{n}) &= (m, \underline{n} - \underline{\beta}), \\ g(m, \underline{n}) &= -(m, \underline{n}) \end{aligned}$$

for  $(m, \underline{n}) \in \mathbb{Z} \times \mathbb{Z}^n$  and  $j = 1, 2, \dots, n$ . The function  $z_{\text{st}}$  is invariant under the transformations  $h_j$ , change sign under  $g$  and  $z_{\text{st}}(f(m, \underline{n})) = z_{\text{st}}(m, \underline{n}) + 2$ . Moreover,  $q_{(m, \underline{n})}$  and  $c_{(m, \underline{n})}^{(j)}$ ,  $j \in 2\mathbb{Z}_{\geq 0}$ , are invariant under all the above transformations (and their inverses). For  $a \in \mathbb{Z}$  put

$$W_a = \{(m, \underline{n}) \in \mathbb{Z} \times \mathcal{S} \mid z_{\text{st}}(m, \underline{n}) \in ]2a, 2(a+1)[ \setminus \{2a+1\}\}.$$

For any  $a \in \mathbb{Z}$  there is a bijection between  $W_a$  and  $W_0$ ; namely  $(m, \underline{n}) \in W_a$  is mapped to  $h_1^{k_1} \dots h_n^{k_n} f^{-a}(m, \underline{n})$ , where  $(k_1, \dots, k_n) \in \mathbb{Z}^n$  is the unique tuple of integers such that  $n_j + a\beta_j + k_j\alpha_j \in \{0, 1, \dots, \alpha_j - 1\}$ . Finally we partition  $W_0$  into the two sets

$$W_0^1 = \{(m, \underline{n}) \in \mathbb{Z} \times \mathcal{S} \mid z_{\text{st}}(m, \underline{n}) \in ]0, 1[\}.$$

and  $W_0^2 = W_0 \setminus W_0^1$  and observe that there is a bijection between  $W_0^1$  and  $W_0^2$  constructed using  $fg$  and the  $h_j$ 's. We have thus shown

**Corollary 5.15.** *Let  $Z(r) = Z(X; r)$ . Then*

$$Z_{\text{int},1}(r; N) = \sqrt{r} \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} \sum_{(m, \underline{n}) \in \mathcal{I}_1} \exp(2\pi i r q_{(m, \underline{n})}) c_{j/2}^{(m, \underline{n})} r^{-\frac{j}{2}},$$

where  $q_{(m, \underline{n})}$  is given by (39) and  $c_{j/2}^{(m, \underline{n})}$  is given by (49).

Finally observe that  $|g(\underline{\mu}, \underline{n})| = 2^{-n}$  for all  $(\underline{\mu}, \underline{n}) \in \Lambda$  by (79). Thus the estimate for  $R_1(N; \delta)$  given at the end of Proposition 5.14 is valid.

**5.1.4. An asymptotic description of  $Z_{\text{int},2}(\delta, \eta)$ 's contribution to  $Z_{\text{int}}(r)$ .** We will follow the same strategy as in the analysis of  $Z_{\text{int},1}(\delta, \eta)$ , i.e. we will partition the expression  $Z_{\text{int},2}(\delta, \eta)$  into simple and manageable pieces, one piece giving the main contribution and the remaining ones adding up to a remainder part, which is small in the large  $r$  limit compared to the main contribution piece.

Let us in the following assume that  $q(m; \lambda) \neq 0$  for all  $m \in \mathbb{Z}$  and all  $\lambda \in \Lambda$ . By periodicity and continuity of  $q$  it then follows that there exists a  $\rho > 0$  such that  $q(z; \lambda) \neq 0$  for all  $z \in \cup_{m \in \mathbb{Z}} D(m, \rho)$  and all  $\lambda \in \Lambda$ . Assume also that the positive parameter  $\eta < \rho$ .

Let  $\lambda \in \Lambda$  be arbitrary but fixed and suppress  $\lambda$  from the notation everywhere. By (101) we have

$$F^+(y; l, \delta, \eta) = \exp(-irA\eta^2) G^+(y; l, \delta, \eta) \exp(irAy^2),$$

where

$$G(y; l, \delta, \eta) = e^{-\pi\delta(y+i\eta+l)^2} \frac{q(y+i\eta+l) \exp(-2rA\eta y)}{\sin^k(\pi y)}$$

and  $G^+$  is the even part of  $G$ , i.e.  $G^+(y; l, \delta, \eta) = (G(y; l, \delta, \eta) + G(-y; l, \delta, \eta))/2$ . We note that  $G^+$  is entire if  $k \leq 1$  and analytic on  $D'(0, 1)$  with a pole in 0 of order  $m > 0$  if  $k > 1$ , where  $m$  is the biggest even integer less than or equal to  $k$ . Write  $G(y; l, \delta, \eta) = \sum_{s=-k}^{\infty} c_s(l, \delta, \eta) y^s$  for  $y \in D'(0, 1)$ . Then

$$G^+(y; l, \delta, \eta) = \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^{\infty} d_{\nu}(l, \delta, \eta) y^{\nu-k},$$

where

$$d_{\nu}(l, \delta, \eta) = \frac{1}{\pi^k \nu!} \partial_y^{(\nu)} \Psi(y; l, \delta, \eta) \Big|_{y=0}, \quad (126)$$

where

$$\Psi(y; l, \delta, \eta) = e^{-\pi\delta(y+i\eta+l)^2} q(y+i\eta+l) \exp(-2rA\eta y) \left( \frac{\pi y}{\sin(\pi y)} \right)^k. \quad (127)$$

By (102) the relevant integrals to examine are  $\int_{C_{\eta}} F^+(y; l, \delta, \eta) dy = \exp(-irA\eta^2) I(l, \delta, \eta)$ , where we write  $C_{\eta}$  for  $C(\lambda, \eta)$  and  $I(l, \delta, \eta) = \int_{C_{\eta}} G^+(y; l, \delta, \eta) \exp(irAy^2) dy$ . Let  $N \in \mathbb{Z}_{\geq 0}$  be arbitrary but fixed in the following and write

$$G^+(y; l, \delta, \eta) = \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N d_{\nu}(l, \delta, \eta) y^{\nu-k} + R_N(y; l, \delta, \eta)$$

for  $y \in D'(0, 1)$ . Put

$$\begin{aligned} I(N; l, \delta, \eta) &= \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N d_\nu(l, \delta, \eta) \int_{C_\eta} y^{\nu-k} e^{irAy^2} dy, \\ \epsilon(N; l, \delta, \eta) &= I(l, \delta, \eta) - I(N; l, \delta, \eta). \end{aligned}$$

To describe  $\epsilon(N; l, \delta, \eta)$ , let  $C_\eta(t) = e^{\frac{i\pi}{4}\text{sign}(A)}t - i\eta$  be a parametrization of  $C_\eta$ , and let  $C_\eta^0$  and  $C_\eta^\infty$  be the restrictions of  $C_\eta(t)$  to  $[-1/2, 1/2]$  and  $\mathbb{R} \setminus ]-1/2, 1/2[$  respectively and put

$$\begin{aligned} I_\infty(l, \delta, \eta) &= \int_{C_\eta^\infty} G^+(y; l, \delta, \eta) e^{irAy^2} dy, \\ J_1(N; l, \delta, \eta) &= \int_{C_\eta^0} R_N(y; l, \delta, \eta) e^{irAy^2} dy, \\ J_2(N; l, \delta, \eta) &= - \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N d_\nu(l, \delta, \eta) \int_{C_\eta^\infty} y^{\nu-k} e^{irAy^2} dy. \end{aligned}$$

Then

$$\epsilon(N; l, \delta, \eta) = I_\infty(l, \delta, \eta) + J_1(N; l, \delta, \eta) + J_2(N; l, \delta, \eta).$$

For  $\lambda \in \Lambda$  put

$$I_\lambda = \left\{ l \in \mathbb{Z} \mid \frac{B(\lambda)}{2\pi} + \frac{A(\lambda)}{\pi} l \in \mathbb{Z} \right\}.$$

**Lemma 5.16.** *There exists a  $\eta_2 > 0$  such that the infinite series*

$$\begin{aligned} Z_{\text{int},2}(N; \delta, \eta) &= \sum_{\lambda \in \Lambda} g(\lambda) e^{-irA\eta^2} \sum_{l \in I_\lambda} \beta(l) I(N; l, \delta, \eta), \\ \Sigma^4(N; \delta, \eta) &= \sum_{\lambda \in \Lambda} g(\lambda) e^{-irA\eta^2} \sum_{l \in I_\lambda} \beta(l) J_1(N; l, \delta, \eta) \\ \Sigma^5(N; \delta, \eta) &= \sum_{\lambda \in \Lambda} g(\lambda) e^{-irA\eta^2} \sum_{l \in I_\lambda} \beta(l) J_2(N; l, \delta, \eta) \end{aligned}$$

are absolutely convergent for all  $\delta > 0$ , all  $\eta \in ]0, \eta_2]$  and all  $N \in \mathbb{Z}_{\geq 0}$ . We can choose  $\eta_2 > 0$  such that all the above series are uniformly convergent w.r.t.  $\eta$  on  $]0, \eta_2]$  for each fixed  $\delta > 0$  and  $N \in \mathbb{Z}_{\geq 0}$ .

Moreover, by choosing  $\eta_2$  sufficiently small, there exists a  $r_0 \in \mathbb{Z}_{\geq 2}$  and a  $\delta_0 \in ]0, \infty[$  such that

$$\Sigma^6(\delta, \eta) = \sum_{\lambda \in \Lambda} g(\lambda) e^{-irA\eta^2} \sum_{l \in I_\lambda} \beta(l) I_\infty(l, \delta, \eta)$$

is absolutely convergent for all  $\delta \in ]0, \delta_0]$ ,  $\eta \in ]0, \eta_2]$  and  $r \in \mathbb{Z}_{\geq r_0}$ , and such that  $\Sigma^6(\delta, \eta)$  is uniformly convergent w.r.t.  $\eta$  on  $]0, \eta_2]$  for each fixed  $\delta \in ]0, \delta_0]$  and  $r \in \mathbb{Z}_{\geq r_0}$ .

The proof is given in Appendix E. By the above lemma (and proof of that lemma) we get

$$\begin{aligned} Z_{\text{int},2}(N; \delta) &:= \lim_{\eta \rightarrow 0_+} Z_{\text{int},2}(N; \delta, \eta) \\ &= \sum_{\lambda \in \Lambda} g(\lambda) \frac{e^{\frac{i\pi}{4}\text{sign}(A)}}{\sqrt{r|A|}} \sum_{l \in I_\lambda} \beta(l, \lambda) \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N d_\nu(l, \delta, 0) \left( \frac{i}{rA} \right)^{\frac{\nu-k}{2}} \Gamma\left( \frac{\nu-k+1}{2} \right), \end{aligned}$$

where we have also used (146) and where  $d_\nu(l, \delta, 0)$  is equal to  $d_\nu(l, \delta, \eta)$  with  $\eta = 0$ . Similarly we find that

$$\Sigma^\nu(N; \delta) := \lim_{\eta \rightarrow 0_+} \Sigma^\nu(N; \delta, \eta) = \sum_{\lambda \in \Lambda} g(\lambda) \sum_{l \in I_\lambda} \beta(l) J_{\nu-3}(N; l, \delta)$$

for  $\nu = 4, 5$ , where  $J_k(N; l, \delta)$  is equal to  $J_k(N; l, \delta, \eta)$  with  $\eta = 0$ ,  $k = 1, 2$ . Finally, for  $\delta \in ]0, \delta_0]$  and  $r \in \mathbb{Z}_{\geq r_0}$  we get

$$\Sigma^6(\delta) := \lim_{\eta \rightarrow 0_+} \Sigma^6(\delta, \eta) = \sum_{\lambda \in \Lambda} g(\lambda) \sum_{l \in I_\lambda} \beta(l) I_\infty(l, \delta),$$

where  $I_\infty(l, \delta)$  is equal to  $I_\infty(l, \delta, \eta)$  with  $\eta = 0$ . (That  $\lim_{\eta \rightarrow 0_+} I_\infty(l, \delta, \eta) = I_\infty(l, \delta, 0)$  and  $\lim_{\eta \rightarrow 0_+} J_k(N; l, \delta, \eta) = J_k(N; l, \delta, 0)$  follows by Lebesgue's theorem on dominated convergence.) Thus, for  $\delta \in ]0, \delta_0]$  and  $r \in \mathbb{Z}_{\geq r_0}$ , we have a decomposition

$$Z_{\text{int},2}(\delta) = Z_{\text{int},2}(N; \delta) + R_2(N; \delta),$$

where

$$R_2(N; \delta) = \sum_{\lambda \in \Lambda} g(\lambda) e^{-irA\eta^2} \sum_{l \in I_\lambda} \beta(l) \epsilon(N; l, \delta) = \Sigma^4(N; \delta) + \Sigma^5(N; \delta) + \Sigma^6(\delta),$$

where

$$\epsilon(N; l, \delta) = \lim_{\eta \rightarrow 0_+} \epsilon(N; l, \delta, \eta) = I_\infty(l, \delta) + J_1(N; l, \delta) + J_2(N; l, \delta).$$

The task is now to calculate

$$Z_{\text{int},2}(r; N) := r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} Z_{\text{int},2}(N; \delta)$$

and estimate  $R_2(N; \delta)$  for small  $\delta$  and large  $r$ . For the next proposition, recall that  $\epsilon(N; l, \delta) = \epsilon(N, \lambda; l, \delta)$  depends on  $\lambda \in \Lambda$ .

**Proposition 5.17.** *With notation from above we have*

$$\begin{aligned} Z_{\text{int},2}(r; N) &= \frac{r}{\pi^k} \sum_{\lambda \in \Lambda} \frac{g(\lambda)}{M_\lambda} \frac{e^{\frac{i\pi}{4} \text{sign}(A)}}{\sqrt{r|A|}} \sum_{l \in I_\lambda \bmod M_\lambda} \beta(l, \lambda) \\ &\times \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N \left( \frac{i}{rA} \right)^{\frac{\nu-k}{2}} \frac{\Gamma\left(\frac{\nu-k+1}{2}\right)}{\nu!} \partial_y^{(\nu)} h(y; l) \Big|_{y=0}, \end{aligned} \quad (128)$$

where

$$h(y; l) = q(y + l) \left( \frac{\pi y}{\sin(\pi y)} \right)^k$$

and  $M_\lambda$  is the least common multiplum of 2,  $m_q$ ,  $b(\lambda)$  and  $P(\lambda)$ , where  $m_q$  and  $P(\lambda)$  are the integers in (81) and (87) respectively, and  $b(\lambda)$  is an integer such that  $b(\lambda)B(\lambda)/\pi \in \mathbb{Z}$ , see (89).

Moreover, let  $\delta_0 > 0$  and  $r_0 \in \mathbb{Z}_{\geq 2}$  be as in Lemma 5.16. For each  $N = 0, 1, 2, \dots$  and each  $\lambda \in \Lambda$  there exists a function  $\delta \mapsto \varepsilon_2(N, \lambda; \delta)$  on  $]0, \delta_0]$  (also depending on  $r$ ) such that

$$\sum_{l \in I_\lambda} |\epsilon(N, \lambda; l, \delta)| \leq \varepsilon_2(N, \lambda; \delta),$$

for all  $\delta \in ]0, \delta_0]$  and  $r \in \mathbb{Z}_{\geq r_0}$  and such that  $r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \varepsilon_2(N, \lambda; \delta)$  exists and is  $O\left(r^{\frac{k-\nu_1+1}{2}}\right)$  on  $\mathbb{Z}_{\geq r_0}$ , where  $\nu_1 \in \{N+1, N+2\}$  such that  $\nu_1 - k \in 2\mathbb{Z}$ . If  $|g(\lambda)|$  is independent of  $r$  it follows that there for each  $N = 0, 1, 2, \dots$  exists a function  $\delta \mapsto \varepsilon_2(N; \delta)$  on  $]0, \delta_0]$  (also depending on  $r$ ) such that

$$|R_2(N; \delta)| \leq \varepsilon_2(N; \delta),$$

for all  $\delta \in ]0, \delta_0]$  and  $r \in \mathbb{Z}_{\geq r_0}$  and such that  $r \lim_{\delta \rightarrow 0+} \sqrt{\delta} \varepsilon_2(N; \delta)$  exists and is  $O\left(r^{\frac{k-\nu_1+1}{2}}\right)$  on  $\mathbb{Z}_{\geq r_0}$ , where  $\nu_1$  is as above.

Proposition 5.17 is proved by applications of Lemma 7.1. The estimate on  $R_2(N; \delta, \eta)$  mainly follows from the proof of Lemma 5.16. We give the details in Appendix E.

Let us finally restrict to the Seifert case, where the relevant data are given in (79). Here we have that  $m_q = 2\mathcal{A} = 2 \prod_{j=1}^n \alpha_j$ . For all  $\lambda \in \Lambda$  we can let  $b(\lambda) = \mathcal{A}$  and  $P(\lambda) = 2\mathcal{A}$ . Therefore, we can put  $M_\lambda = 2\mathcal{A}$  in Proposition 5.17 and find that

$$\begin{aligned} Z_{\text{int},2}(r; N) &= \frac{e^{\frac{i\pi}{4} \text{sign}(E)}}{\pi^{n+a_\epsilon g-2}} \sqrt{\frac{2r}{\pi|E|}} \sum_{\underline{\mu} \in \{\pm 1\}^n} \sum_{\underline{n} \in \mathcal{S}} \frac{g(\underline{\mu}, \underline{n})}{2\mathcal{A}} \sum_{\substack{l \in \mathbb{Z} \bmod 2\mathcal{A} \\ \sum_{j=1}^n \frac{n_j + \frac{1}{2}\beta_j l}{\alpha_j} \in \mathbb{Z}}} \beta(l, \underline{n}) \\ &\quad \times \sum_{\substack{\nu=0 \\ \nu-n \in 2\mathbb{Z}}}^N \left( \frac{i2}{r\pi E} \right)^{\frac{\nu-n-a_\epsilon g+2}{2}} \frac{\Gamma\left(\frac{\nu-n-a_\epsilon g+3}{2}\right)}{\nu!} \partial_y^{(\nu)} h(y; l) \Big|_{y=0}, \end{aligned}$$

where

$$h(y; l) = \exp \left( -i\pi \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) (y + l) \right) \left( \frac{\pi y}{\sin(\pi y)} \right)^{n+a_\epsilon g-2}.$$

For  $l \in \mathbb{Z}$  arbitrary but fixed we put  $n'_j = n_j + \frac{1}{2}\beta_j l$  for  $\underline{n} \in \mathbb{Z}^n$  and obtain by (79) and (97) that

$$g(\underline{\mu}, \underline{n}) \beta(l, \underline{n}) e^{-i\pi \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) l} = \left( \frac{i}{2} \right)^n \left( \prod_{j=1}^n \mu_j \right) k(l, \underline{\mu}, \underline{n}'),$$

where

$$k(l, \underline{\mu}, \underline{n}') = (-1)^{nl} e^{2\pi i r q(l, \underline{n}')} \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} n'_j - \frac{1}{2} \sigma_j l \right) \right],$$

where  $q(l, \underline{n}')$  is given by (40). Therefore

$$\begin{aligned} Z_{\text{int},2}(r; N) &= \frac{e^{\frac{i\pi}{4} \text{sign}(E)}}{\pi^{n+a_\epsilon g-2} \mathcal{A}} \left( \frac{i}{2} \right)^n \sqrt{\frac{r}{2\pi|E|}} \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \sum_{\underline{n} \in \mathcal{S}} \sum_{\substack{l \in \mathbb{Z} \bmod 2\mathcal{A} \\ \underline{n}' = \underline{n} + \frac{1}{2}\beta \underline{l} : \sum_{j=1}^n \frac{n'_j}{\alpha_j} \in \mathbb{Z}}} k(l, \underline{\mu}, \underline{n}') \\ &\quad \times \sum_{\substack{\nu=0 \\ \nu-n \in 2\mathbb{Z}}}^N \left( \frac{i2}{r\pi E} \right)^{\frac{\nu-n-a_\epsilon g+2}{2}} \frac{\Gamma\left(\frac{\nu-n-a_\epsilon g+3}{2}\right)}{\nu!} \partial_y^{(\nu)} h(y; 0) \Big|_{y=0}. \end{aligned}$$

Since  $k(l, \underline{\mu}, \underline{n}')$  and the condition  $\sum_{j=1}^n \frac{n'_j}{\alpha_j} \in \mathbb{Z}$  are invariant under the transformations  $\frac{1}{2}\mathbb{Z}^n \rightarrow \frac{1}{2}\mathbb{Z}^n$ ,  $\underline{n}' \mapsto \underline{n}' + \alpha_j \underline{e}_j$ ,  $j = 1, 2, \dots, n$ , we get

$$Z_{\text{int},2}(r; N) = \frac{e^{\frac{i\pi}{4} \text{sign}(E)}}{\pi^{n+a_\epsilon g-2} \mathcal{A}} \left( \frac{i}{2} \right)^n \sqrt{\frac{r}{2\pi|E|}} \sum_{\substack{\nu=0 \\ \nu-n \in 2\mathbb{Z}}}^N \left( \frac{i2}{r\pi E} \right)^{\frac{\nu-n-a_\epsilon g+2}{2}} \frac{\Gamma\left(\frac{\nu-n-a_\epsilon g+3}{2}\right)}{\nu!}$$



$$\times \sum_{l \in \mathbb{Z} \bmod 2A} \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \sum_{\underline{n}' \in J_l : \sum_{j=1}^n \frac{n'_j}{\alpha_j} \in \mathbb{Z}} k(l, \underline{\mu}, \underline{n}') \partial_y^{(\nu)} h(y; 0) \Big|_{y=0},$$

where  $J_l$  is given by (115). For fixed  $(\underline{\mu}, \underline{n}')$  we have that  $k(l, \underline{\mu}, \underline{n}')$  is periodic in  $l$  with a period of 2. Moreover, the sum  $\sum_{\underline{n}' \in J_l : \sum_{j=1}^n \frac{n'_j}{\alpha_j} \in \mathbb{Z}}$  is periodic in  $l$  with a period of 2. Therefore

$$\begin{aligned} Z_{\text{int},2}(r; N) &= \frac{e^{\frac{i\pi}{4}\text{sign}(E)}}{\pi^{n+a_\epsilon g-2}} \left(\frac{i}{2}\right)^n \sqrt{\frac{r}{2\pi|E|}} \sum_{\substack{\nu=0 \\ \nu-n \in 2\mathbb{Z}}}^N \left(\frac{i2}{r\pi E}\right)^{\frac{\nu-n-a_\epsilon g+2}{2}} \frac{\Gamma\left(\frac{\nu-n-a_\epsilon g+3}{2}\right)}{\nu!} \\ &\times \sum_{l \in \mathbb{Z} \bmod 2} \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \sum_{\underline{n}' \in J_l : \sum_{j=1}^n \frac{n'_j}{\alpha_j} \in \mathbb{Z}} k(l, \underline{\mu}, \underline{n}') \partial_y^{(\nu)} h(y; 0) \Big|_{y=0}. \end{aligned}$$

Finally, if  $J'_l$  is the set in (116), we can change the sum  $\sum_{\underline{n}' \in J_l : \sum_{j=1}^n \frac{n'_j}{\alpha_j} \in \mathbb{Z}}$  to

$$\sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\underline{n}' \in J'_l : \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \in \mathbb{Z}} \frac{1}{\prod_{j=1}^n \text{Sym}_{\mathbb{Z}_\pm} \left( \frac{n'_j}{\alpha_j} \right)}$$

if we everywhere change  $n'_j$  to  $\mu'_j n'_j$ . By this we end up with

$$\begin{aligned} Z_{\text{int},2}(r; N) &= \frac{e^{\frac{i\pi}{4}\text{sign}(E)}}{\pi^{n+a_\epsilon g-2}} \left(\frac{i}{2}\right)^n \sqrt{\frac{r}{2\pi|E|}} \sum_{\substack{\nu=0 \\ \nu-n \in 2\mathbb{Z}}}^N \left(\frac{i2}{r\pi E}\right)^{\frac{\nu-n-a_\epsilon g+2}{2}} \frac{\Gamma\left(\frac{\nu-n-a_\epsilon g+3}{2}\right)}{\nu!} \\ &\times \sum_{l \in \mathbb{Z} \bmod 2} \sum_{\underline{\mu} \in \{\pm 1\}^n} \sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\underline{n}' \in J'_l : \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \in \mathbb{Z}} \frac{1}{\prod_{j=1}^n \text{Sym}_{\mathbb{Z}_\pm} \left( \frac{n'_j}{\alpha_j} \right)} \\ &\times \left( \prod_{j=1}^n \mu_j \right) (-1)^{nl} \exp(2\pi i r q(l, \underline{n}')) \\ &\times \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \partial_y^{(\nu)} h(y; 0) \Big|_{y=0}. \end{aligned}$$

By evaluating the sum over  $\underline{\mu}$  we arrive at

**Corollary 5.18.** *Let  $Z(r) = Z(X; r)$ . Then*

$$Z_{\text{int},2}(r; N) = \sqrt{r} \sum_{\substack{\nu=0 \\ \nu-n \in 2\mathbb{Z}}}^N \sum_{(l, \underline{n}') \in \mathcal{I}_2} \exp(2\pi i r q(l, \underline{n}')) c_{\nu'}^{(l, \underline{n}')} r^{-\nu'}$$

where  $q(l, \underline{n}')$  is given by (40) and  $c_{\nu'}^{(l, \underline{n}')}$  is given by (50), where  $\nu' = (\nu - n - a_\epsilon g + 2)/2$ .  $\square$

As already noted after Corollary 5.15 we have that  $|g(\underline{\mu}, \underline{n})| = 2^{-n}$  for all  $(\underline{\mu}, \underline{n}) \in \Lambda$  so the estimate for  $R_2(N; \delta, \eta)$  given at the end of Proposition 5.17 is valid.

## 6. PROOF OF THEOREM 4.4

In this section we calculate the large  $r$  asymptotics of  $Z(r)$  in (76) in case  $A = 0$ . Like in the case  $A \neq 0$  one can calculate this asymptotics by finding a good deformation of the integration contour (the real axes) of the integral

$$I(m, \lambda) = \int_{-\infty}^{\infty} f(z; m, \lambda) dz$$

to some other contour (depending on  $(m, \lambda)$ ). However, in this case we do not have a natural choice of a good such deformation, since the steepest descent method is not applicable. In the case  $A \neq 0$  we saw that by deforming the real axes in  $I(m, \lambda)$  to the steepest descent contour we obtain an integrand with an exponential decay of the form  $e^{-ct^2}$  with  $c$  a positive constant. It turns out that we can again obtain integrands with an exponential decay by deforming the real axes to a line parallel to the real line and lying in the upper or lower half plane depending on the sign of a certain parameter. Like in the case  $A \neq 0$  this leads to a decomposition of  $Z(r)$  into a residue part and a part being equal to a limit of a certain infinite sum of integrals. If  $a_0 - k\pi < 0$  we can show that this limit is equal to zero,  $a_0$  being the constant in (80). However, if  $a_0 - k\pi \geq 0$ , we have not been able to find a nice reduction of the expression involving the infinite sum of integrals. Let us give the details. Let  $q(z)$  be an  $r$  independent function satisfying (80) and (81). Let us write  $q(z) = q_0(z)q_1(z)$ , where

$$q_0(z; \lambda) = e^{\pi i \zeta(\lambda) z} \quad (129)$$

with  $\zeta(\lambda) \in \mathbb{Q}$  for all  $\lambda \in \Lambda$ . In particular  $q_1$  is independent of  $r$  and satisfies (80) and (81) with the numbers  $m_q$ ,  $a_n$  and  $b_n$  replaced by some other numbers  $m'_q = m_{q_1}$ ,  $a'_n$  and  $b'_n$ . We will use this splitting in a flexible way. In particular we are interested in the extreme cases, with  $q_0$  respectively  $q_1$  being constantly one. We have

$$f(z; m, \lambda) = e^{-\pi \delta z^2} q_1(z) \frac{\exp\left(2\pi i r \left[m + \frac{B}{2\pi} + \frac{1}{2r}\zeta\right] z\right)}{\sin^k(\pi(z - i\eta))}$$

with  $B = B(\lambda)$  and  $\zeta = \zeta(\lambda)$ , so

$$Z(r) = r \lim_{\delta \rightarrow 0+} \sqrt{\delta} \lim_{\eta \rightarrow 0+} \sum_{m \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) \int_{-\infty}^{\infty} e^{-\pi \delta z^2} q_1(z) \frac{e^{2\pi i r \nu(m, \lambda) z}}{\sin^k(\pi(z - i\eta))} dz,$$

where  $\nu(m, \lambda) = m + \frac{B}{2\pi} + \frac{1}{2r}\zeta$ . For  $R \in \mathbb{Z} + \frac{1}{2}$ ,  $\nu \in \mathbb{R}$  and  $\rho \in ]0, \infty[$ , let  $\gamma(t) = \gamma_{(\nu, \rho, R)}(t) = R + i \operatorname{sign}(\nu)t$ ,  $t \in [0, \rho]$ . Here  $\operatorname{sign}(0) \in \{\pm 1\}$  (both options will be used). Then

$$\begin{aligned} \int_{\gamma} f(z; m, \lambda) dz &= i \operatorname{sign}(\nu) e^{2\pi i r \nu R} e^{-\pi \delta R^2} \\ &\times \int_0^{\rho} e^{\pi \delta t^2} q_1(R + i \operatorname{sign}(\nu)t) \frac{e^{-i 2\pi \delta R \operatorname{sign}(\nu)t} e^{-2\pi t|\nu|t}}{\sin^k(\pi(R + i(\operatorname{sign}(\nu)t - \eta)))} dt, \end{aligned}$$

so

$$\left| \int_{\gamma} f(z; m, \lambda) dz \right| \leq b'_0 e^{-\pi \delta R^2} \int_0^{\rho} e^{\pi \delta t^2} \frac{e^{a'_0(m'_q + t)} e^{-2\pi t|\nu|t}}{|\sin(\pi(R + i(\operatorname{sign}(\nu)t - \eta)))|^k} dt.$$

By (132) and our choice of  $R$  we have  $|\sin(\pi(R + i(\operatorname{sign}(\nu)t - \eta)))| \geq 1$  so

$$\left| \int_{\gamma} f(z; m, \lambda) dz \right| \leq b'_0 e^{a'_0 m'_q} e^{-\pi \delta R^2} \int_0^{\rho} e^{\pi \delta t^2} e^{a'_0 t} e^{-2\pi t|\nu|t} dt.$$

Here the integral on the right-hand side is finite and independent of  $R$ , hence  $\int_\gamma f(z; m, \lambda) dz$  converges to zero as  $|R| \rightarrow \infty$ . Now put  $M(m, \lambda) = 0$  if  $\text{sign}(\nu(m, \lambda)) = -1$  and

$$M(m, \lambda) = 2\pi i \sum_{l \in \mathbb{Z}} \text{Res}_{z=l+i\eta} \{f(z; m, \lambda)\}$$

if  $\text{sign}(\nu(m, \lambda)) = +1$ . For the following, let  $\eta_0 > 0$  be arbitrary but fixed, and let  $\kappa_\nu^\rho(t) = t + i\text{sign}(\nu)\rho$ ,  $t \in \mathbb{R}$ , for  $\rho > \eta_0$ . Moreover, let  $J$  be any subset of  $I = \{(m, \lambda) \in \mathbb{Z} \times \Lambda \mid \nu(m, \lambda) = 0\}$  and let  $\text{sign}(\nu(m, \lambda)) = 1$  for  $(m, \lambda) \in J$  and  $\text{sign}(\nu(m, \lambda)) = -1$  for  $(m, \lambda) \in I \setminus J$  and put

$$\begin{aligned} Z_{\text{int}}^J(\delta, \eta) &= \sum_{m \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) \int_{\kappa_\nu^\rho(m, \lambda)} f(z; m, \lambda) dz, \\ Z_{\text{polar}}^J(\delta, \eta) &= \sum_{(m, \lambda) \in \mathbb{Z} \times \Lambda} g(\lambda) M(m, \lambda). \end{aligned}$$

**Lemma 6.1.** *For any  $J \subseteq I$  the infinite series  $Z_{\text{int}}^J(\delta, \eta)$  and  $Z_{\text{polar}}^J(\delta, \eta)$  are absolutely convergent for all  $\delta > 0$  and  $\eta \in ]0, \eta_0]$ . In particular*

$$Z(r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} (Z_{\text{int}}^J(\delta, \eta) + Z_{\text{polar}}^J(\delta, \eta))$$

for any such subset  $J$ .

The proof of this lemma is given in Appendix F. By the estimate (156) the series  $Z_{\text{int}}^J(\delta, \eta)$  is uniformly convergent with respect to  $\eta$  on an interval on the form  $]0, \eta_0]$ . Using this and Lebesgue's dominated convergence theorem we get

$$Z_{\text{int}}^J(\delta) := \lim_{\eta \rightarrow 0_+} Z_{\text{int}}^J(\delta, \eta) = \sum_{m \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) \int_{\kappa_\nu^\rho} f_0(z; m, \lambda) dz,$$

where

$$f_0(z; m, \lambda) = e^{-\pi \delta z^2} \frac{q_1(z) e^{2\pi i r \nu(m, \lambda) z}}{\sin^k(\pi z)}.$$

Let  $d_\rho = \sqrt{e^{2\pi\rho} + e^{-2\pi\rho} - 2}$ . Similar to (156) we have

$$\left| \int_{\kappa_\nu^\rho} f_0(z; m, \lambda) dz \right| \leq b'_0 e^{a'_0(m'_q + \rho)} \frac{1}{\sqrt{\delta}} \left( \frac{2}{d_\rho} \right)^k e^{\pi \delta \rho^2} e^{-2\pi r \rho |\nu|}$$

and then

$$\left| \sqrt{\delta} Z_{\text{int}}^J(\delta) \right| \leq b'_0 e^{a'_0(m'_q + \rho)} \left( \frac{2}{d_\rho} \right)^k e^{\pi \delta \rho^2} \sum_{\lambda \in \Lambda} |g(\lambda)| \sum_{m \in \mathbb{Z}} e^{-2\pi r \rho |m + \frac{B}{2\pi} + \frac{1}{2r} \zeta|}$$

for all  $\rho > 0$ . (Note that this estimate is independent of  $J$ .) Therefore (below we show that  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}^J(\delta, \eta)$  exists, hence that  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} Z_{\text{int}}^J(\sqrt{\delta})$  exists)

$$\left| \lim_{\delta \rightarrow 0_+} \sqrt{\delta} Z_{\text{int}}^J(\delta) \right| \leq b'_0 e^{a'_0(m'_q + \rho)} \left( \frac{2}{d_\rho} \right)^k \sum_{\lambda \in \Lambda} |g(\lambda)| \sum_{m \in \mathbb{Z}} e^{-2\pi r \rho |m + \frac{B}{2\pi} + \frac{1}{2r} \zeta|}$$

Here

$$\begin{aligned} \sum_{m \in \mathbb{Z}} e^{-2\pi r \rho |m + \frac{B}{2\pi} + \frac{1}{2r} \zeta|} &\leq \int_{-\infty}^{\infty} e^{-2\pi r \rho |x|} dx + \max_{x \in \mathbb{R}} \left\{ e^{-2\pi r \rho |x|} \right\} \\ &= 2 \int_0^{\infty} e^{-2\pi r \rho x} dx + 1 = 1 + \frac{1}{\pi r \rho}. \end{aligned}$$

In the limit  $\rho \rightarrow \infty$  we have  $d_\rho \sim e^{\pi\rho}$  so  $e^{a'_0\rho}/d_\rho^k \sim e^{(a'_0-k\pi)\rho}$ , and since  $Z_{\text{int}}^J(\delta)$  is independent of  $\rho \in ]0, \infty[$  we find by letting  $\rho \rightarrow \infty$  that

$$\lim_{\delta \rightarrow 0_+} \sqrt{\delta} Z_{\text{int}}^J(\delta) = 0$$

for any  $J \subseteq I$  if  $a'_0 - k\pi < 0$ . Thus, in that case we have that

$$Z(r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}^J(\delta, \eta)$$

for any  $J$ , hence the right-hand side is independent of  $J$  in case  $a'_0 - k\pi < 0$ . Thus we can conclude that non of the terms with  $(m, \lambda) \in I$  contribute to  $Z(r)$  in that case. For  $(m, \lambda) \in I$  we have

$$\lim_{\eta \rightarrow 0_+} M(m, \lambda) = 2\pi i \sum_{l \in \mathbb{Z}} (-1)^{kl} \text{Res}_{z=0} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l)}{\sin^k(\pi z)} \right\}.$$

Here  $(-1)^{kl} q_1(z+l)$  is periodic in  $l$  with a period of  $2m'_q$ , thus by the standard argument we obtain

$$\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} M(m, \lambda) = \frac{\pi i}{m'_q} \sum_{l=0}^{2m'_q-1} (-1)^{kl} \text{Res}_{z=0} \left\{ \frac{q_1(z+l)}{\sin^k(\pi z)} \right\}$$

and by the above argument this limit is zero in case  $a'_0 - k\pi < 0$ . If  $q_1(z) = 1$  (so  $m'_q = 1$  and  $a'_0 = 0$ ), this is the trivial statement that

$$\sum_{l=0}^1 (-1)^{kl} \text{Res}_{z=0} \left\{ \frac{1}{\sin^k(\pi z)} \right\} = 0.$$

We calculate  $r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}^J(\delta, \eta)$  by using some of the same techniques as in the case  $A \neq 0$ , however, this case is much simpler. Indeed we have

$$\begin{aligned} Z_{\text{polar}}^J(\delta, \eta) &= A_+^J(\delta, \eta) - A_-^J(\delta, \eta) + 2\pi i \sum_{l \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) e^{2\pi i r (\frac{B}{2\pi} + \frac{1}{2r} \zeta) l} (-1)^{kl} \\ &\quad \times \sum_{m \in \mathbb{Z}: \nu(m, \lambda) \geq 0} \frac{1}{\text{Sym}_\pm(\nu(m, \lambda))} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l)}{\sin^k(\pi(z-i\eta))} \right\}, \end{aligned}$$

where

$$A_j^J(\delta, \eta) = \pi i \sum_{(m, \lambda) \in J_j} g(\lambda) \sum_{l \in \mathbb{Z}} (-1)^{kl} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l)}{\sin^k(\pi(z-i\eta))} \right\}$$

for  $j = +, -$ , where  $J_+ = J$  and  $J_- = I \setminus J$ . Since the summation limit in the sum over  $m$  is independent of  $l$  we can immediately use Lemma 5.7 to obtain

$$\begin{aligned} Z_{\text{polar}}^J(\delta, \eta) &= A_+^J(\delta, \eta) - A_-^J(\delta, \eta) + 2\pi i \sum_{l \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) e^{2\pi i r (\frac{B}{2\pi} + \frac{1}{2r} \zeta) l} (-1)^{kl} \\ &\quad \times \left( \frac{i}{2} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l) e^{2\pi i r (\frac{B}{2\pi} + \frac{1}{2r} \zeta) z}}{\sin^k(\pi(z-i\eta))} \cot(\pi r z) \right\} \right. \\ &\quad \left. + \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq \lfloor \frac{B}{2\pi} + \frac{1}{2r} \zeta \rfloor}} \frac{\text{sign}(\frac{B}{2\pi} + \frac{1}{2r} \zeta) \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l) e^{-2\pi i r \text{sign}(\frac{B}{2\pi} + \frac{1}{2r} \zeta)(m - \lfloor \frac{B}{2\pi} + \frac{1}{2r} \zeta \rfloor) z}}{\sin^k(\pi(z-i\eta))} \right\}}{\text{Sym}_\pm(m) \text{Sym}_\pm(m - \lfloor \frac{B}{2\pi} + \frac{1}{2r} \zeta \rfloor)} \right). \end{aligned}$$

Like in the case  $A \neq 0$  we get by uniform convergence that

$$Z_{\text{polar}}^J(\delta) := \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}^J(\delta, \eta) = Z_{\text{polar}}^J(\delta, 0).$$

Finally, since  $B(\lambda)$  and  $\zeta(\lambda)$  are rational for all  $\lambda \in \Lambda$  we arrive via Lemma 7.1 at

**Theorem 6.2.** *The limit  $Z_{\text{polar}}^J(r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} Z_{\text{polar}}^J(\delta)$  exists. In fact, there exists a family of positive integers  $\{M_\lambda\}_{\lambda \in \Lambda}$  such that*

$$\begin{aligned} Z_{\text{polar}}^J(r) = & r \left( A_+^J(r) - A_-^J(r) \right) + 2\pi i r \sum_{\lambda \in \Lambda} \frac{g(\lambda)}{M_\lambda} \sum_{l \in \mathbb{Z} \bmod M_\lambda} e^{2\pi i r \left( \frac{B}{2\pi} + \frac{1}{2r} \zeta \right) l} (-1)^{kl} \\ & \times \left( \frac{i}{2} \text{Res}_{z=0} \left\{ \frac{q_1(z+l) e^{2\pi i r \left( \frac{B}{2\pi} + \frac{1}{2r} \zeta \right) z}}{\sin^k(\pi z)} \cot(\pi r z) \right\} \right. \\ & \left. + \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq \left\lfloor \frac{B}{2\pi} + \frac{1}{2r} \zeta \right\rfloor}} \frac{\text{sign} \left( \frac{B}{2\pi} + \frac{1}{2r} \zeta \right) \text{Res}_{z=0} \left\{ \frac{q_1(z+l) e^{-2\pi i r \text{sign} \left( \frac{B}{2\pi} + \frac{1}{2r} \zeta \right) \left( m - \left\lfloor \frac{B}{2\pi} + \frac{1}{2r} \zeta \right\rfloor \right) z}}{\sin^k(\pi z)} \right\}}{\text{Sym}_\pm(m) \text{Sym}_\pm \left( m - \left\lfloor \frac{B}{2\pi} + \frac{1}{2r} \zeta \right\rfloor \right)} \right), \end{aligned}$$

where

$$A_j^J(r) = \pi i \sum_{(m, \lambda) \in J_j} \frac{g(\lambda)}{M_\lambda} \sum_{l \in \mathbb{Z} \bmod M_\lambda} (-1)^{kl} \text{Res}_{z=0} \left\{ \frac{q_1(z+l)}{\sin^k(\pi z)} \right\}$$

for  $j = +, -$ . If  $a'_0 - \pi k < 0$ , then  $Z(r) = Z_{\text{polar}}^J(r)$  for any  $J \subseteq I$ . In particular,  $A_j^J(r) = 0$ ,  $j = +, -$  in that case.

The disadvantage with the above expression for  $Z(r)$  is that the summation limit depends on  $r$  (except in the extreme case where  $q_1 = q$ , where  $\zeta(\lambda) = 0$  for all  $\lambda \in \Lambda$ ). Let us derive a slightly different expression valid for all large  $r$  avoiding this problem. Let  $\lambda \in \Lambda$ . If  $B(\lambda) \in \mathbb{R} \setminus 2\pi\mathbb{Z}$  or  $\zeta(\lambda) \geq 0$  we can choose a positive integer  $r_\lambda$  such that

$$\left\{ m \in \mathbb{Z} \mid m + \frac{B(\lambda)}{2\pi} + \frac{1}{2r} \zeta(\lambda) \geq 0 \right\} = \left\{ m \in \mathbb{Z} \mid m + \frac{B(\lambda)}{2\pi} \geq 0 \right\}$$

for all  $r \geq r_\lambda$ . If  $B(\lambda) \in 2\pi\mathbb{Z}$  and  $\zeta(\lambda) < 0$  we can choose  $r_\lambda$  such that

$$\left\{ m \in \mathbb{Z} \mid m + \frac{B(\lambda)}{2\pi} + \frac{1}{2r} \zeta(\lambda) \geq 0 \right\} = \left\{ m \in \mathbb{Z} \mid m + \frac{B(\lambda)}{2\pi} > 0 \right\}$$

for all  $r \geq r_\lambda$ . Put  $r_0 = \max\{r_\lambda\}_{\lambda \in \Lambda}$ . For  $r \geq r_0$  we have

$$\begin{aligned} Z_{\text{polar}}^I(\delta, \eta) &= A_1(\delta, \eta) + 2\pi i \sum_{l \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) e^{2\pi i r \left( \frac{B}{2\pi} + \frac{1}{2r} \zeta \right) l} (-1)^{kl} \\ &\quad \times \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} \geq 0}} \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \frac{q_1(z+l) e^{2\pi i r \nu(m, \lambda) z}}{\sin^k(\pi(z-i\eta))} \right\}, \end{aligned}$$

where

$$A_1(\delta, \eta) = -2\pi i \sum_{l \in \mathbb{Z}} \sum_{\substack{\lambda \in \Lambda \\ B \in 2\pi\mathbb{Z}; \zeta < 0}} g(\lambda) e^{\pi i \zeta l} (-1)^{kl} \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \frac{q_1(z+l) e^{\pi i \zeta z}}{\sin^k(\pi(z-i\eta))} \right\}$$

We therefore have

$$\begin{aligned} Z_{\text{polar}}^I(\delta, \eta) &= A_1(\delta, \eta) + A_2(\delta, \eta) + 2\pi i \sum_{l \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) e^{2\pi i r (\frac{B}{2\pi} + \frac{1}{2r}\zeta)l} (-1)^{kl} \\ &\quad \times \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} \geq 0}} \frac{1}{\text{Sym}_{\pm}(m + \frac{B}{2\pi})} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l)e^{2\pi i r \nu(m, \lambda)z}}{\sin^k(\pi(z-i\eta))} \right\}, \end{aligned}$$

where

$$A_2(\delta, \eta) = \pi i \sum_{l \in \mathbb{Z}} \sum_{\substack{\lambda \in \Lambda \\ B \in 2\pi\mathbb{Z}}} g(\lambda) e^{\pi i \zeta l} (-1)^{kl} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l)e^{\pi i \zeta z}}{\sin^k(\pi(z-i\eta))} \right\}.$$

We note that

$$A_1(\delta, \eta) + A_2(\delta, \eta) = A_+(\delta, \eta) + A_0(\delta, \eta) - A_-(\delta, \eta),$$

where

$$A_j(\delta, \eta) = \pi i \sum_{l \in \mathbb{Z}} \sum_{\lambda \in \Lambda_j} g(\lambda) e^{\pi i \zeta l} (-1)^{kl} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l)e^{\pi i \zeta z}}{\sin^k(\pi(z-i\eta))} \right\}$$

for  $j = +, 0, -$ , where  $\Lambda_j = \{\lambda \in \Lambda \mid B \in 2\pi\mathbb{Z}, j\zeta(\lambda) > 0\}$  for  $j = +, -$  and  $\Lambda_0 = \{\lambda \in \Lambda \mid B \in 2\pi\mathbb{Z}, \zeta(\lambda) = 0\}$ . Similarly, we find that

$$\begin{aligned} Z_{\text{polar}}^{\emptyset}(\delta, \eta) &= A_+(\delta, \eta) - A_0(\delta, \eta) - A_-(\delta, \eta) + 2\pi i \sum_{l \in \mathbb{Z}} \sum_{\lambda \in \Lambda} g(\lambda) e^{2\pi i r (\frac{B}{2\pi} + \frac{1}{2r}\zeta)l} (-1)^{kl} \\ &\quad \times \sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} \geq 0}} \frac{1}{\text{Sym}_{\pm}(m + \frac{B}{2\pi})} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l)e^{2\pi i r \nu(m, \lambda)z}}{\sin^k(\pi(z-i\eta))} \right\}. \end{aligned}$$

By letting  $I_+ = I$  and  $I_- = \emptyset$  we therefore have

**Theorem 6.3.** *There exists a family of positive integers  $\{M_\lambda\}_{\lambda \in \Lambda}$  such that*

$$\begin{aligned} Z_{\text{polar}}^{I_\tau}(r) &= r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}^{I_\tau}(\delta, \eta) \\ &= r (A_+ + \tau A_0 - A_-) + 2\pi i r \sum_{\lambda \in \Lambda} \frac{g(\lambda)}{M_\lambda} \sum_{l \in \mathbb{Z} \bmod M_\lambda} e^{2\pi i r (\frac{B}{2\pi} + \frac{1}{2r}\zeta)l} (-1)^{kl} \\ &\quad \times \left( \frac{i}{2} \text{Res}_{z=0} \left\{ \frac{q_1(z+l)e^{2\pi i r (\frac{B}{2\pi} + \frac{1}{2r}\zeta)z}}{\sin^k(\pi z)} \cot(\pi r z) \right\} \right. \\ &\quad \left. + \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq \lfloor \frac{B}{2\pi} \rfloor}} \frac{\text{sign}(B) \text{Res}_{z=0} \left\{ \frac{q_1(z+l)e^{2\pi i r (\text{sign}(B)m + \frac{B}{2\pi} + \frac{1}{2r}\zeta)z}}{\sin^k(\pi z)} \right\}}{\text{Sym}_{\pm}(m) \text{Sym}_{\pm}(m - \lfloor \frac{B}{2\pi} \rfloor)} \right) \end{aligned}$$

for all  $r \geq r_0$  and  $\tau = +, -$ , where  $A_j = \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} A_j(\delta, \eta)$  is given by

$$A_j = \pi i \sum_{\lambda \in \Lambda_j} \frac{g(\lambda)}{M_\lambda} \sum_{l \in \mathbb{Z} \bmod M_\lambda} e^{\pi i \zeta l} (-1)^{kl} \text{Res}_{z=0} \left\{ \frac{q_1(z+l)e^{\pi i \zeta z}}{\sin^k(\pi z)} \right\}$$

for  $j = +, 0, -$ . If  $a'_0 - k\pi < 0$ , then  $Z(r) = Z_{\text{polar}}^{I_\tau}(r)$ ,  $\tau = +, -$ . In particular,  $A_0 = 0$  in this case.

If  $q_1 = q$  (so  $\zeta(\lambda) = 0$  for all  $\lambda \in \Lambda$ ) and  $a_0 - k\pi < 0$ , then  $A_0 = 0$  and  $A_- = A_+ = 0$  since  $\Lambda_j = \emptyset$ ,  $j = -, +$ . Note that Theorem 6.3 is a corollary of Theorem 6.2 in this case. In fact  $(m, \lambda) \mapsto \lambda$  defines a bijection  $I \rightarrow \Lambda_0$  and  $A_0 = A_+^I(r) = A_-^\emptyset(r)$  (which is independent of  $r$  in this case).

Let us finally take a closer look at the Seifert case, i.e. the case  $Z(r) = Z(X; r)$ . Then  $q(z) = \exp\left(-i\pi\left(\sum_{j=1}^n \frac{\mu_j}{\alpha_j}\right)z\right)$  and  $B = -2\pi \sum_{j=1}^n \frac{n_j}{\alpha_j}$ . By using Theorem 6.2 and Theorem 6.3 directly we would end up with a result similar to Corollary 5.10. As argued below that corollary we obtain in this way a sum with pairs of terms with the same Chern–Simons value. We will therefore follow a strategy similar to the one followed after Corollary 5.10.

Recall that we always assume that  $a_\epsilon g$  is even. We will further in what follows assume that  $n + a_\epsilon g - 2 > 0$ . This is not a serious assumption. Since we treat the case  $n = 0$  as the case  $n = 1$  with  $(\alpha_1, \beta_1) = (1, 0)$  we have  $n + a_\epsilon g - 2 > 0$  except in case  $g = 0$  (hence  $\epsilon = o$ ) and  $n = 1, 2$ . Since we further assume that  $E = 0$ , we find that  $X = S^2 \times S^1$  in these cases (compare with [JN, Sect. 4 pp. 28–31] and the discussion following Theorem 4.4).

Let us first put  $q_0(z) = q(z)$ , so  $\zeta = \zeta(\underline{\mu}) = -\sum_{j=1}^n \frac{\mu_j}{\alpha_j}$ . Since  $q_1 = 1$  (so  $a'_0 = 0$ ) we get from the general argument that

$$Z(X; r) = r \lim_{\delta \rightarrow 0+} \sqrt{\delta} \lim_{\eta \rightarrow 0+} Z_{\text{polar}}^J(\delta, \eta),$$

where

$$\begin{aligned} Z_{\text{polar}}^J(\delta, \eta) &= 2\pi i \sum_{l \in \mathbb{Z}} \sum_{\underline{\mu} \in \{\pm 1\}^n} \sum_{\underline{n} \in \mathcal{S}} g(\underline{\mu}, \underline{n}) \\ &\times \sum_{\substack{m \in \mathbb{Z} \\ \text{sign}(\nu(m, \underline{\mu}, \underline{n}))=1}} \text{Res}_{z=l+i\eta} \left\{ e^{-\pi\delta z^2} \frac{e^{2\pi i r \nu(m, \underline{\mu}, \underline{n})z}}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))} \right\}, \end{aligned}$$

where  $\nu(m, \underline{\mu}, \underline{n}) = m - \sum_{j=1}^n \frac{n_j}{\alpha_j} - \frac{1}{2r} \sum_{j=1}^n \frac{\mu_j}{\alpha_j}$  and  $g(\underline{\mu}, \underline{n})$  is given by (79). Here  $\text{sign}(\nu(m, \underline{\mu}, \underline{n})) = 1$  if  $(m, \underline{\mu}, \underline{n}) \in J$  and  $\text{sign}(\nu(m, \underline{\mu}, \underline{n})) = -1$  if  $(m, \underline{\mu}, \underline{n}) \in I \setminus J$  (so here the terms with  $\nu(m, \underline{\mu}, \underline{n}) = 0$  do not contribute to  $Z(X; r)$ ). Let us specialize to the case  $J = I$  and let us write  $Z_{\text{polar}}^I$  for  $Z_{\text{polar}}^J$ . Like in the  $E \neq 0$  case we first rewrite the above expression by introducing  $n'_j = n_j + \frac{1}{2}\beta_j l$ . This leads to

$$\begin{aligned} Z_{\text{polar}}(\delta, \eta) &= A_+^I(\delta, \eta) + 2\pi i \left(\frac{i}{2}\right)^n \sum_{l \in \mathbb{Z}} (-1)^{nl} \sum_{\underline{n}' \in J_l} \exp(2\pi i r q(l, \underline{n}')) \\ &\times \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} n'_j - \frac{1}{2} \sigma_j l \right) \right] \\ &\times \sum_{\substack{m \in \mathbb{Z} \\ \nu(m, \underline{\mu}, \underline{n}') \geq 0}} \frac{1}{\text{Sym}_\pm(\nu(m, \underline{\mu}, \underline{n}'))} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{e^{2\pi i r \nu(m, \underline{\mu}, \underline{n}')z}}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))} \right\}, \end{aligned}$$

where  $J_l$  is given by (115) and

$$A_+^I(\delta, \eta) = \pi i \left(\frac{i}{2}\right)^n \sum_{l \in \mathbb{Z}} (-1)^{nl} \sum_{\underline{n}' \in J_l} \exp(2\pi i r q(l, \underline{n}'))$$

$$\begin{aligned}
& \times \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} n'_j - \frac{1}{2} \sigma_j l \right) \right] \\
& \times \sum_{\substack{m \in \mathbb{Z} \\ \nu(m, \underline{\mu}, \underline{n}') = 0}} \operatorname{Res}_{z=i\eta} \left\{ e^{-\pi \delta(z+l)^2} \frac{1}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))} \right\}.
\end{aligned}$$

Note that  $\sum_{j=1}^n \frac{n'_j}{\alpha_j} = \sum_{j=1}^n \frac{n_j}{\alpha_j}$  since  $E = 0$ . Like in the case  $E \neq 0$  we can change the sum  $\sum_{\underline{n}' \in J_l}$  to

$$\sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\underline{n}' \in J'_l} \frac{1}{\prod_{j=1}^n \operatorname{Sym}_{\mathbb{Z}_\pm} \left( \frac{n'_j}{\alpha_j} \right)}$$

if we at the same time substitute  $\mu'_j n'_j$  for  $n'_j$  everywhere. This leads to the identity

$$\begin{aligned}
Z_{\text{polar}}(\delta, \eta) &= A_+^I(\delta, \eta) + 2\pi i \left( \frac{i}{2} \right)^n \sum_{l \in \mathbb{Z}} (-1)^{nl} \sum_{\underline{n}' \in J'_l} \frac{1}{\prod_{j=1}^n \operatorname{Sym}_{\mathbb{Z}_\pm} \left( \frac{n'_j}{\alpha_j} \right)} \exp(2\pi i r q(l, \underline{n}')) \\
&\times \sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\substack{m \in \mathbb{Z} \\ \nu(m, \underline{\mu}, \underline{n}', \underline{\mu}') \geq 0}} \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \frac{1}{\operatorname{Sym}_\pm(\nu(m, \underline{\mu}, \underline{n}', \underline{\mu}'))} \\
&\times \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \\
&\times \operatorname{Res}_{z=i\eta} \left\{ e^{-\pi \delta(z+l)^2} \frac{e^{2\pi i r \nu(m, \underline{\mu}, \underline{n}', \underline{\mu}') z}}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))} \right\},
\end{aligned}$$

where  $J'_l$  is given by (116) and  $\nu(m, \underline{\mu}, \underline{n}', \underline{\mu}') = m - \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} - \frac{1}{2r} \sum_{j=1}^n \frac{\mu_j}{\alpha_j}$ . By an application of Lemma 5.7 we thus find that

$$\begin{aligned}
Z_{\text{polar}}(\delta, \eta) &= A_+^I(\delta, \eta) + 2\pi i \left( \frac{i}{2} \right)^n \sum_{l \in \mathbb{Z}} (-1)^{nl} \sum_{\underline{n}' \in J'_l} \frac{1}{\prod_{j=1}^n \operatorname{Sym}_{\mathbb{Z}_\pm} \left( \frac{n'_j}{\alpha_j} \right)} \\
&\times \exp(2\pi i r q(l, \underline{n}')) \left( Z_0^{(l, \underline{n}')}(\delta, \eta) + \tilde{Z}_1^{(l, \underline{n}')}(\delta, \eta) \right),
\end{aligned}$$

where  $Z_0^{(l, \underline{n}')}(\delta, \eta)$  is given by (118) (with  $E = 0$ ) and

$$\begin{aligned}
\tilde{Z}_1^{(l, \underline{n}')}(\delta, \eta) &= - \sum_{\underline{\mu}' \in \{\pm 1\}^n} \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \\
&\times \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |a|}} \frac{\operatorname{sign}(a)}{\operatorname{Sym}_\pm(m) \operatorname{Sym}_\pm(m - |a|)} \\
&\times \operatorname{Res}_{z=i\eta} \left\{ e^{-\pi \delta(z+l)^2} \frac{\exp(2\pi i r \operatorname{sign}(a)(m - |a|)z)}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))} \right\},
\end{aligned}$$



where  $a = a(\underline{\mu}, \underline{\mu}', \underline{n}') = \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} + \frac{1}{2r} \sum_{j=1}^n \frac{\mu_j}{\alpha_j}$ . Since the terms with  $\nu(m, \underline{\mu}, \underline{n}) = 0$  do not contribute to  $Z(X; r)$  we have  $\lim_{\delta \rightarrow 0+} \sqrt{\delta} \lim_{\eta \rightarrow 0+} A_+^I(\delta, \eta) = 0$ . The first part of Theorem 4.4 thus follows by calculating  $\lim_{\delta \rightarrow 0+} \sqrt{\delta} \lim_{\eta \rightarrow 0+} Z_{\text{polar}}(\delta, \eta)$  like in the case  $E \neq 0$ .

Like Theorem 6.2 the disadvantage with the above obtained expression for  $Z(X; r)$  is that the summation limit in  $\tilde{Z}_1^{(l, \underline{n}')} (r)$ , see Theorem 4.4, depends on the level  $r$ . Let us proceed like in the proof of Theorem 6.3 to obtain a slightly different expression for  $Z(X; r)$ . In fact, there exists a positive integer  $r_0$  (depending on  $X$ ) such that

$$\begin{aligned} Z_{\text{polar}}^{I_\tau}(\delta, \eta) &= A_+(\delta, \eta) + \tau A_0(\delta, \eta) - A_-(\delta, \eta) \\ &+ 2\pi i \left(\frac{i}{2}\right)^n \sum_{l \in \mathbb{Z}} (-1)^{nl} \sum_{\underline{n}' \in J_l'} \frac{1}{\prod_{j=1}^n \text{Sym}_{\mathbb{Z}_{\pm}} \left(\frac{n'_j}{\alpha_j}\right)} \exp(2\pi i r q(l, \underline{n}')) \sum_{\underline{\mu}' \in \{\pm 1\}^n} \\ &\times \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \sum_{\substack{m \in \mathbb{Z} \\ m \geq \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j}}} \frac{1}{\text{Sym}_{\pm} \left(m - \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j}\right)} \\ &\times \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \\ &\times \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \frac{e^{2\pi i r \nu(m, \underline{\mu}, \underline{n}', \underline{\mu}') z}}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))} \right\} \end{aligned}$$

for all  $r \geq r_0$ , where  $\nu(m, \underline{\mu}, \underline{n}', \underline{\mu}') = m - \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} - \frac{1}{2r} \sum_{j=1}^n \frac{\mu_j}{\alpha_j}$  as above and

$$\begin{aligned} A_j(\delta, \eta) &= \pi i \left(\frac{i}{2}\right)^n \sum_{l \in \mathbb{Z}} (-1)^{nl} \sum_{\underline{n}' \in J_l'} \frac{1}{\prod_{j=1}^n \text{Sym}_{\mathbb{Z}_{\pm}} \left(\frac{n'_j}{\alpha_j}\right)} \exp(2\pi i r q(l, \underline{n}')) \\ &\times \sum_{(\underline{\mu}, \underline{\mu}') \in \Lambda(\underline{n}', j)} \left( \prod_{j=1}^n \mu_j \right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \\ &\times \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \frac{\exp \left[ -\pi i \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right]}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))} \right\} \end{aligned}$$

for  $j = +, 0, -$ , where

$$\Lambda(\underline{n}', j) = \left\{ (\underline{\mu}, \underline{\mu}') \in \{\pm 1\}^n \times \{\pm 1\}^n \mid -\text{sign} \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) = j, \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \in \mathbb{Z} \right\},$$

where  $\text{sign}(0) = 0$ . (Note the minus in front of  $\text{sign} \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right)$  due to the fact that  $\zeta = \zeta(\underline{\mu}) = -\sum_{j=1}^n \frac{\mu_j}{\alpha_j}$ .) By Lemma 5.7 we then find for  $r \geq r_0$  that

$$\begin{aligned} Z_{\text{polar}}^{I_\tau}(\delta, \eta) &= \pi i \left(\frac{i}{2}\right)^n \sum_{l \in \mathbb{Z}} (-1)^{nl} \sum_{\underline{n}' \in J_l'} \frac{1}{\prod_{j=1}^n \text{Sym}_{\mathbb{Z}_{\pm}} \left(\frac{n'_j}{\alpha_j}\right)} \exp(2\pi i r q(l, \underline{n}')) \left( 2Z_0^{(l, \underline{n}')}(\delta, \eta) \right. \\ &\quad \left. + 2Z_1^{(l, \underline{n}')}(\delta, \eta) + A_+^{(l, \underline{n}')}(\delta, \eta) + \tau A_0^{(l, \underline{n}')}(\delta, \eta) - A_-^{(l, \underline{n}')}(\delta, \eta) \right), \end{aligned}$$

where  $Z_0^{(l, \underline{n}')}(\delta, \eta)$  and  $Z_1^{(l, \underline{n}')}(\delta, \eta)$  are given as in the case  $E \neq 0$ , i.e. they are given by respectively (118) and (119) with  $E = 0$ , and where

$$A_j^{(l, \underline{n}')}(\delta, \eta) = \sum_{(\underline{\mu}, \underline{\mu}') \in \Lambda(\underline{n}', j)} \left( \prod_{j=1}^n \mu_j \right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} \mu'_j n'_j - \frac{1}{2} \sigma_j l \right) \right] \\ \times \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \frac{\exp \left[ -\pi i \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right]}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))} \right\}$$

for  $j = +, 0, -$ . This expression leads to

$$Z(X; r) = Z_{\text{polar}}(X; r) + r Z_{\text{spec}}(X; r) \quad (130)$$

for  $r \geq r_0$ , where  $Z_{\text{polar}}(X; r)$  is as in Theorem 4.1 (with  $E = 0$ ) and

$$Z_{\text{spec}}(X; r) = \sum_{(l, \underline{n}') \in \mathcal{I}_2^a} b_{(l, \underline{n}')} \exp(2\pi i r q(l, \underline{n}')) \left( A_+^{(l, \underline{n}')} - A_-^{(l, \underline{n}')} \right),$$

where  $\mathcal{I}_2$  and  $b_{(l, \underline{n}')}$  are as in Theorem 4.1 and

$$A_j^{(l, \underline{n}')} = \frac{\pi i}{2} \left( \frac{i}{4} \right)^n \sum_{(\underline{\mu}, \underline{\mu}') \in \Lambda(\underline{n}', j)} \left( \prod_{j=1}^n \mu_j \right) \exp \left( 2\pi i \sum_{j=1}^n \mu_j \frac{\rho_j}{\alpha_j} \mu'_j n'_j \right) \\ \times \text{Res}_{z=0} \left\{ \frac{\exp \left[ -\pi i \left( \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \right) z \right]}{\sin^{n+a_\epsilon g-2}(\pi z)} \right\}$$

for  $j = +, -$ . (Note that we necessarily have  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} A_0(\delta, \eta) = 0$ .)

Now put  $q_1(z) = q(z)$  (so  $q_0(z) = 1$ ,  $\zeta = 0$ ). In this case (80) is valid with  $a_0 = \pi \sum_{j=1}^n \frac{1}{\alpha_j}$  (and  $b_0 = 1$ ). Assume in the following that  $a_0 < \pi(n + a_\epsilon g - 2)$ . Then we have from the general case, see Theorem 6.2. that

$$Z(X; r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \lim_{\eta \rightarrow 0_+} Z_{\text{polar}}^J(\delta, \eta),$$

where

$$Z_{\text{polar}}^J(\delta, \eta) = 2\pi i \left( \frac{i}{2} \right)^n \sum_{l \in \mathbb{Z}} (-1)^{nl} \sum_{\underline{n}' \in J_l} \exp(2\pi i r q(l, \underline{n}')) \\ \times \sum_{\underline{\mu} \in \{\pm 1\}^n} \left( \prod_{j=1}^n \mu_j \right) \exp \left[ 2\pi i \sum_{j=1}^n \mu_j \left( \frac{\rho_j}{\alpha_j} n'_j - \frac{1}{2} \sigma_j l \right) \right] \\ \times \sum_{\substack{m \in \mathbb{Z} \\ \text{sign}(\nu(m, \underline{n}'))=1}} \text{Res}_{z=i\eta} \left\{ e^{-\pi \delta (z+l)^2} \frac{q(z) e^{2\pi i r \nu(m, \underline{n}') z}}{\sin^{n+a_\epsilon g-2}(\pi(z-i\eta))} \right\},$$

where  $\nu(m, \underline{n}') = m - \sum_{j=1}^n \frac{n'_j}{\alpha_j}$  and  $\text{sign}(\nu(m, \underline{n}')) = 1$  if  $(m, \underline{n}') \in J$  and  $\text{sign}(\nu(m, \underline{n}')) = -1$  if  $(m, \underline{n}') \in I \setminus J$ , where  $J$  is any subset of  $I$ . By the same procedure as used above we immediately get that  $Z(X; r) = Z_{\text{polar}}(X; r)$ , where  $Z_{\text{polar}}(X; r)$  is as in Theorem 4.1 (with  $E = 0$ ). A comparison with (130) reveals that  $Z_{\text{spec}}(X; r) = 0$  for  $r \geq r_0$  in case  $a_0 < \pi(n + a_\epsilon g - 2)$ , and since the  $q(l, \underline{n}')$ 's are rational, we in fact have that  $Z_{\text{spec}}(X; r) = 0$  for all levels  $r$  if  $a_0 < \pi(n + a_\epsilon g - 2)$ .

Let  $(l, \underline{n}') \in \mathcal{I}_2$  be fixed. If  $\underline{\mu}' \in \{\pm 1\}^n$  satisfies that  $\sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \in \mathbb{Z}$ , then the same is true for  $-\underline{\mu}'$ . Moreover, if  $\underline{\mu} \in \{\pm 1\}^n$  satisfies that  $\sum_{j=1}^n \frac{\mu_j}{\alpha_j} < 0$ , then  $-\underline{\mu}$  satisfies the opposite inequality. These observations show that  $Z_{\text{spec}}(X; r)$  is given by the formula in Theorem 4.4.

The level  $r_0$  is chosen so that we for all  $r \geq r_0$ ,  $l \in \mathbb{Z}$  and  $(\underline{\mu}, \underline{\mu}', \underline{n}') \in \{\pm 1\}^n \times \{\pm 1\}^n \times J'_l$  have

$$\left\{ m \in \mathbb{Z} \left| m - \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} - \frac{1}{2r} \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \geq 0 \right. \right\} = \left\{ m \in \mathbb{Z} \left| m - \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} \geq 0 \right. \right\}$$

in case  $\sum_{j=1}^n \frac{\mu_j}{\alpha_j} \notin \mathbb{Z}$  or  $\sum_{j=1}^n \frac{\mu_j}{\alpha_j} \leq 0$ , and

$$\left\{ m \in \mathbb{Z} \left| m - \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} - \frac{1}{2r} \sum_{j=1}^n \frac{\mu_j}{\alpha_j} \geq 0 \right. \right\} = \left\{ m \in \mathbb{Z} \left| m - \sum_{j=1}^n \frac{\mu'_j n'_j}{\alpha_j} > 0 \right. \right\}$$

if  $\sum_{j=1}^n \frac{\mu_j}{\alpha_j} \in \mathbb{Z}$  and  $\sum_{j=1}^n \frac{\mu_j}{\alpha_j} > 0$ . In particular we can put  $r_0 = 2$  (the minimum level) for  $n = 1$ . In general we see that  $r_0$  only depends on the pairs of Seifert invariants  $(\alpha_1, \beta_1), \dots, (\alpha_n, \beta_n)$  (so is independent of the data for the base space, i.e. the genus  $g$  and the invariant  $\epsilon$ ).

## 7. APPENDICES

We have in the following appendices collected material of a technical nature.

**7.1. Appendix A. The periodicity lemma.** The following lemma is proven in bigger generality than needed in this paper. However, to make parts of the ideas in this paper useable to other situations, in particular to multi-dimensional cases, we give it in this generality. One can find an even more general version of the periodicity lemma in [HT, Appendix], where it is used in a multi-dimensional setting.

**Lemma 7.1.** *Let  $F : \mathbb{Z}^n \rightarrow \mathbb{C}$  be a function periodic in all variables with a period of  $N_i$  in the  $i$ th variabel. Let  $P_i(x) = A_i x^2 + B_i x + C_i$  be polynomials with complex coefficients such that  $\text{Re}(A_i) > 0$  for all  $i$ . Then*

$$\begin{aligned} & \sum_{k_1=0}^{N_1-1} \dots \sum_{k_n=0}^{N_n-1} F(k_1, \dots, k_n) \\ &= \left( \prod_{i=1}^n \sqrt{A_i N_i} \right) \lim_{\varepsilon \rightarrow 0_+} \varepsilon^{n/2} \sum_{(k_1, \dots, k_n) \in \mathbb{Z}^n} e^{-\pi \varepsilon f(\varepsilon)(P_1(k_1) + \dots + P_n(k_n))} F(k_1, \dots, k_n) \end{aligned}$$

for all functions  $f : ]0, \varepsilon_0] \rightarrow ]0, \infty[$  with  $\lim_{\varepsilon \rightarrow 0_+} f(\varepsilon) = 1$ , where  $\varepsilon_0 > 0$  is arbitrary but fixed.

**Proof.** The series  $\sum_{(k_1, \dots, k_n) \in \mathbb{Z}^n} e^{-\pi \varepsilon f(\varepsilon)(P_1(k_1) + \dots + P_n(k_n))} F(k_1, \dots, k_n)$  is absolutely convergent by periodicity of  $F$ , hence

$$\begin{aligned} & \sum_{(k_1, \dots, k_n) \in \mathbb{Z}^n} e^{-\pi \varepsilon f(\varepsilon)(P_1(k_1) + \dots + P_n(k_n))} F(k_1, \dots, k_n) \\ &= \sum_{j_1=0}^{N_1-1} \dots \sum_{j_n=0}^{N_n-1} F(j_1, \dots, j_n) \sum_{(k_1, \dots, k_n) \in \mathbb{Z}^n} e^{-\pi \varepsilon f(\varepsilon)(P_1(j_1 + k_1 N_1) + \dots + P_n(j_n + k_n N_n))}. \end{aligned}$$

Thus we have to show that

$$\lim_{\varepsilon \rightarrow 0_+} \sqrt{\varepsilon} \sqrt{AN} Q(\varepsilon) = 1, \quad (131)$$

where  $Q(\varepsilon) = \sum_{k \in \mathbb{Z}} e^{-\pi \varepsilon f(\varepsilon) P(j+kN)}$  for a polynomial  $P(x) = Ax^2 + Bx + C$  with  $\operatorname{Re}(A) > 0$ . Let  $j \in \{0, 1, \dots, N-1\}$  be fixed. By Poisson's formula (72) we have

$$Q(\varepsilon) = \sum_{m \in \mathbb{Z}} \int_{-\infty}^{\infty} e^{2\pi i m x} e^{-\pi \varepsilon f(\varepsilon) P(j+xN)} dx$$

which is a sum of Gaussian integrals. Thus

$$Q(\varepsilon) = \frac{e^{-\pi k(\varepsilon)C}}{N \sqrt{Ak(\varepsilon)}} \exp\left(\frac{\pi}{4} \frac{B^2 k(\varepsilon)}{A}\right) \sum_{m \in \mathbb{Z}} e^{-\pi i \frac{2jA+B}{AN} m} \exp\left(-\frac{\pi}{Ak(\varepsilon)N^2} m^2\right),$$

where  $k(\varepsilon) = \varepsilon f(\varepsilon)$  from which it follows that  $Q(\varepsilon)$  is absolutely convergent. The term coming from  $m = 0$  gives the contribution

$$\lim_{\varepsilon \rightarrow 0_+} \frac{e^{-\pi k(\varepsilon)C}}{\sqrt{f(\varepsilon)}} \exp\left(\frac{\pi}{4} \frac{B^2 k(\varepsilon)}{A}\right) = 1$$

to the left-hand side of (131). To see that the remaining part is zero we simply note that

$$\begin{aligned} & \left| \sum_{m \in \mathbb{Z} \setminus \{0\}} e^{-\pi i \frac{2jA+B}{AN} m} \exp\left(-\frac{\pi}{Ak(\varepsilon)N^2} m^2\right) \right| \\ & \leq \exp\left(-\frac{1}{2} \frac{\pi}{Ak(\varepsilon)N^2}\right) \sum_{m \in \mathbb{Z} \setminus \{0\}} \exp\left(-\frac{1}{2} \frac{\pi}{Ak(\varepsilon)N^2} m^2 + \frac{\pi \operatorname{Im}(B)}{AN} m\right). \end{aligned}$$

where the factor in front of the  $m$ -sum in the right-hand side converges to zero as  $\varepsilon \rightarrow 0_+$  and the  $m$ -sum in the right-hand side is majorized by the convergent and  $\varepsilon$ -independent series  $\sum_{m \in \mathbb{Z} \setminus \{0\}} \exp\left(-\frac{1}{2} \frac{\pi}{2A\varepsilon_0 N^2} m^2 + \frac{\pi \operatorname{Im}(B)}{AN} m\right)$  on an interval on the form  $]0, \varepsilon_1]$ , where  $\varepsilon_1 \leq \varepsilon_0$  is chosen such that  $f(\varepsilon) \leq 2$  for all  $\varepsilon \in ]0, \varepsilon_1]$ .  $\square$

**7.2. Appendix B. Proofs of Lemma 5.4–Lemma 5.6.** For the following proofs we need certain estimates. Let  $a_\eta(t; m, \lambda) = |\sin(\pi(\gamma_{(m, \lambda)}(t) - i\eta))|$ , where  $\gamma_{(m, \lambda)}$  is the steepest descent contour, see (90). For  $u, v \in \mathbb{R}$  we have

$$|\sin(u + iv)| = \frac{1}{2} \sqrt{e^{2v} + e^{-2v} - 2 \cos(2u)} \quad (132)$$

and therefore

$$a_\eta(t; m, \lambda) = \frac{1}{2} \sqrt{e^{2\pi(\operatorname{sign}(A) \frac{t}{\sqrt{2}} - \eta)} + e^{-2\pi(\operatorname{sign}(A) \frac{t}{\sqrt{2}} - \eta)} - 2 \cos\left(2\pi \left[z_{\text{st}}(m, \lambda) + \frac{t}{\sqrt{2}}\right]\right)}. \quad (133)$$

Let  $\eta_0 > 0$  be as in (88). Then  $z_{\text{st}}(m, \lambda) + \operatorname{sign}(A)\eta \in \mathbb{R} \setminus \mathbb{Z}$  for all  $(m, \lambda) \in \mathbb{Z} \times \Lambda$  and all  $\eta \in ]0, \eta_0]$ . Since the function  $x \mapsto e^x + e^{-x}$  is strictly decreasing on  $] -\infty, 0]$  and strictly increasing on  $[0, \infty]$  and takes the value 2 for  $x = 0$  we thus have

$$d_\eta := \inf_{t \in \mathbb{R}, m \in \mathbb{Z}, \lambda \in \Lambda} |a_\eta(t; m, \lambda)| > 0 \quad (134)$$

for every fixed  $\eta \in ]0, \eta_0]$ . Moreover, letting  $\rho = \sqrt{2}\eta_0$ , there exists a  $\alpha \in [0, 1[$  such that  $\cos\left(2\pi \left[z_{\text{st}}(m, \lambda) + t/\sqrt{2}\right]\right) \leq \alpha$  for all  $t \in [-\rho, \rho]$  and all  $(m, \lambda) \in W$ . (We use here that  $\{z_{\text{st}}(m, \lambda) \pmod{\mathbb{Z}} \mid (m, \lambda) \in \mathbb{Z} \times \Lambda\}$  is finite. If we only assume (88) then put  $\rho = \eta_0$ .) Let

$s = -\text{sign}(A)\frac{t}{\sqrt{2}}$ . Then  $a_\eta(t; m, \lambda) \geq \frac{1}{2}\sqrt{e^{2\pi(s+\eta)} + e^{-2\pi(s+\eta)} - 2}$ . Letting  $\eta_1 = \frac{\rho}{2\sqrt{2}} < \eta_0$  we then get  $a_\eta(t; m, \lambda) \geq \frac{1}{2}\sqrt{e^{\pi\eta_1} + e^{-\pi\eta_1} - 2}$  for  $|t| > \rho$  and  $\eta \in [0, \eta_1]$ . We conclude that

$$a_\eta(t; m, \lambda) \geq M := \min \left\{ \frac{1}{\sqrt{2}}\sqrt{1 - \alpha}, \frac{1}{2}\sqrt{e^{\pi\eta_1} + e^{-\pi\eta_1} - 2} \right\} > 0 \quad (135)$$

for all  $t \in \mathbb{R}$ ,  $\eta \in [0, \eta_1]$  and  $(m, \lambda) \in W$ .

**Proof of Lemma 5.4** Let  $(m, \lambda) \in \mathbb{Z} \times \Lambda$  be fixed in the following and let  $C_{\text{sd}} = C_{\text{sd}}(m, \lambda)$ ,  $\gamma = \gamma_{(m, \lambda)}$ ,  $z_{\text{st}} = z_{\text{st}}(m, \lambda)$ , and  $f(z) = f(z; m, \lambda)$ . Assume first that  $k > 0$ . Observe that  $\text{sign}(A)(\text{Re}z_l - z_{\text{st}}) > \text{Im}z_l$  is equivalent to  $\text{sign}(A)(l - z_{\text{st}}) > \eta$ , explaining the summation range in the sum of residues in (95). We will first show that the integral  $\int_{C_{\text{sd}}} f(z)dz$  is convergent for all sufficiently small positive  $\eta$ . By (77), (92) and (134) we have

$$|f(\gamma(t))| \leq \frac{1}{d_\eta^k} |e^{-\pi\delta\gamma(t)^2} q(\gamma(t))| \exp(-r|A|t^2)$$

for every  $\delta > 0$  and  $\eta \in ]0, \eta_0]$ . Here  $|e^{-\pi\delta\gamma(t)^2}| = e^{-\pi\delta z_{\text{st}}^2} e^{-\pi\sqrt{2}z_{\text{st}}\delta t}$ , so by (80) we get

$$|f(\gamma(t))| \leq \frac{b_0}{d_\eta^k} e^{-\pi\delta z_{\text{st}}^2} e^{-\pi\sqrt{2}z_{\text{st}}\delta t} e^{a_0(|z_{\text{st}}|+|t|)} \exp(-r|A|t^2)$$

where the right-hand side is integrable. Thus  $\int_{C_{\text{sd}}} f(z)dz$  is convergent and

$$\begin{aligned} \left| \int_{C_{\text{sd}}} f(z)dz \right| &\leq \frac{b_0}{d_\eta^k} e^{-\pi\delta z_{\text{st}}^2} e^{a_0|z_{\text{st}}|} \int_{-\infty}^{\infty} e^{-\pi\sqrt{2}z_{\text{st}}\delta t} (e^{a_0t} + e^{-a_0t}) \exp(-r|A|t^2) dt \\ &= \frac{b_0}{d_\eta^k} e^{-\pi\delta z_{\text{st}}^2} e^{a_0|z_{\text{st}}|} \sqrt{\frac{\pi}{r|A|}} \left( \exp\left(\frac{a_+^2}{4r|A|}\right) + \exp\left(\frac{a_-^2}{4r|A|}\right) \right) \end{aligned} \quad (136)$$

where  $a_\pm = \pi\delta\sqrt{2}z_{\text{st}} \pm a_0$ .

Next we estimate the contributions coming from integrals along curves connecting  $C_{\text{sd}}$  by the real axes. Let  $R_n = n + \frac{1}{2}$  for  $n \in \mathbb{Z}$  and let  $\tau_n(t) = R_n + it$ ,  $t \in [a, b]$ , where  $a, b \in \mathbb{R}$ . Then

$$\begin{aligned} \left| \int_{\tau_n} f(z)dz \right| &\leq e^{-\pi\delta R_n^2} \int_a^b \frac{|q(\tau_n(t))|}{|\sin^k(\pi(\gamma(t) - i\eta))|} e^{\pi\delta t^2} |e^{rQ_m(\tau_n(t))}| dt \\ &\leq b_0 e^{a_0|R_n|} e^{-\pi\delta R_n^2} \int_a^b \frac{e^{a_0|t|}}{|\sin^k(\pi(\tau_n(t) - i\eta))|} e^{\pi\delta t^2} e^{2rA(z_{\text{st}}-R_n)t} dt \end{aligned}$$

by (80). By (132) we have  $|\sin(\pi(\tau_n(t) - i\eta))| \geq \frac{1}{\sqrt{2}}\sqrt{1 - \cos(2\pi R_n)} = 1$  so

$$\left| \int_{\tau_n} f(z)dz \right| \leq b_0 e^{a_0|R_n|} e^{-\pi\delta R_n^2} \int_a^b (e^{a_0t} + e^{-a_0t}) e^{\pi\delta t^2} e^{2rA(z_{\text{st}}-R_n)t} dt.$$

In our case  $\{a, b\} = \{0, \text{sign}(A)(R_n - z_{\text{st}})\}$ , where we assume that  $a \leq b$ . For  $c \in \mathbb{R}$  we have

$$\begin{aligned} &\int_a^b e^{ct} e^{\pi\delta t^2} e^{2rA(z_{\text{st}}-R_n)t} dt \\ &= \int_a^b e^{\pi\delta t^2} e^{-\text{sign}(A)[r|A|(R_n - z_{\text{st}}) - \text{sign}(A)c]t} e^{-\text{sign}(A)r|A|(R_n - z_{\text{st}})t} dt. \end{aligned}$$

We can choose a  $N_c \in \mathbb{N}$  such that

$$\text{sign}(A)(r|A|(R_n - z_{\text{st}}) - \text{sign}(A)c)t \geq 0$$

for all  $t \in [a, b]$  and all  $n \in \mathbb{Z}$  with  $|n| > N_c$ . Therefore

$$\left| \int_{\tau_n} f(z) dz \right| \leq 2b_0 e^{a_0 |R_n|} e^{-\pi \delta R_n^2} \int_a^b e^{-[r|A| \operatorname{sign}(A)(R_n - z_{\text{st}}) - \pi \delta t]t} dt$$

for all  $n \in \mathbb{Z}$  with  $|n| > N = \max\{N_{-a_0}, N_{a_0}\}$ . For  $|n| > N$  and  $\delta > 0$  with  $r|A| \geq \pi \delta$  we get

$$\begin{aligned} \left| \int_{\tau_n} f(z) dz \right| &\leq 2b_0 e^{a_0 |R_n|} e^{-\pi \delta R_n^2} \int_a^b e^{-r|A|[\operatorname{sign}(A)(R_n - z_{\text{st}}) - t]t} dt \\ &\leq 2b_0 e^{a_0 |R_n|} e^{-\pi \delta R_n^2} (b - a) = 2|R_n - z_{\text{st}}| b_0 e^{a_0 |R_n|} e^{-\pi \delta R_n^2} \end{aligned}$$

which converges to zero as  $|n| \rightarrow \infty$ . Now, since  $\int_{-\infty}^{\infty} f(z) dz$  is convergent for all  $\delta > 0$  and all  $\eta \in ]0, \eta_0]$ , it follows by the residue theorem and the above that

$$\sum_{\substack{l \in \mathbb{Z} \\ \operatorname{sign}(A)(l - z_{\text{st}}) > \eta}} \operatorname{Res}_{z=z_l(\eta)} \{f(z)\}$$

is convergent and (95) is satisfied for all  $\eta \in ]0, \eta_0]$  and all  $\delta \in ]0, r|A|/\pi]$ . We will actually see in what follows that this last sum is absolutely convergent.

The case  $k \leq 0$  is handled as above but is easier since the sin-factor is included in the function  $q$  and  $\eta$  is not present. Note also that in this case we do not need the assumption (88).  $\square$

For the next proof and later we note the following elementary fact which is stated for the sake of easy reference. Let  $a > 0$ ,  $\delta > 0$  and  $b \in \mathbb{C}$  be constants and let  $j \geq 1$ . Then

$$\left( \frac{d}{dz} \right)^j e^{-a\delta(z+b)^2} = P_j(z+b, \delta) e^{-a\delta(z+b)^2}, \quad (137)$$

where  $P_j(y, \delta)$  is a polynomial in  $\delta$  and  $y$ . By induction we find that  $P_j(y, \delta) = \sum_{k=0}^j a_k^j(\delta) (\delta y)^k$  where  $a_k^j(\delta) = \sum_{s=0}^{m_k^j} b_k^j(s) \delta^s$  is a polynomial such that for every  $j \geq 1$  either  $a_0^j(\delta) = 0$  or else  $b_0^j(0) = 0$  and  $m_k^j \geq 1$ . Of importance here is that all terms in a derivative of  $e^{-a\delta(z+b)^2}$  are of the form  $Az^k \delta^j e^{-a\delta(z+b)^2}$ ,  $A \in \mathbb{C}$ ,  $k \in \mathbb{N}$ ,  $j > 0$  (i.e. there are no terms with  $j = 0$ ). In what follows  $P_0(y, \delta) = 1$  by definition.

Throughtout we will use the following elementary fact: If  $l$  is a nonnegative integer and  $P(z)$  is a polynomial of degree  $\geq 1$  with complex coefficients with leading coefficient having positive real part, then  $\sum_{k=1}^{\infty} k^l e^{-P(k)}$  is absolutely convergent and  $\sum_{k \in \mathbb{Z}} k^l e^{-P(k)}$  is absolutely convergent if  $P$  is of even degree.

**Proof of Lemma 5.5** Let  $\eta_0$  be as in (88) and let  $\eta \in ]0, \eta_0]$  and  $\lambda \in \Lambda$  be arbitrary but fixed in the following. Write  $C_{\text{sd}}(m)$ ,  $f(z; m)$  and  $z_{\text{st}}(m)$  for  $C_{\text{sd}}(m, \lambda)$ ,  $f(z; m, \lambda)$  and  $z_{\text{st}}(m, \lambda)$  respectively,  $m \in \mathbb{Z}$ . By (136) we have

$$\left| \int_{C_{\text{sd}}(m)} f(z; m) dz \right| \leq C e^{-\pi \delta z_{\text{st}}^2(m)} e^{a_0 |z_{\text{st}}(m)|} \left( \exp \left( \frac{a_+^2}{4r|A|} \right) + \exp \left( \frac{a_-^2}{4r|A|} \right) \right)$$

for all  $\delta > 0$ , where  $C$  and  $a_0$  are positive constants independent of  $m$ , and  $a_{\pm} = \pi \delta \sqrt{2} z_{\text{st}}(m) \pm a_0$ . We have

$$e^{-\pi \delta z_{\text{st}}^2(m)} \exp \left( \frac{a_{\pm}^2}{4r|A|} \right) = e^{-\pi \delta z_{\text{st}}^2(m)} \exp \left( \frac{1}{4r|A|} (a_0^2 \pm 2\sqrt{2}\pi \delta a_0 z_{\text{st}}(m)) \right),$$

where  $d = 1 - \frac{\pi\delta}{2r|A|}$ . By this and (82) we get

$$\sum_{m \in \mathbb{Z}} \left| \int_{C_{sd}(m)} f(z; m) dz \right| \leq C' \sum_{\mu = \pm 1} e^{\mu a} \sum_{m \in \mathbb{Z}} e^{\frac{\pi a_0}{|A|} |m|} e^{(b + \mu c)m} \exp \left( -\frac{\pi^3 \delta}{A^2} dm^2 \right),$$

where  $C'$ ,  $a$ ,  $b$  and  $c$  are constants independent of  $m$ . If we choose  $\delta \in ]0, 2r|A|/\pi[$  we have  $d > 0$  and

$$\sum_{m \in \mathbb{Z}} \left| \int_{C_{sd}(m)} f(z; m) dz \right| < \infty.$$

Next let us consider the sum  $\Sigma_2$ . Let  $w = z - l$ . Then  $f(z; m) = \beta(l)\phi(w; l, m)$ , where  $\beta(l)$  and  $\phi$  are given by (97). Since  $A$  and  $B$  are assumed to be real we get

$$|\Sigma_2| \leq \sum_{m \in \mathbb{Z}} \sum_{\substack{l \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi} l > \frac{|A|}{\pi} \eta}} |\text{Res}_{z=i\eta} \{\phi(w; l, m)\}|.$$

Write  $\phi(w; l, m) = \frac{F(w; l, m)}{\sin^k(\pi(w - i\eta))}$ , where  $F(w; l, m) = e^{-\pi\delta(w+l)^2} q(w+l) e^{ir(Aw^2 + 2\pi(m + \frac{B}{2\pi} + \frac{A}{\pi}l)w)}$ . For  $|w - i\eta| < 1$  we have  $\frac{1}{\sin^k(\pi(w - i\eta))} = \sum_{j=-k}^{\infty} c_j (w - i\eta)^j$ , where the  $c_j$ 's are independent of  $l$  and  $m$ . Moreover,  $F(w; l, m) = \sum_{j=0}^{\infty} a_j (w - i\eta)^j$  for  $w \in \mathbb{C}$ . Thus

$$\text{Res}_{w=i\eta} \{\phi(w; l, m)\} = \sum_{j=0}^{k-1} a_j c_{-1-j}.$$

We have  $a_j = \frac{1}{j!} \frac{d^j F(w; l, m)}{dw^j}$ , and thus by Leibnitz' formula and (137)

$$\begin{aligned} a_j &= \frac{1}{j!} \sum_{\substack{k_1, k_2, k_3, k_4 \geq 0 \\ k_1 + k_2 + k_3 + k_4 = j}} \frac{j!}{k_1! k_2! k_3! k_4!} \left. \frac{d^{k_1} e^{-\pi\delta(w+l)^2}}{dw^{k_1}} \right|_{w=i\eta} \\ &\quad \times \left. \frac{d^{k_2} e^{2\pi ir(m + \frac{B}{2\pi} + \frac{A}{\pi}l)w}}{dw^{k_2}} \right|_{w=i\eta} \left. \frac{d^{k_3} q(w+l)}{dw^{k_3}} \right|_{w=i\eta} \left. \frac{d^{k_4} e^{irAw^2}}{dw^{k_4}} \right|_{w=i\eta} \\ &= \sum_{\substack{k_1, k_2, k_3, k_4 \geq 0 \\ k_1 + k_2 + k_3 + k_4 = j}} \frac{M_{k_4}(\eta)}{k_1! k_2! k_3! k_4!} e^{-irA\eta^2} P_{k_1}(i\eta + l, \delta) e^{-\pi\delta(i\eta+l)^2} \\ &\quad \times \left( 2\pi ir \left( m + \frac{B}{2\pi} + \frac{A}{\pi} l \right) \right)^{k_2} e^{-2\pi r(m + \frac{B}{2\pi} + \frac{A}{\pi} l)\eta} \left. \frac{d^{k_3} q(w+l)}{dw^{k_3}} \right|_{w=i\eta}, \end{aligned}$$

where  $P_m(i\eta + l, \delta)$ ,  $m \in \{0, 1, 2, \dots\}$ , is a polynomial in  $l$  for fixed  $\eta > 0$  and  $\delta > 0$  by (137), and  $M_{k_4}(\eta)$  is a constant independent of  $m$  and  $l$ . We therefore get

$$\begin{aligned} |\Sigma_2| &\leq \sum_{j=0}^{k-1} |c_{-1-j}| \sum_{\substack{k_1, k_2, k_3, k_4 \geq 0 \\ k_1 + k_2 + k_3 + k_4 = j}} \frac{|M_{k_4}(\eta)|}{k_1! k_2! k_3! k_4!} \sum_{m \in \mathbb{Z}} \sum_{\substack{l \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi} l > \frac{|A|}{\pi} \eta}} |P_{k_1}(i\eta + l, \delta)| \\ &\quad \times \left| q^{(k_3)}(i\eta + l) \right| e^{-\pi\delta(l^2 - \eta^2)} \left( 2\pi r \left( m + \frac{B}{2\pi} + \frac{A}{\pi} l \right) \right)^{k_2} e^{-2\pi r(m + \frac{B}{2\pi} + \frac{A}{\pi} l)\eta}. \end{aligned}$$

By (80) and (81) we have  $|q^{(k_3)}(i\eta + l)| \leq b_{k_3} e^{a_{k_3}(\eta + m_q)}$ , and for  $l \in \mathbb{Z}$  fixed we have that

$$\sum_{\substack{m \in \mathbb{Z} \\ m + \frac{B}{2\pi} + \frac{A}{\pi}l > \frac{|A|}{\pi}\eta}} \left( 2\pi r \left( m + \frac{B}{2\pi} + \frac{A}{\pi}l \right) \right)^{k_2} e^{-2\pi r(m + \frac{B}{2\pi} + \frac{A}{\pi}l)\eta} = \sum_{\substack{m' \in \mathbb{Z} + \frac{B}{2\pi} + \frac{A}{\pi}l \\ m' > \frac{|A|}{\pi}\eta}} (2\pi r m')^{k_2} e^{-2\pi r m' \eta}.$$

is bounded from above by

$$\beta_{k_2}(\eta) = \max_{x \in [0, \infty[} f(x) + \int_0^\infty f(x) dx = \left( \frac{k_2}{\eta} \right)^{k_2} e^{-k_2} + \frac{k_2!}{2\pi r \eta^{k_2+1}},$$

where  $f(x) = (2\pi r x)^{k_2} e^{-2\pi r x}$ . But then

$$\begin{aligned} |\Sigma_2| &\leq \sum_{j=0}^{k-1} |c_{-1-j}| \sum_{\substack{k_1, k_2, k_3, k_4 \geq 0 \\ k_1 + k_2 + k_3 + k_4 = j}} \frac{|M_{k_4}(\eta)| |\beta_{k_2}(\eta) b_{k_3}| e^{a_{k_3}(\eta + m_q)}}{k_1! k_2! k_3! k_4!} \\ &\quad \times \sum_{l \in \mathbb{Z}} |P_{k_1}(i\eta + l, \delta)| e^{-\pi \delta (l^2 - \eta^2)}, \end{aligned}$$

where the right-hand side is convergent.  $\square$

**Proof of Lemma 5.6** Let  $\lambda \in \Lambda$ ,  $l \in \mathbb{Z}$ ,  $\eta \in ]0, 1[$ , and  $\delta > 0$  be fixed in the following and let  $F(y) = F(y; l, \lambda, \delta, \eta)$  and let  $C(\rho) = C(\lambda, \rho)$  for all  $\rho \in \mathbb{R}$ . To show the first identity in Lemma 5.6, first note that

$$\begin{aligned} \int_{C(\eta)} F^-(y) dy &= \frac{1}{2} \left( \int_{C(\eta)} F^-(y) dy - \int_{C(-\eta)} F^-(y) dy \right) \\ &= \pi i \operatorname{Res}_{y=0} F^-(y) = \pi i \operatorname{Res}_{y=0} F(y), \end{aligned} \quad (138)$$

where we in the last equality use that  $F^+(y)$  has zero residue in  $y = 0$  as an even function. The first equality simply follows by the fact that  $\int_{C(-\rho)} f(-y) dy = \int_{C(\rho)} f(y) dy$  for all  $\rho > 0$  and every complex function such that  $\int_{C(\rho)} f(y) dy$  exists. By this we have

$$\begin{aligned} -\frac{1}{2} \int_{C(-\eta)} F^-(y) dy &= \frac{1}{4} \left( \int_{C(-\eta)} F(-y) dy - \int_{C(-\eta)} F(y) dy \right) \\ &= \frac{1}{4} \left( \int_{C(\eta)} F(y) dy - \int_{C(\eta)} F(-y) dy \right) = \frac{1}{2} \int_{C(\eta)} F^-(y) dy. \end{aligned}$$

The second equality in (138) follows by the residue theorem and the following estimation (since  $y = 0$  is the only pole for  $F^-(y)$  between the contours  $C(-\eta)$  and  $C(\eta)$  since  $\eta \in ]0, 1[$ ): Let  $\tau_R(t) = iR + t$ ,  $t \in [a, b]$ ,  $R \in \mathbb{R}$ , where  $a = \operatorname{sign}(A)R - \eta$  and  $b = \operatorname{sign}(A)R + \eta$  (so  $\tau_R(t)$  is a contour parallel with the real axes and connecting  $C(-\eta)$  and  $C(\eta)$ ). Since

$$\int_{\tau_R} F(-y) dy = \int_{\tau_{-R}} F(y) dy$$

we only have to show that  $|\int_{\tau_R} F(y) dy|$  converges to zero as  $|R|$  converges to infinity. By (132) one finds  $|\sin(\pi \tau_R(t))| \geq \frac{1}{4} e^{\pi |R|}$  for  $|R|$  sufficiently large. By this and (80) we therefore get

$$\left| \int_{\tau_R} F(y) dy \right| \leq b_0 4^k e^{-\pi k |R|} e^{a_0(|R| + l + \eta)} \int_a^b \left| e^{-\pi \delta (iR + t + l + i\eta)^2} \right| e^{-2rA(R + \eta)t} dt$$



for  $|R|$  sufficiently large. Now, for  $|R|$  large,  $a$  and  $b$  have the same sign and

$$\sup_{t \in [a, b]} e^{-2rA(R+\eta)t} \leq e^{-2rA(R+\eta)a} + e^{-2rA(R+\eta)b} \leq 2e^{-2r|A|R^2} e^{2r|A||R|\eta} e^{2r|A|\eta^2}$$

so

$$\begin{aligned} \left| \int_{\tau_R} F(y) dy \right| &\leq 2b_0 4^k e^{-\pi k|R|} e^{a_0(|R|+l+\eta)} e^{-2r|A|R^2} \\ &\quad \times \exp(2r|A||R|\eta) \exp(2r|A|\eta^2) \int_a^b \left| e^{-\pi\delta(iR+t+l+i\eta)^2} \right| dt. \end{aligned}$$

Here  $\int_a^b \left| e^{-\pi\delta(iR+t+l+i\eta)^2} \right| dt \leq e^{\pi\delta(R+\eta)^2} \int_{-\infty}^{\infty} e^{-\pi\delta t^2} dt = \frac{1}{\sqrt{\delta}} e^{\pi\delta(R+\eta)^2}$  so

$$\left| \int_{\tau_R} F(y) dy \right| \leq c e^{d|R|} e^{-(2r|A|-\pi\delta)R^2},$$

where  $c$  and  $d$  are constants independent of  $R$ , so the right-hand side converges to zero as  $|R|$  converges to infinity if  $\delta \in ]0, 2r|A|/\pi[$ .

The last claim about the absolute convergency is proved by the same technique as used in the proof of the absolute convergency of  $\Sigma_2$  in Lemma 5.5. However, the proof is shorter here since we only have a single sum to consider.  $\square$

### 7.3. Appendix C. Proofs of Lemma 5.8 and Proposition 5.9.

**Proof of Lemma 5.8** We will follow the same strategy as in the last part of the proof of Lemma 5.5. In particular we let  $c_j$ ,  $j = -k, -k+1, \dots$  be as in that proof. Let

$$\begin{aligned} \tau(z) &= \cot(\pi r z) \sum_{\nu \in \Gamma(l)} K_l(z; \nu) e^{-2\pi i r V z}, \\ \theta(z) &= \sum_{\nu \in \Gamma(l)} \sum_{\substack{m \in \mathbb{Z} \\ 0 \leq m \leq |V|}} K_l(z; \nu) e^{2\pi i r \text{sign}(V)(m-|V|)z}. \end{aligned}$$

By (111),  $\tau$  is analytic in a neighborhood of  $z = 0$ , in fact on  $D(0, 1/r)$ . For  $\eta \in ]0, 1/r[$  we have

$$\begin{aligned} e^{-\pi\delta(z+l)^2} \tau(z) &= \sum_{j=0}^{\infty} a_j(l, \eta) (z - i\eta)^j, \\ e^{-\pi\delta(z+l)^2} \theta(z) &= \sum_{j=0}^{\infty} b_j(l, \eta) (z - i\eta)^j \end{aligned}$$

for  $z$  in a neighborhood of  $i\eta$ . We thus have

$$\begin{aligned} &\left| \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{\tau(z)}{\sin^k(\pi(z-i\eta))} \right\} \right| \leq \sum_{j=0}^{k-1} |c_{-1-j}| |a_j(l, \eta)| \\ &\leq \sum_{j=0}^{k-1} |c_{-1-j}| \frac{1}{j!} \sum_{n=0}^j \binom{j}{n} |P_n(l + i\eta, \delta)| e^{-\pi\delta(l^2 - \eta^2)} \left| \left( \frac{d}{dz} \right)^{j-n} \tau(z) \right|_{z=i\eta} \end{aligned}$$

and

$$\left| \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{\theta(z)}{\sin^k(\pi(z-i\eta))} \right\} \right| \leq \sum_{j=0}^{k-1} |c_{-1-j}| |b_j(l, \eta)|$$

$$\leq \sum_{j=0}^{k-1} |c_{-1-j}| \frac{1}{j!} \sum_{n=0}^j \binom{j}{n} |P_n(l + i\eta, \delta)| e^{-\pi\delta(l^2 - \eta^2)} \left| \left( \frac{d}{dz} \right)^{j-n} \theta(z) \right|_{z=i\eta}.$$

Now let  $\eta_2 \in ]0, 1/r[$  be fixed and observe that  $A_j = \sup_{l \in \mathbb{Z}, \eta \in [0, \eta_2]} \left| \left( \frac{d}{dz} \right)^j \tau(z) \right|_{z=i\eta}$  and  $B_j = \sup_{l \in \mathbb{Z}, \eta \in [0, \eta_2]} \left| \left( \frac{d}{dz} \right)^j \theta(z) \right|_{z=i\eta}$  are both finite since both  $\tau$  and  $\theta$  are periodic in  $l$  with a period of  $L$  by the assumptions on  $\Gamma(l)$  and  $K_l$ . By the above we get

$$|Z^l(\delta, \eta)| \leq \pi e^{\pi\delta\eta_2^2} \sum_{j=0}^{k-1} |c_{-1-j}| \frac{1}{j!} \sum_{n=0}^j \binom{j}{n} (A_{j-n} + 2B_{j-n}) e^{-\pi\delta l^2} |P_n(l + i\eta, \delta)|.$$

By using the explicit expression for the polynomials  $P_n$ , see (137), we can now appeal to Weierstrass' test for uniform convergence. Note finally that

$$\begin{aligned} \lim_{\eta \rightarrow 0_+} \operatorname{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{\tau(z)}{\sin^k(\pi(z-i\eta))} \right\} &= \sum_{j=0}^{k-1} c_{-1-j} \lim_{\eta \rightarrow 0_+} a_j(l, \eta) \\ &= \sum_{j=0}^{k-1} c_{-1-j} a_j(l, 0) = \operatorname{Res}_{z=0} \left\{ e^{-\pi\delta(z+l)^2} \frac{\tau(z)}{\sin^k(\pi z)} \right\} \end{aligned}$$

since  $a_j(l, \eta) = \frac{1}{j!} \left( \frac{d}{dz} \right)^j e^{-\pi\delta(z+l)^2} \tau(z) \Big|_{z=0}$ , and similarly with the other residue term in  $Z^l(\delta, \eta)$ .  $\square$

For the proof of Proposition 5.9 we need the following small lemma which we state for the sake of easy reference.

**Lemma 7.2.** *Let  $P(z) = Az^2 + Bz + C$  be a polynomial with complex coefficients such that  $\operatorname{Re}(A) > 0$ . Then*

$$\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \sum_{k \in \mathbb{Z}} |k\delta|^l e^{-\delta P(k)} = 0$$

for all  $l \in \{1, 2, \dots\}$  and

$$\lim_{\delta \rightarrow 0_+} \sqrt{\delta} \sum_{k \in \mathbb{Z}} \delta e^{-\delta P(k)} = 0.$$

The first statement follows by comparing the sums by integrals of the form  $\int_0^\infty x^l e^{-\delta a(x+d)^2} dx$ . The second statement is a simple corollary of the periodicity lemma, Lemma 7.1.

**Proof of Proposition 5.9** Let us use the notation from the proof of Lemma 5.8. Thus we have  $Z_{\text{polar}}(r) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \sum_{l \in \mathbb{Z}} Z^l(\delta)$ , where

$$Z^l(\delta) = 2\pi i e^{-\pi\delta l^2} \sum_{j=0}^{k-1} c_{-1-j} \frac{1}{j!} \sum_{n=0}^j \binom{j}{n} P_n(l, \delta) \left( \frac{d}{dz} \right)^{j-n} \left( \frac{i}{2} \tau(z) + \theta(z) \right) \Big|_{z=0}.$$

For  $j \in \{0, 1, \dots, k-1\}$  and  $n \in \{0, 1, \dots, j\}$  we let

$$C(j, n, \delta) = 2\pi i c_{-1-j} \frac{1}{j!} \binom{j}{n} \sum_{l \in \mathbb{Z}} P_n(l, \delta) e^{-\pi\delta l^2} \left( \frac{d}{dz} \right)^{j-n} \left( \frac{i}{2} \tau(z) + \theta(z) \right) \Big|_{z=0}.$$

Then

$$|C(j, n, \delta)| \leq 2\pi |c_{-1-j}| \frac{1}{j!} \binom{j}{n} \left( \frac{1}{2} A_j + B_j \right) \sum_{l \in \mathbb{Z}} |P_n(l, \delta)| e^{-\pi \delta l^2}.$$

By Lemma 7.2 and (137) we therefore get

$$\begin{aligned} Z_{\text{polar}}(r) &= r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \sum_{j=0}^{k-1} \sum_{n=0}^j C(j, n, \delta) = r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \sum_{j=0}^{k-1} C(j, 0, \delta) \\ &= 2\pi i r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \sum_{l \in \mathbb{Z}} e^{-\pi \delta l^2} \sum_{j=0}^{k-1} c_{-1-j} \frac{1}{j!} \left( \frac{d}{dz} \right)^j \left( \frac{i}{2} \tau(z) + \theta(z) \right) \Big|_{z=0} \\ &= \frac{2\pi i r}{L} \sum_{l=0}^{L-1} \sum_{j=0}^{k-1} c_{-1-j} \frac{1}{j!} \left( \frac{d}{dz} \right)^j \left( \frac{i}{2} \tau(z) + \theta(z) \right) \Big|_{z=0}, \end{aligned}$$

where the last identity follows by Lemma 7.1 and the fact that  $\tau$  and  $\theta$  are periodic in  $l$  with a period of  $L$ .  $\square$

#### 7.4. Appendix D. Proofs of Lemma 5.11, Lemma 5.13 and Proposition 5.14.

**Proof of Lemma 5.11** By following the first part of the proof of Lemma 5.4 leading to (136) and by the remarks before the proof of Lemma 5.4 we find that

$$\begin{aligned} \left| \int_{C_{\text{sd}}(m, \lambda)} f(z; m, \lambda) dz \right| &\leq \frac{b_0}{M^k} \sqrt{\frac{\pi}{r|A|}} e^{-\pi \delta z_{\text{st}}^2(m, \lambda)} e^{a_0 |z_{\text{st}}(m, \lambda)|} \\ &\quad \times \left( \exp \left( \frac{a_+^2(m, \lambda)}{4r|A|} \right) + \exp \left( \frac{a_-^2(m, \lambda)}{4r|A|} \right) \right) \end{aligned}$$

for all  $\eta \in [0, \eta_1]$  and all  $(m, \lambda) \in W$ . Here  $a_0$  and  $b_0$  are the constants from (80) and  $a_{\pm}(m, \lambda) = \pi \delta \sqrt{2} z_{\text{st}}(m, \lambda) \pm a_0$ . By following the first part of the proof of Lemma 5.5 we find that

$$\sum_{(m, \lambda) \in W} \frac{|g(\lambda)|}{\sqrt{|A|}} e^{-\pi \delta z_{\text{st}}^2(m, \lambda)} e^{a_0 |z_{\text{st}}(m, \lambda)|} \left( \exp \left( \frac{a_+^2(m, \lambda)}{4r|A|} \right) + \exp \left( \frac{a_-^2(m, \lambda)}{4r|A|} \right) \right)$$

is convergent for  $\delta \in ]0, 2rA_0/\pi[$ , where  $A_0$  is given by (94).  $\square$

**Proof of Lemma 5.13** Keep  $N \in \mathbb{Z}_{\geq 0}$  fixed in the following. For  $\lambda \in \Lambda$ , let  $W_\lambda = \{m \in \mathbb{Z} \mid (m, \lambda) \in W\}$ . We write  $z_{\text{st}}$  for  $z_{\text{st}}(m, \lambda)$ . Let us start by showing that  $Z_{\text{int}, 1}(N; \delta)$  and  $\Sigma^1(N; \delta)$  are absolutely convergent. By Cauchy's integral formula for the derivatives we have

$$a_j(\delta) = \frac{1}{j!} \kappa^{(j)}(z_{\text{st}}; \delta) = \frac{1}{2\pi i} \int_{|z - z_{\text{st}}| = 3\rho} \frac{\kappa(z; \delta)}{(z - z_{\text{st}})^{j+1}} dz.$$

Let  $L(\rho, \delta) = \sup_{|z - z_{\text{st}}| = 3\rho} |\kappa(z; \delta)|$ . Then

$$|a_j(\delta)| \leq (3\rho)^{-j} L(\rho, \delta). \quad (139)$$

By (91) and (121) we get  $L(\rho, \delta) \leq C_\rho \sup_{|z| = 3\rho} \left| e^{-\pi \delta (z + z_{\text{st}})^2} \sin^{-k}(\pi(z + z_{\text{st}})) \right|$ . Since  $z_{\text{st}} \in \mathbb{R}$  we get by (132) that

$$|\sin(\pi(z + z_{\text{st}}))| = \frac{1}{2} \sqrt{e^{2\pi \text{Im}(z)} + e^{-2\pi \text{Im}(z)} - 2 \cos(2\pi(z_{\text{st}} + \text{Re}(z)))}$$

$$\geq \frac{1}{\sqrt{2}} \sqrt{1 - \cos(2\pi(z_{\text{st}} + \text{Re}(z)))}.$$

By our choice of  $\rho$  we have that  $z_{\text{st}}(m, \lambda) + \text{Re}(z) \notin \mathbb{Z} + [-\rho, \rho]$  for all  $(m, \lambda) \in W$  and all  $z \in \mathbb{C}$  with  $|z| = 3\rho$ , so  $|\sin(\pi(z + z_{\text{st}}(m, \lambda)))| \geq M_\rho := \sqrt{1 - \cos(2\pi\rho)} > 0$  for all  $(m, \lambda) \in W$  and all  $z \in \mathbb{C}$  with  $|z| = 3\rho$ . Therefore  $L(\rho, \delta) \leq C_\rho M_\rho^{-k} \sup_{|z|=3\rho} |e^{-\pi\delta(z+z_{\text{st}})^2}|$ . Now use that

$$\begin{aligned} \sup_{|z|=3\rho} |e^{-\pi\delta(z+z_{\text{st}})^2}| &= e^{-\pi\delta z_{\text{st}}^2} \sup_{|z|=3\rho} |e^{-\pi\delta(z^2+2z_{\text{st}}z)}| \\ &\leq e^{-\pi\delta z_{\text{st}}^2} \left( \sup_{|z|=3\rho} e^{-\pi\delta \text{Re}(z^2)} \right) \left( \sup_{|z|=3\rho} e^{-2\pi\delta z_{\text{st}} \text{Re}(z)} \right) \\ &= e^{-\pi\delta z_{\text{st}}^2} e^{9\pi\delta\rho^2} e^{6\pi\delta|z_{\text{st}}|\rho}. \end{aligned}$$

Letting  $C'_\rho = C_\rho M_\rho^{-k} e^{9\pi\delta\rho^2}$  and  $d = 6\pi\rho$  we see that

$$L(\rho, \delta) \leq C'_\rho e^{d\delta|z_{\text{st}}|} e^{-\pi\delta z_{\text{st}}^2}. \quad (140)$$

By (139) and (140) we therefore get

$$\begin{aligned} \sum_{m \in W_\lambda} \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} \left| a_j(\delta) \left( \frac{i}{rA} \right)^{\frac{j+1}{2}} \Gamma\left(\frac{j+1}{2}\right) \right| &\leq C_N^{(0)} \Sigma(\lambda; \delta), \\ \sum_{m \in W_\lambda} \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} \left| a_j(\delta) \left( \frac{i}{rA} \right)^{\frac{j+1}{2}} \Gamma\left(\frac{j+1}{2}, 4|A|\rho^2 r\right) \right| &\leq C_N^{(1)} \Sigma(\lambda; \delta), \end{aligned}$$

where

$$\Sigma(\lambda; \delta) = \sum_{m \in \mathbb{Z}} e^{-\pi\delta z_{\text{st}}(m)^2} \left( e^{d\delta z_{\text{st}}(m)} + e^{-d\delta z_{\text{st}}(m)} \right) \quad (141)$$

with  $z_{\text{st}}(m) = z_{\text{st}}(m, \lambda)$ , and where

$$\begin{aligned} C_N^{(0)} &= C'_\rho \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} (rA_0)^{-\frac{j+1}{2}} (3\rho)^{-j} \Gamma\left(\frac{j+1}{2}\right), \\ C_N^{(1)} &= C_N^{(1)}(r) = C'_\rho \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} (rA_0)^{-\frac{j+1}{2}} (3\rho)^{-j} \Gamma\left(\frac{j+1}{2}, 4A_0\rho^2 r\right) \end{aligned} \quad (142)$$

are independent of  $\lambda$ . Since  $z_{\text{st}}$  is a real affine expression in  $m$  for  $\lambda \in \Lambda$  fixed, the sum  $\Sigma(\lambda; \delta)$  is convergent for all  $\delta > 0$ .

Next we show that  $\Sigma_\nu^2(N; \delta)$  are absolutely convergent. By (122) and (123) we have

$$R_N(z; \delta) = \sum_{j=N+1}^{\infty} a_j(\delta) (z - z_{\text{st}})^j$$

for  $z \in D(z_{\text{st}}, 4\rho)$ . By (139) we have for  $|z - z_{\text{st}}| \leq 2\rho$  that

$$|R_N(z; \delta)| \leq 3L(\rho, \delta) \left( \frac{|z - z_{\text{st}}|}{3\rho} \right)^{N+1}.$$

Therefore

$$|E_\nu(N; \delta)| \leq \int_0^{2\rho} |R_N(\gamma((-1)^\nu t); \delta)| \exp(-r|A|t^2) dt$$

$$\begin{aligned}
&\leq 3L(\rho, \delta)(3\rho)^{-N-1} \int_0^\infty t^{N+1} \exp(-r|A|t^2) dt \\
&= \frac{3}{2} \Gamma\left(\frac{N+2}{2}\right) (3\rho)^{-N-1} (r|A|)^{-(N+2)/2} L(\rho, \delta).
\end{aligned}$$

Note that this estimate is independent of  $\nu \in \{0, 1\}$  and that the only quantity depending on  $m$  is  $L(\rho, \delta)$ . Therefore we have for  $\nu \in \{0, 1\}$  that

$$\sum_{m \in W_\lambda} |E_\nu(N; \delta)| \leq C'_N r^{-(N+2)/2} \sum_{m \in W_\lambda} L(\rho, \delta),$$

where  $C'_N = \frac{1}{2A_0^{(N+2)/2}} (3\rho)^{-N-1} \Gamma\left(\frac{N+2}{2}\right)$ . By (140) we therefore get

$$\sum_{m \in W_\lambda} |E_\nu(N; \delta)| \leq C_N^{(2)} r^{-(N+2)/2} \Sigma(\lambda; \delta) \quad (143)$$

for  $\nu \in \{0, 1\}$ , where  $\Sigma(\lambda; \delta)$  is the convergent sum in (141) and

$$C_N^{(2)} = C'_\rho C'_N = \frac{C_\rho}{2M_\rho^k A_0^{(N+2)/2}} e^{9\pi\delta\rho^2} (3\rho)^{-N-1} \Gamma\left(\frac{N+2}{2}\right). \quad (144)$$

Note that the right-hand side of (143) is independent of  $\nu$  and  $C_N^{(2)}$  is independent of  $\lambda$ .

Let us finally consider the sums  $\Sigma_\nu^3$ . We begin by estimating the integrals

$$J_\nu(\delta) = \int_{2\rho}^\infty \psi((-1)^\nu t; \delta) \exp(-r|A|t^2) dt.$$

By the remarks above the proof of Lemma 5.4 we have

$$|\psi(t; \delta)| \leq \frac{1}{M^k} \left| e^{-\pi\delta\gamma(t)^2} q(\gamma(t)) \right| = \frac{1}{M^k} e^{-\pi\delta z_{\text{st}}^2} e^{-\sqrt{2}\pi\delta z_{\text{st}} t} |q(\gamma(t))|.$$

By our periodicity assumptions on  $q$  and  $z_{\text{st}}$  we have

$$q(z + z_{\text{st}}(m + m_q H, \lambda); \lambda) = q(z + z_{\text{st}}(m, \lambda); \lambda)$$

for all  $(m, \lambda) \in \mathbb{Z} \times \Lambda$  and all  $z \in \mathbb{C}$ . Combining this with (80) we get

$$|q(\gamma((-1)^\nu t))| \leq b_0 e^{a_0 K} e^{a_0 |t|}$$

for all  $(m, \lambda) \in \mathbb{Z} \times \Lambda$  and  $t \in \mathbb{R}$ , where

$$K = \max \{ |z_{\text{st}}(m, \lambda)| \mid m = 0, 1, \dots, m_q H - 1, \lambda \in \Lambda \}.$$

We therefore have the estimate

$$|J_\nu(\delta)| \leq K_1 e^{-\pi\delta z_{\text{st}}^2} \int_{2\rho}^\infty e^{(a_0 - (-1)^\nu \sqrt{2}\pi\delta z_{\text{st}})t} e^{-r|A|t^2} dt,$$

where  $K_1 = \frac{b_0}{M^k} e^{a_0 K}$ . Here  $e^{-(-1)^\nu \sqrt{2}\pi\delta z_{\text{st}} t} = e^{(-1)^\nu \pi \frac{B}{\sqrt{2}A} \delta t} e^{(-1)^\nu \frac{\pi^2}{\sqrt{2}A} \delta m t}$ . Choose  $r_\lambda \in \mathbb{Z}_{\geq 2}$  such that  $e^{a_0 t} \leq e^{\frac{1}{2}r|A|t^2}$  for all  $t \geq 2\rho$  and all  $r \geq r_\lambda$ , and choose  $\delta_\lambda > 0$  such that  $e^{(-1)^\nu \pi \frac{B}{\sqrt{2}A} \delta t} \leq e^{\frac{1}{4}r|A|t^2}$  for all  $t \geq 2\rho$ ,  $r \geq r_\lambda$  and  $\delta \in [0, \delta_\lambda]$ . Then

$$|J_\nu(\delta)| \leq K_1 e^{-\pi\delta z_{\text{st}}^2} \int_{2\rho}^\infty e^{c_\delta |m|t} e^{-at^2} dt$$

for  $r \geq r_\lambda$  and  $\delta \in ]0, \delta_\lambda]$ , where  $c_\delta = \frac{\pi^2}{\sqrt{2}|A|}\delta$  and  $a = \frac{1}{4}r|A|$ . Now use that  $e^{c_\delta|m|t} \leq e^{\frac{a}{2}t^2}$  if and only if  $2c_\delta|m|/a \leq t$ . If  $2\rho \geq 2c_\delta|m|/a$  we therefore get

$$\int_{2\rho}^{\infty} e^{c_\delta|m|t} e^{-at^2} dt \leq \int_{2\rho}^{\infty} e^{-\frac{a}{2}t^2} dt = \sqrt{\frac{2}{|A|}} r^{-1/2} \Gamma\left(\frac{1}{2}, \frac{|A|\rho^2}{2} r\right).$$

For  $2\rho < 2c_\delta|m|/a$  we have

$$\int_{2\rho}^{\infty} e^{c_\delta|m|t} e^{-at^2} dt \leq \int_{2\rho}^{2c_\delta|m|/a} e^{c_\delta|m|t} e^{-at^2} dt + \int_{2c_\delta|m|/a}^{\infty} e^{-\frac{a}{2}t^2} dt.$$

Here

$$\int_{2c_\delta|m|/a}^{\infty} e^{-\frac{a}{2}t^2} dt \leq \int_{2\rho}^{\infty} e^{-\frac{a}{2}t^2} dt = \sqrt{\frac{2}{|A|}} r^{-1/2} \Gamma\left(\frac{1}{2}, \frac{|A|\rho^2}{2} r\right),$$

and

$$\begin{aligned} \int_{2\rho}^{2c_\delta|m|/a} e^{c_\delta|m|t} e^{-at^2} dt &\leq e^{-4a\rho^2} \int_{2\rho}^{2c_\delta|m|/a} e^{c_\delta|m|t} dt \\ &\leq e^{-r|A|\rho^2} \int_0^{2c_\delta|m|/a} e^{c_\delta|m|t} dt \leq \frac{2c_\delta|m|}{a} e^{\frac{2}{a}c_\delta^2 m^2} e^{-r|A|\rho^2}. \end{aligned}$$

We therefore get

$$\sum_{m \in W_\lambda} |J_\nu(\delta)| \leq K_1 \sum_{m \in W_\lambda} e^{-\pi\delta z_{st}^2} \left( \sqrt{\frac{2}{r|A|}} \Gamma\left(\frac{1}{2}, \frac{|A|\rho^2}{2} r\right) + \frac{2c_\delta|m|}{a} e^{\frac{2}{a}c_\delta^2 m^2} e^{-r|A|\rho^2} \right)$$

for  $r \geq r_\lambda$  and  $\delta \in ]0, \delta_\lambda]$ . Here

$$\frac{2c_\delta|m|}{a} e^{\frac{2}{a}c_\delta^2 m^2} e^{-\pi\delta z_{st}^2} = \frac{8\pi^2}{\sqrt{2}A^2} e^{-\pi\delta(\frac{B}{2A})^2} \frac{1}{r} \delta |m| e^{-\pi\delta(\frac{\pi}{A})^2 (1 - \frac{4\pi}{r|A|}\delta) m^2} e^{-\delta(\frac{\pi}{A})^2 Bm}.$$

Finally choose  $\delta_\lambda > 0$  so small that  $\frac{4\pi}{rA}\delta \leq \frac{1}{2}$  for all  $r \in \mathbb{Z}_{\geq 2}$  and all  $\delta \in ]0, \delta_\lambda]$ . Then

$$\sum_{m \in \mathbb{Z}} \frac{2c_\delta|m|}{a} e^{\frac{2}{a}c_\delta^2 m^2} e^{-\pi\delta z_{st}^2} \leq K_2 \frac{1}{r} \sum_{m \in \mathbb{Z}} \delta |m| e^{-\delta P(m; \lambda)},$$

where  $K_2 = \frac{8\pi^2}{\sqrt{2}A_0^2}$  and  $P(m; \lambda) = \frac{\pi^3}{2A^2} m^2 + (\frac{\pi}{A})^2 Bm$ . We therefore get that  $\sum_{m \in W_\lambda} |J_\nu(\delta)| \leq A_3(\lambda; \delta)$  for  $\nu \in \{0, 1\}$ ,  $r \geq r_\lambda$  and  $\delta \in ]0, \delta_\lambda]$ , where

$$A_3(\lambda; \delta) = K_1 \left( \sqrt{\frac{2}{A_0 r}} \Gamma\left(\frac{1}{2}, \frac{A_0 \rho^2}{2} r\right) \sum_{m \in \mathbb{Z}} e^{-\pi\delta z_{st}^2} + \frac{K_2}{r} e^{-rA_0 \rho^2} \sum_{m \in \mathbb{Z}} \delta |m| e^{-\delta P(m; \lambda)} \right) \quad (145)$$

is convergent. Finally put  $r_0 = \max\{r_\lambda \mid \lambda \in \Lambda\}$  and  $\delta_0 = \min\{\delta_\lambda \mid \lambda \in \Lambda\}$ .  $\square$

**Proof of Proposition 5.14** The idea of the proof is to use the periodicity lemma, Lemma 7.1. To this end we have to show that the summand in the infinite sum over  $m$  is periodic in  $m$  (for each fixed  $\lambda \in \Lambda$ ). Keep  $\lambda \in \Lambda$  fixed. First note that

$$irAz_{st}(m)^2 = \pi ir \frac{P}{H} m^2 + Bir \frac{P}{H} m + c,$$

where  $c$  is a constant independent of  $m$ . By (89) it follows that  $\exp(-irAz_{\text{st}}(m)^2)$  is periodic in  $m$ , let's say with a period of  $N_\lambda$ . By (122) and (137) we get

$$a_j(\delta) = \frac{1}{j!} \left. \frac{d^{(j)}\kappa(z; \delta)}{dz^j} \right|_{z=z_{\text{st}}} = \frac{1}{j!} \sum_{n=0}^j \binom{j}{n} P_n(z_{\text{st}}, \delta) e^{-\pi\delta z_{\text{st}}^2} \left. \frac{d^{(j-n)}h(z)}{dz^{j-n}} \right|_{z=z_{\text{st}}},$$

where  $h(z) = h(z; \lambda) = q(z; \lambda)/\sin^k(\pi z)$ . Put

$$C_{j,n,\lambda}(\delta) = \sum_{\substack{m \in \mathbb{Z} \\ z_{\text{st}} \in \mathbb{R} \setminus \mathbb{Z}}} g(\lambda) \exp(-irAz_{\text{st}}^2) P_n(z_{\text{st}}, \delta) e^{-\pi\delta z_{\text{st}}^2} \left. \frac{d^{(j-n)}h(z)}{dz^{j-n}} \right|_{z=z_{\text{st}}}.$$

By (81)  $h$  is periodic in  $z$  with a period of  $2m_q$ , so the derivatives of  $h$  in  $z_{\text{st}}$  are periodic in  $m$  with a period of  $2m_q H$ . We conclude that the functions  $m \mapsto G_j(m, \lambda)$ ,  $j \in \mathbb{Z}_{\geq 0}$ , in (124) are periodic with a period of  $2m_q H N_\lambda$ . Moreover, there exists for each  $\lambda \in \Lambda$  a constant  $K_N(\lambda)$  such that

$$\sup_{m \in \mathbb{Z}} \left| \left. \frac{d^{(n)}h(z; \lambda)}{dz^n} \right|_{z=z_{\text{st}}(\lambda, m)} \right| \leq K_N(\lambda)$$

for all  $n = 0, 1, \dots, N$ . But then  $\lim_{\delta \rightarrow 0+} \sqrt{\delta} |C_{j,n,\lambda}(\delta)| = 0$  since

$$\lim_{\delta \rightarrow 0+} \sqrt{\delta} \sum_{m \in \mathbb{Z}} |P_n(z_{\text{st}}, \delta)| e^{-\pi\delta z_{\text{st}}^2} = 0$$

for all  $j$  and  $n$  with  $1 \leq n \leq j \leq N$  by Lemma 7.2. Note here that

$$\sum_{m \in \mathbb{Z}} |P_n(z_{\text{st}}, \delta)| e^{-\pi\delta z_{\text{st}}^2} \leq \sum_{s=0}^n \left( \frac{\pi}{|A|} \right)^s \sum_{m \in \mathbb{Z}} |a_s^n(\delta)| \delta^s \left| m + \frac{B}{2\pi} \right|^s e^{-\delta P(m)},$$

where the  $a_s^n(\delta)$  are described below (137) and  $P(m) = \pi z_{\text{st}}(m)^2 = \pi \left( \frac{\pi}{A} m + \frac{B}{2A} \right)^2$ . Thus

$$\begin{aligned} r \lim_{\delta \rightarrow 0+} \sqrt{\delta} Z_{\text{int},1}(N; \delta) &= r \sum_{\substack{0 \leq j \leq N \\ j \in 2\mathbb{Z}}} \frac{1}{j!} \Gamma\left(\frac{j+1}{2}\right) \left(\frac{i}{r}\right)^{\frac{j+1}{2}} \sum_{\lambda \in \Lambda} A^{-\frac{j+1}{2}} g(\lambda) \\ &\quad \times \lim_{\delta \rightarrow 0+} \sqrt{\delta} \sum_{\substack{m \in \mathbb{Z} \\ z_{\text{st}} \in \mathbb{R} \setminus \mathbb{Z}}} e^{-\pi\delta z_{\text{st}}^2} G_j(m, \lambda) \end{aligned}$$

by Lemma 5.13. The identity (125) now follows by the periodicity lemma and the fact that  $\Gamma(m + 1/2)/(2m)! = \sqrt{\pi}/(4^m m!)$ .

Let us next show the result about the remainder term  $R_1(N; \delta)$ . We stress again that we do not show that  $\lim_{\delta \rightarrow 0+} \sqrt{\delta} R_1(N; \delta)$  exists, see the discussion below (105). By the proof of Lemma 5.13 we can let

$$\varepsilon_1(N, \lambda; \delta) = \left( C_N^{(1)} + 2C_N^{(2)} r^{-\frac{N+2}{2}} \right) \Sigma(\lambda; \delta) + 2A_3(\lambda; \delta),$$

where  $\Sigma(\lambda; \delta)$  is given by (141),  $A_3(\lambda; \delta)$  is given by (145) and  $C_N^{(1)}$  and  $C_N^{(2)}$  are given by respectively (142) and (144). We note that  $C_N^{(1)}$  and  $C_N^{(2)}$  are continuous in  $\delta$  on  $[0, \delta_0]$ . Moreover, since the polynomial  $z_{\text{st}}(m)$  in  $m$  has  $(\pi/A)^2$  as leading coefficient, we get immediately from Lemma 7.1 that

$$\lim_{\delta \rightarrow 0+} \sqrt{\delta} \Sigma(\lambda; \delta) = \frac{A}{\pi}.$$

By (145), Lemma 7.1 and Lemma 7.2 we get

$$\lim_{\delta \rightarrow 0_+} \sqrt{\delta} A_3(\lambda; \delta) = \frac{K_1 A}{\pi} \sqrt{\frac{2}{A_0 r}} \Gamma\left(\frac{1}{2}, \frac{A_0 \rho^2}{2} r\right).$$

We note that  $C_N^{(2)}$  is independent of  $r$ . For  $\alpha$  and  $u_0$  positive we have the following estimate

$$\Gamma(\alpha, u_0) = \int_{u_0}^{\infty} u^{\alpha-1} e^{-u} du \leq e^{-\frac{u_0}{2}} \int_0^{\infty} u^{\alpha-1} e^{-\frac{u}{2}} du = 2^\alpha \Gamma(\alpha) e^{-\frac{u_0}{2}}$$

finalizing the proof.  $\square$

### 7.5. Appendix E. Proofs of Lemma 5.16 and Proposition 5.17.

**Proof of Lemma 5.16** For the following proof, recall that  $\beta(l) = \beta(l, \lambda)$  has norm 1 for all  $(l, \lambda) \in \mathbb{Z} \times \Lambda$ , see (97). We parametrize  $C_\eta$  by  $C_\eta(t) = e^{\frac{i\pi}{4}\text{sign}(A)t - i\eta}$ ,  $t \in \mathbb{R}$ . Let

$$\tilde{C}_\eta(t) = e^{-\frac{\pi i}{4}\text{sign}(A)} C_\eta(t) = t - \frac{1}{\sqrt{2}} \text{sign}(A) \eta - \frac{i}{\sqrt{2}} \eta,$$

$t \in \mathbb{R}$ , and get

$$I(N; l, \delta, \eta) = \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N d_\nu(l, \delta, \eta) \left( e^{\frac{i\pi}{4}\text{sign}(A)} \right)^{\nu-k+1} \int_{\tilde{C}_\eta} (z^2)^{\frac{\nu-k}{2}} e^{-r|A|z^2} dz.$$

For  $s \in \mathbb{C}$  and  $z \in \mathbb{C}$  we let  $z^s = \exp(s \log(z))$ , where  $\log(re^{i\theta}) = \log(r) + i\theta$ ,  $r > 0$ ,  $\theta \in [0, 2\pi[$ , with  $\log(r) \in \mathbb{R}$  being the usual (principal) logarithm of a positive real number. In particular,  $r^s > 0$  for  $r \in ]0, \infty[$  and  $s \in \mathbb{Z}$ . Let us also in the following text reserve the symbol  $\sqrt{\cdot}$  for the continuous square root on  $\mathbb{C} \setminus ]-\infty, 0]$ , being positive on the positive real axes. Using these conventions we have

$$\int_{\tilde{C}_\eta} (z^2)^{\frac{\nu-k}{2}} e^{-r|A|z^2} dz = -\frac{1}{2} \left( \frac{1}{r|A|} \right)^{\frac{\nu-k}{2}} \frac{1}{\sqrt{r|A|}} \int_{\chi_\eta} z^{\frac{\nu-k}{2}-\frac{1}{2}} e^{-z} dz,$$

where  $\chi_\eta(t) = r|A|\tilde{C}_\eta(t)^2$  is shown on Figure 1.

We have the following integral representation for the gamma function due to Hankel valid for all  $s \in \mathbb{C} \setminus \mathbb{Z}$ :

$$\Gamma(s+1) = \frac{e^{-i\pi s}}{2i \sin(\pi s)} \int_{\infty}^{(0+)} z^s e^{-z} dz.$$

Here the symbol  $\int_{\infty}^{(0+)}$  means that we integrate along a contour which originates at  $+\infty$ , runs in toward the origin just above the real axis, circles the origin once counterclockwise, and then returns to  $+\infty$  just below the real axes, see the contour  $\Gamma$  on Figure 2.

We have

$$\Gamma(s+1) = \frac{e^{-i\pi s}}{2i \sin(\pi s)} \int_{\chi_\eta} z^s e^{-z} dz$$

for all  $s \in \mathbb{R} \setminus \mathbb{Z}$  and all fixed  $\eta > 0$ . To see this, note that the integrand is holomorphic on  $\mathbb{C} \setminus [0, \infty[$ . By Cauchy's theorem we therefore only have to show that  $\int_{\zeta_R^\pm} z^s e^{-z} dz$  converges to zero as  $R$  converges to infinite, where  $\zeta_R^\pm(t) = R + it$ ,  $t \in [0, t_\eta^\pm(R)]$ ,  $R > 0$ , and

$$t_\eta^\pm(R) = \frac{1}{\sqrt{2}} \left( \eta \text{sign}(A) \pm \sqrt{\eta^2 + \frac{2R}{r|A|}} \right)$$






FIGURE 1.




FIGURE 2. Contour of integration for representation of the gamma function.

are the two  $t$ -values for which the line  $R + it$  intersects the contour  $\chi_\eta$ . But this follows by the estimate

$$\left| \int_{\varsigma_R^\pm} z^s e^{-z} dz \right| \leq e^{-R} \left| \int_0^{t_\eta^\pm(R)} |R + it|^s dt \right| \leq |t_\eta^\pm(R)| (R^s + (R + |t_\eta^\pm(R)|)^s) e^{-R}.$$

Therefore

$$I(N; l, \delta, \eta) = \frac{e^{\frac{i\pi}{4}\text{sign}(A)}}{\sqrt{r|A|}} \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N d_\nu(l, \delta, \eta) \left(\frac{i}{rA}\right)^{\frac{\nu-k}{2}} \Gamma\left(\frac{\nu-k+1}{2}\right). \quad (146)$$

To show that  $Z_{\text{int},2}(N; \delta, \eta)$  is absolutely convergent it is enough to show that  $\sum_{l \in \mathbb{Z}} |d_\nu(l, \delta, \eta)|$  is convergent for all  $\nu \in \mathbb{Z}_{\geq 0}$  and all  $\lambda \in \Lambda$  (where  $\lambda$  is suppressed from the notation). By (126) and (137) we have

$$d_\nu(l, \delta, \eta) = \frac{1}{\pi^k} \sum_{j=0}^{\nu} \frac{1}{j!(\nu-j)!} P_j(i\eta + l, \delta) e^{-\pi\delta(i\eta+l)^2} \partial_y^{(\nu-j)} h_\eta(y)|_{y=0}, \quad (147)$$

where  $h_\eta(y) = q(y + i\eta + l) e^{-2rA\eta y} \left(\frac{\pi y}{\sin(\pi y)}\right)^k$ . Let  $\eta_3 > 0$ . By the periodicity assumption on  $q$  (and continuity) we have that

$$M_\nu(\eta_3) := \max \left\{ \left| \partial_y^{(j)} h_\eta(y)|_{y=0} \right| : j \in \{0, 1, \dots, \nu\}, \lambda \in \Lambda, l \in \mathbb{Z}, \eta \in [0, \eta_3] \right\}$$

is finite for  $\nu \in \mathbb{Z}_{\geq 0}$ . Using notation from (137) we therefore get

$$|d_\nu(l, \delta, \eta)| \leq \frac{M_\nu(\eta_3)}{\pi^k} e^{\pi\delta\eta_3^2} \sum_{j=0}^{\nu} \frac{1}{j!(\nu-j)!} \sum_{n=0}^j |a_n^j(\delta)| (\delta(\eta_3 + |l|))^n e^{-\pi\delta l^2}, \quad (148)$$

where the  $a_n^j(\delta)$ 's are polynomials in  $\delta$  independent of  $l$ . It follows that  $\sum_{l \in \mathbb{Z}} |d_\nu(l, \delta, \eta)|$  is convergent. In fact, we see that this series is uniformly convergent w.r.t.  $\eta$  on  $[0, \eta_3]$ .

Let us next show that  $\Sigma^5(N; \delta, \eta)$  is absolutely convergent. To this end we consider the series  $\sum_{l \in \mathbb{Z}} |J_2(N; l, \delta, \eta)|$  for  $\lambda \in \Lambda$  arbitrary but fixed. Here

$$|J_2(N; l, \delta, \eta)| \leq \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N |d_\nu(l, \delta, \eta)| \int_{|t| \geq \frac{1}{2}} |C_\eta(t)|^{\nu-k} \left| e^{irAC_\eta(t)^2} \right| dt.$$

We have  $\left| e^{irAC_\eta(t)^2} \right| = e^{-r|A|t^2} e^{\sqrt{2}r|A|\eta t}$ . Moreover,  $|C_\eta(t)| \leq |t| + \eta$  and  $|C_\eta(t)| \geq |t|/\sqrt{2}$ , so

$$|J_2(N; l, \delta, \eta)| \leq (2\sqrt{2})^k \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N 2^\nu |d_\nu(l, \delta, \eta)| \int_{|t| \geq \frac{1}{2}} |t|^\nu e^{\sqrt{2}r|A|\eta t} e^{-r|A|t^2} dt$$

for  $\eta \leq 1/2$ . For  $t \geq 2\sqrt{2}\eta$  we have  $\exp(\sqrt{2}r|A|\eta t) \leq \exp(\frac{1}{2}r|A|t^2)$ , so

$$|J_2(N; l, \delta, \eta)| \leq (2\sqrt{2})^k \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N 2^\nu |d_\nu(l, \delta, \eta)| \int_{|t| \geq \frac{1}{2}} |t|^\nu e^{-\frac{1}{2}r|A|t^2} dt$$

for  $\eta \in ]0, \eta']$ , where  $\eta' = \frac{1}{4\sqrt{2}}$ . Therefore

$$|J_2(N; l, \delta, \eta)| \leq \frac{1}{4} (2\sqrt{2})^k \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N \left(\frac{8}{r|A|}\right)^{\frac{\nu}{2}+1} |d_\nu(l, \delta, \eta)| \Gamma\left(\frac{\nu+1}{2}, \frac{1}{8}|A|r\right).$$

This implies that  $\sum_{l \in \mathbb{Z}} |J_2(N; l, \delta, \eta)| \leq A_5(N, \lambda; \delta, \eta)$ , where

$$A_5(N, \lambda; \delta, \eta) = \frac{1}{4} (2\sqrt{2})^k \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N \left( \frac{8}{r|A|} \right)^{\frac{\nu}{2}+1} \Gamma \left( \frac{\nu+1}{2}, \frac{1}{8}|A|r \right) \sum_{l \in \mathbb{Z}} |d_\nu(l; \delta, \eta)|. \quad (149)$$

Thus  $\Sigma^5(N; \delta, \eta)$  is absolutely convergent.

Next we show that  $\Sigma^4(N; \delta, \eta)$  is absolutely convergent. It is enough to prove that the series  $\sum_{l \in \mathbb{Z}} |J_1(N; l, \delta, \eta)|$  is convergent. Thus we have to estimate  $R_N(y; l, \delta, \eta) = \sum_{\substack{\nu=N+1 \\ \nu-k \in 2\mathbb{Z}}}^\infty d_\nu(l, \delta, \eta)$  for  $y = C_\eta(t)$ ,  $|t| \leq 1/2$ . By (126) and Cauchy's formula we have

$$d_\nu(l, \delta, \eta) = \frac{1}{\pi^k} \frac{1}{2\pi i} \int_{|z|=\rho} \frac{\Psi(z; l, \delta, \eta)}{z^{\nu+1}} dz$$

for  $\nu \in \mathbb{Z}_{\geq 0}$  and  $\rho \in ]0, 1[$ . Therefore  $|d_\nu(l, \delta, \eta)| \leq \pi^{-k} \rho^{-\nu} L(\rho, l, \delta, \eta)$ , where  $L(\rho, l, \delta, \eta) = \max_{|z|=\rho} |\Psi(z; l, \delta, \eta)|$ . Let  $\nu_1 \in \{N+1, N+2\}$  such that  $\nu_1 - k \in 2\mathbb{Z}$ . Assume first that  $\nu_1 \geq k$ . Then

$$|R_N(y; l, \delta, \eta)| \leq \frac{1}{\pi^k} L(\rho, l, \delta, \eta) \sum_{\substack{\nu=\nu_1 \\ \nu-k \in 2\mathbb{Z}}}^\infty \frac{1}{\rho^\nu} |y|^{\nu-k} \leq \frac{1}{\pi^k} L(\rho, l, \delta, \eta) \frac{1}{\rho^{\nu_1}} |y|^{\nu_1-k} \sum_{\nu=0}^\infty \left( \frac{|y|}{\rho} \right)^\nu$$

for  $\rho \in ]0, 1[$  and  $|y| < \rho$ . Now let  $\rho = 3/4$  and let  $L(l, \delta, \eta) = L(3/4, l, \delta, \eta)$ . Then

$$|R_N(y; l, \delta, \eta)| \leq \frac{9}{\pi^k} \left( \frac{4}{3} \right)^{\nu_1} L(l, \delta, \eta) |y|^{\nu_1-k}$$

for  $|y| \leq 2/3$ . For  $\eta \leq 1/6$  we have  $|C_\eta(t)| \leq 2/3$  for all  $t \in [-1/2, 1/2]$  so

$$|J_1(N; l, \delta, \eta)| \leq \frac{18}{\pi^k} \left( \frac{4}{3} \right)^{\nu_1} \exp \left( \frac{r|A|\eta}{\sqrt{2}} \right) L(l, \delta, \eta) \int_0^{1/2} (t + \eta)^{\nu_1-k} e^{-r|A|t^2} dt.$$

By (127) and the assumptions on  $q$ , see (80) and (81), we have

$$L(\rho, l, \delta, \eta) \leq M_1(\rho)^k b_0 e^{a_0(\rho+\eta+m_q)} e^{2r|A|\eta\rho} \max_{|z|=\rho} \left| e^{-\pi\delta(z+i\eta+l)^2} \right|,$$

where  $M_1(\rho) = \max_{|z|=\rho} |(\pi z)/\sin(\pi z)|$ . Here  $\max_{|z|=\rho} \left| e^{-\pi\delta(z+i\eta+l)^2} \right| \leq e^{-\pi\delta l^2 + 2\pi\delta\rho|l|} e^{\pi\delta(\eta+\rho)^2}$  so

$$L(l, \delta, \eta) = L(3/4, l, \delta, \eta) = C(\delta, \eta) e^{\frac{3}{2}|A|r\eta} \left( e^{-\frac{3}{2}\pi\delta l} + e^{\frac{3}{2}\pi\delta l} \right) e^{-\pi\delta l^2},$$

where  $C(\delta, \eta) = M_1(3/4)^k b_0 e^{a_0(3/4+\eta+m_q)} e^{\pi\delta(\eta+3/4)^2}$ . Therefore  $\sum_{l \in \mathbb{Z}} |J_1(N; l, \delta, \eta)|$  is bounded from above by

$$A_4^0(N, \lambda; \delta, \eta) := C_1(\lambda; \delta, \eta) \int_0^\infty (t + \eta)^{\nu_1-k} e^{-r|A|t^2} dt \sum_{l \in \mathbb{Z}} \left( e^{-\frac{3}{2}\pi\delta l} + e^{\frac{3}{2}\pi\delta l} \right) e^{-\pi\delta l^2}, \quad (150)$$

where

$$C_1(\lambda; \delta, \eta) = \frac{18C(\delta, \eta)}{\pi^k} \left( \frac{4}{3} \right)^{\nu_1} \exp \left( \left( \frac{3}{2} + \frac{1}{\sqrt{2}} \right) |A|r\eta \right).$$

We note that  $A_4^0(N, \lambda; \delta, \eta)$  is convergent, hence  $\Sigma_4(N; \delta, \eta)$  is absolutely convergent.

Assume next that  $\nu_1 < k$  and write

$$R_N(y; l, \delta, \eta) = \sum_{\substack{\nu=\nu_1 \\ \nu-k \in 2\mathbb{Z}}}^{k-2} d_\nu(l, \delta, \eta) y^{\nu-k} + R(y; l, \delta, \eta).$$

Then  $J_1(N; l, \delta, \eta) = I_3(N; l, \delta, \eta) + J_3(l, \delta, \eta)$ , where

$$\begin{aligned} I_3(N; l, \delta, \eta) &= \sum_{\substack{\nu=\nu_1 \\ \nu-k \in 2\mathbb{Z}}}^{k-2} d_\nu(l, \delta, \eta) \int_{C_\eta^0} y^{\nu-k} e^{irAy^2} dy, \\ J_3(l, \delta, \eta) &= \int_{C_\eta^0} R(y; l, \delta, \eta) e^{irAy^2} dy. \end{aligned}$$

Let

$$\Sigma^7(\delta, \eta) = \sum_{\lambda \in \Lambda} g(\lambda) e^{-irA\eta^2} \sum_{l \in I_\lambda} \beta(l) J_3(l, \delta, \eta).$$

According to the above we have that  $\sum_{l \in \mathbb{Z}} |J_3(l, \delta, \eta)|$  is bounded from above by an expression equal to the right-hand side of (150) with  $\nu_1$  replaced by  $k$ , i.e. by

$$A_4^1(\lambda; \delta, \eta) := C_1(\lambda; \delta, \eta) \int_0^\infty e^{-r|A|t^2} dt \sum_{l \in \mathbb{Z}} \left( e^{-\frac{3}{2}\pi\delta l} + e^{\frac{3}{2}\pi\delta l} \right) e^{-\pi\delta l^2}, \quad (151)$$

where  $C_1(\lambda, \delta, \eta)$  is as in (150) with  $\nu_1$  replaced by  $k$ . The series  $A_4^1(\delta, \eta)$  is convergent, hence  $\Sigma^7(\delta, \eta)$  is absolutely convergent. We further partition  $I_3(N; l, \delta, \eta) = I_4(N; l, \delta, \eta) + J_4(N; l, \delta, \eta)$  with

$$\begin{aligned} I_4(N; l, \delta, \eta) &= \sum_{\substack{\nu=\nu_1 \\ \nu-k \in 2\mathbb{Z}}}^{k-2} d_\nu(l, \delta, \eta) \int_{C_\eta} y^{\nu-k} e^{irAy^2} dy, \\ J_4(N; l, \delta, \eta) &= - \sum_{\substack{\nu=\nu_1 \\ \nu-k \in 2\mathbb{Z}}}^{k-2} d_\nu(l, \delta, \eta) \int_{C_\eta^\infty} y^{\nu-k} e^{irAy^2} dy. \end{aligned}$$

Put

$$\begin{aligned} \Sigma^8(N; \delta, \eta) &= \sum_{\lambda \in \Lambda} g(\lambda) e^{-irA\eta^2} \sum_{l \in I_\lambda} \beta(l) I_4(N; l, \delta, \eta), \\ \Sigma^9(N; \delta, \eta) &= \sum_{\lambda \in \Lambda} g(\lambda) e^{-irA\eta^2} \sum_{l \in I_\lambda} \beta(l) J_4(N; l, \delta, \eta). \end{aligned}$$

As in the case  $Z_{\text{int},2}(N; \delta, \eta)$  we get that  $\sum_{l \in \mathbb{Z}} |I_4(N; l, \delta, \eta)|$  is bounded from above by

$$A_4^2(N, \lambda; \delta, \eta) := \sum_{\substack{\nu=\nu_1 \\ \nu-k \in 2\mathbb{Z}}}^{k-2} \left( \frac{1}{r|A|} \right)^{\frac{\nu-k+1}{2}} \left| \Gamma \left( \frac{\nu-k+1}{2} \right) \right| \sum_{l \in \mathbb{Z}} |d_\nu(l, \delta, \eta)|, \quad (152)$$

thus  $\Sigma^8(N; \delta, \eta)$  is absolutely convergent. Moreover, similar to the case  $\Sigma^5(N; \delta, \eta)$  we get that  $\sum_{l \in \mathbb{Z}} |J_4(N; l, \delta, \eta)|$  is bounded from above by

$$A_4^3(N, \lambda; \delta, \eta) := \frac{1}{4} (2\sqrt{2})^k \sum_{\substack{\nu=\nu_1 \\ \nu-k \in 2\mathbb{Z}}}^{k-2} \left( \frac{8}{r|A|} \right)^{\frac{\nu}{2}+1} \Gamma \left( \frac{\nu+1}{2}, \frac{1}{8}|A|r \right) \sum_{l \in \mathbb{Z}} |d_\nu(l; \delta, \eta)| \quad (153)$$

showing that  $\Sigma^9(N; \delta, \eta)$  is absolutely convergent.

Finally let us consider  $\Sigma^6(\delta, \eta)$ . By (132) we have

$$|\sin(\pi C_\eta(t))| \geq \frac{1}{2} \sqrt{e^{2v_\eta(t)} + e^{-2v_\eta(t)} - 2},$$

where  $v_\eta(t) = \pi \left( \frac{1}{\sqrt{2}} \text{sign}(A)t - \eta \right)$ . Let  $\eta_2 < \frac{1}{2\sqrt{2}}$  and let  $M_2 = \inf_{\eta \in [0, \eta_2], |t| \geq \frac{1}{2}} |\sin(\pi C_\eta(t))| > 0$ . Then

$$\sum_{l \in \mathbb{Z}} |I_\infty(l, \delta, \eta)| \leq \frac{1}{M_2^k} \sum_{l \in \mathbb{Z}} \int_{|t| \geq \frac{1}{2}} f(t; l, \delta, \eta) dt$$

for  $\eta \in [0, \eta_2]$  and  $\delta > 0$ , where

$$f(t; l, \delta, \eta) = \left( |\tilde{G}(C_\eta(t); l, \delta, \eta)| + |\tilde{G}(-C_\eta(t); l, \delta, \eta)| \right) \left| e^{irAC_\eta(t)^2} \right|,$$

where  $\tilde{G}(y; l, \delta, \eta) = e^{-\pi\delta(y+i\eta+l)^2} q(y+i\eta+l) e^{-2rA\eta y}$ . By (80), (81) and the fact that  $|C_\eta(t)| \leq |t| + \eta$  we get

$$|q(\pm C_\eta(t) + i\eta + l)| \leq b_0 e^{a_0(2\eta+m_q+|t|)}.$$

Moreover,  $|e^{-2rA\eta C_\eta(t)}| \leq e^{\sqrt{2}r|A|\eta t}$ , and  $|e^{irAC_\eta(t)^2}| \leq e^{-r|A|t^2} e^{\sqrt{2}r|A|\eta|t|}$  so we have

$$f(t; l, \delta, \eta) \leq b_0 e^{a_0(2\eta+m_q)} e^{a_0|t|} e^{2\sqrt{2}r|A|\eta|t|} e^{-r|A|t^2} \sum_{\mu \in \{\pm 1\}} \left| e^{-\pi\delta(\mu C_\eta(t) + i\eta + l)^2} \right|.$$

Here  $|e^{-\pi\delta(-C_\eta(t) + i\eta + l)^2}| = g(t; l) e^{4\pi\delta\eta^2} e^{-2\sqrt{2}\pi\delta\eta \text{sign}(A)t}$  and  $|e^{-\pi\delta(C_\eta(t) + i\eta + l)^2}| = g(t; l)$ , where  $g(t; l) = e^{-\pi\delta l^2} e^{-\sqrt{2}\pi\delta l t}$  so

$$f(t; l, \delta, \eta) \leq C_2(\delta, \eta) e^{-\pi\delta l^2} e^{\sqrt{2}\pi\delta|l||t|} e^{D(\delta, \eta)|t|} e^{2\sqrt{2}r|A|\eta|t|} \exp(-r|A|t^2),$$

where  $C_2(\delta, \eta) = b_0 e^{a_0(2\eta+m_q)} e^{4\pi\delta\eta^2}$  and  $D(\delta, \eta) = a_0 + 2\sqrt{2}\pi\delta\eta$ . Choose for each  $\lambda \in \Lambda$  a  $r_\lambda \in \mathbb{Z}_{\geq 2}$  so

$$e^{(D(\delta, \eta) + 2\sqrt{2}r|A|\eta)t} \leq \exp\left(\frac{1}{2}r|A|t^2\right)$$

for all  $t \geq 1/2$ ,  $\delta, \eta \in [0, 1/2]$  and  $r \geq r_\lambda$ . For  $\delta \in [0, 1/2]$ ,  $\eta \in [0, \eta_2]$ , and  $r \geq r_\lambda$  we then get

$$\int_{\frac{1}{2}}^{\infty} e^{\sqrt{2}\pi\delta|l|t} e^{D(\delta, \eta)t} e^{2\sqrt{2}r|A|\eta t} e^{-r|A|t^2} dt \leq \int_{\frac{1}{2}}^{\infty} e^{\sqrt{2}\pi\delta|l|t} e^{-\frac{1}{2}r|A|t^2} dt.$$

We therefore have

$$\sum_{l \in \mathbb{Z}} |I_\infty(l, \delta, \eta)| \leq \frac{2C_2(\delta, \eta)}{M_2^k} \sum_{l \in \mathbb{Z}} e^{-\pi\delta l^2} \int_{\frac{1}{2}}^{\infty} e^{\sqrt{2}\pi\delta|l|t} e^{-\frac{1}{2}r|A|t^2} dt$$

for  $\delta \in [0, 1/2]$ ,  $\eta \in [0, \eta_2]$  and  $r \geq r_\lambda$ . For  $t > 0$  we have that  $e^{\sqrt{2}\pi\delta|l|t} \leq e^{\frac{r}{4}|A|t^2}$  if and only if  $t \geq c\delta|l|$ , where  $c = 4\sqrt{2}\pi/(r|A|)$ . If  $c\delta|l| \leq \frac{1}{2}$  we therefore get

$$\int_{\frac{1}{2}}^{\infty} e^{\sqrt{2}\pi\delta|l|t} e^{-\frac{1}{2}r|A|t^2} dt \leq \int_{\frac{1}{2}}^{\infty} e^{-\frac{1}{4}r|A|t^2} dt = \frac{1}{\sqrt{r|A|}} \int_{\frac{r|A|}{16}}^{\infty} u^{\frac{1}{2}-1} e^{-u} du = \frac{1}{\sqrt{r|A|}} \Gamma\left(\frac{1}{2}, \frac{1}{16}r|A|\right).$$

For  $\frac{1}{2} < c\delta|l|$  we have

$$\int_{\frac{1}{2}}^{\infty} e^{\sqrt{2}\pi\delta|l|t} e^{-\frac{1}{2}r|A|t^2} dt \leq \int_{\frac{1}{2}}^{c\delta|l|} e^{\sqrt{2}\pi\delta|l|t} e^{-\frac{1}{2}r|A|t^2} dt + \int_{c\delta|l|}^{\infty} e^{-\frac{1}{4}r|A|t^2} dt.$$

Here  $\int_{c\delta|l|}^{\infty} e^{-\frac{1}{4}r|A|t^2} dt \leq \int_{\frac{1}{2}}^{\infty} e^{-\frac{1}{4}r|A|t^2} dt = \frac{1}{\sqrt{r|A|}} \Gamma\left(\frac{1}{2}, \frac{1}{16}r|A|\right)$  and

$$\int_{\frac{1}{2}}^{c\delta|l|} e^{\sqrt{2}\pi\delta|l|t} e^{-\frac{1}{2}r|A|t^2} dt \leq e^{-\frac{1}{16}r|A|} \int_0^{c\delta|l|} e^{\sqrt{2}\pi\delta|l|t} dt \leq e^{-\frac{1}{16}r|A|} c\delta|l| e^{\sqrt{2}\pi\delta c\delta^2 l^2}.$$

By putting everything together we get that  $\sum_{l \in \mathbb{Z}} |I_\infty(l, \delta, \eta)|$  is bounded from above by

$$\frac{2C_2(\delta, \eta)}{M_2^k} \sum_{l \in \mathbb{Z}} e^{-\pi \delta l^2} \left( \frac{1}{\sqrt{r|A|}} \Gamma\left(\frac{1}{2}, \frac{1}{16} r|A|\right) + \frac{4\sqrt{2}\pi e^{-\frac{1}{16}r|A|}}{r|A|} \delta |l| e^{\frac{8\pi\delta}{r|A|}\pi\delta l^2} \right)$$

for  $\delta \in ]0, 1/2]$ ,  $\eta \in ]0, \eta_2]$  and  $r \in \mathbb{Z}_{\geq r_\lambda}$ . By choosing  $\delta_0 \in ]0, 1/2]$  so small that  $\frac{8\pi\delta_0}{A_0} \leq \frac{1}{2}$  we get  $\sum_{l \in \mathbb{Z}} |I_\infty(l, \delta, \eta)| \leq A_6(\lambda; \delta, \eta)$ , where

$$A_6(\lambda; \delta, \eta) = C_3(\delta, \eta) \left( \frac{1}{\sqrt{r|A|}} \Gamma\left(\frac{1}{2}, \frac{1}{16} |A|r\right) \sum_{l \in \mathbb{Z}} e^{-\pi \delta l^2} + C_4(r) \sum_{l \in \mathbb{Z}} \delta |l| e^{-\frac{1}{2}\pi \delta l^2} \right) \quad (154)$$

for all  $\eta \in [0, \eta_2]$ ,  $\delta \in ]0, \delta_0]$  and  $r \in \mathbb{Z}_{\geq r_\lambda}$ . Here  $C_3(\delta, \eta) = 2C_2(\delta, \eta)/M_2^k$  is independent of  $r$  and  $C_4(r) = \frac{4\sqrt{2}\pi}{|A|r} e^{-\frac{1}{16}|A|r}$ . Finally put  $r_0 = \max_{\lambda \in \Lambda} \{r_\lambda\}$ . The claims about uniform convergence immediately follows from the above.  $\square$

**Proof of Proposition 5.17** By the remarks following Lemma 5.16 we have

$$Z_{\text{int},2}(N; \delta) = \frac{1}{\pi^k} \sum_{\lambda \in \Lambda} g(\lambda) \frac{e^{\frac{i\pi}{4} \text{sign}(A)}}{\sqrt{r|A|}} (D(N, \lambda; \delta) + E(N, \lambda; \delta)),$$

where

$$\begin{aligned} D(N, \lambda; \delta) &= \sum_{l \in I_\lambda} \beta(l, \lambda) \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N \left( \frac{i}{rA} \right)^{\frac{\nu-k}{2}} \frac{\Gamma\left(\frac{\nu-k+1}{2}\right)}{\nu!} e^{-\pi \delta l^2} \partial_y^{(\nu)} h(y)|_{y=0}, \\ E(N, \lambda; \delta) &= \sum_{l \in I_\lambda} \beta(l, \lambda) \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N \left( \frac{i}{rA} \right)^{\frac{\nu-k}{2}} \Gamma\left(\frac{\nu-k+1}{2}\right) \\ &\quad \times \sum_{j=1}^{\nu} \frac{1}{j!(\nu-j)!} P_j(l, \delta) e^{-\pi \delta l^2} \partial_y^{(\nu-j)} h(y; l)|_{y=0}, \end{aligned}$$

where  $h(y; l) = h_0(y) = q(y+l) \left( \frac{\pi y}{\sin(\pi y)} \right)^k$ . By Lemma 7.2 together with the remarks about the polynomials  $P_j(l, \delta)$  in connection to (137) we have

$$\lim_{\delta \rightarrow 0_+} \sqrt{\delta} E(N, \lambda; \delta) = 0.$$

By (81) the derivatives of  $y \mapsto h(y; l)$  in  $y = 0$  are periodic in  $l$  with a period of  $m_q$ . Moreover, the function  $\beta(l, \lambda)$  is periodic in  $l$  for each fixed  $\lambda$  with a period equal to the least common multipulum of 2,  $P(\lambda)$  and  $b(\lambda)$ , where  $P(\lambda)$  and  $b(\lambda)$  are as in Proposition 5.17. Finally we observe that if  $l_0 \in I_\lambda$  then  $l_0 + P(\lambda)\mathbb{Z} \subseteq I_\lambda$ . Thus (128) follows by Lemma 7.1.

Next let us estimate the remainder term  $R_2(N; \delta)$ . By the proof of Lemma 5.16 and the remarks after that lemma we have

$$\epsilon(N, \lambda; \delta) \leq A_4(N, \lambda; \delta) + A_5(N, \lambda; \delta) + A_6(\lambda; \delta),$$

where  $A_\nu(N, \lambda; \delta) = \lim_{\eta \rightarrow 0_+} A_\nu(N, \lambda; \delta, \eta) = A_\nu(N, \lambda; \delta, 0)$  for  $\nu = 4, 5$ , and  $A_6(\lambda; \delta)$  is equal to  $\lim_{\eta \rightarrow 0_+} A_6(\lambda; \delta, \eta) = A_6(\lambda; \delta, 0)$ , where  $A_5(N, \lambda; \delta, \eta)$  and  $A_6(\lambda; \delta, \eta)$  are given by respectively (149) and (154), and where  $A_4(N, \lambda; \delta, \eta) = A_4^0(N, \lambda; \delta, \eta)$  in case  $\nu_1 \geq k$  and

$$A_4(N, \lambda; \delta, \eta) = A_4^1(\lambda; \delta, \eta) + A_4^2(N, \lambda; \delta, \eta) + A_4^3(N, \lambda; \delta, \eta) \quad (155)$$

in case  $\nu_1 < k$ , where  $A_4^0(N, \lambda; \delta, \eta)$ ,  $A_4^1(\lambda; \delta, \eta)$ ,  $A_4^2(N, \lambda; \delta, \eta)$  and  $A_4^3(N, \lambda; \delta, \eta)$  are given by respectively (150), (151), (152) and (153). Recall here that  $\nu_1 \in \{N+1, N+2\}$  such that  $\nu_1 - k \in 2\mathbb{Z}$ .

As in the calculation of  $Z_{\text{int},2}(r; N)$  above we get

$$A_5(N, \lambda; \delta) \leq \tilde{A}_5(N, \lambda; \delta) := \frac{1}{4} \left( \frac{2\sqrt{2}}{\pi} \right)^k (D(N, \lambda; \delta) + E(N, \lambda; \delta)),$$

with

$$D(N, \lambda; \delta) = \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N \left( \frac{8}{r|A|} \right)^{\frac{\nu}{2}+1} \frac{1}{\nu!} \Gamma\left(\frac{\nu+1}{2}, \frac{1}{8}|A|r\right) \sum_{l \in \mathbb{Z}} e^{-\pi \delta l^2} \left| \partial_y^{(\nu)} h(y; l) \right|_{y=0}$$

and with  $\lim_{\delta \rightarrow 0_+} \sqrt{\delta} E(N, \lambda; \delta) = 0$ . We let

$$\begin{aligned} A_5(r; N, \lambda) &= r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} \tilde{A}_5(N, \lambda; \delta) = \frac{r}{4} \left( \frac{2\sqrt{2}}{\pi} \right)^k \lim_{\delta \rightarrow 0_+} \sqrt{\delta} D(N, \lambda; \delta) \\ &= \sum_{\substack{\nu=0 \\ \nu-k \in 2\mathbb{Z}}}^N c_\nu r^{-\frac{\nu}{2}} \Gamma\left(\frac{\nu+1}{2}, \frac{1}{8}|A|r\right), \end{aligned}$$

where  $c_\nu = \frac{1}{4m_q \nu!} \left( \frac{2\sqrt{2}}{\pi} \right)^k \left( \frac{8}{|A|} \right)^{\frac{\nu}{2}+1} \sum_{l=0}^{m_q-1} \left| \partial_y^{(\nu)} h(y; l) \right|_{y=0}$  are independent of  $r$ .

Next we get

$$A_6(r; \lambda) := r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} A_6(\lambda; \delta) = C \sqrt{\frac{r}{|A|}} \Gamma\left(\frac{1}{2}, \frac{1}{16}|A|r\right),$$

where  $C = C_3(0, 0)$  is independent of  $r$ .

Let us finally consider  $A_4(N, \lambda; \delta)$ . We first consider the case  $\nu_1 \geq k$ . In this case  $A_4(N, \lambda; \delta)$  is equal to  $A_4^0(N, \lambda; \delta, \eta)$  in (150) with  $\eta = 0$ . We have

$$A_4^0(r; N, \lambda) := r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} A_4(N, \lambda; \delta) = C r^{\frac{k-\nu_1+1}{2}},$$

where  $C = \frac{1}{2} C_1(\lambda; 0, 0) \Gamma\left(\frac{\nu_1-k+1}{2}\right) |A|^{\frac{k-\nu_1-1}{2}}$  is independent of  $r$ . (Actually  $C_1(\lambda; 0, 0)$  is independent of  $\lambda$ .)

Next assume that  $\nu_1 < k$ . Here  $A_4(N, \lambda; \delta)$  is given by the right-hand side of (155) with  $\eta = 0$ . Recall here that  $A_4^1(\lambda; \delta, \eta)$  is simply equal to  $A_4^0(N, \lambda; \delta, \eta)$  with  $\nu_1$  replaced by  $k$ , hence

$$A_4^1(r; \lambda) := r \lim_{\delta \rightarrow 0_+} \sqrt{\delta} A_4^1(\lambda; \delta, 0) = C \sqrt{r},$$

where  $C$  is a constant independent of  $r$ .

The analysis of  $A_4^2(N, \lambda; \delta, 0)$  and  $A_4^3(N, \lambda; \delta, 0)$  are similar to the analysis of  $A_5(N, \lambda; \delta)$ , the main point being to analyse the series  $\sum_{l \in \mathbb{Z}} |d_\nu(l; \delta, 0)|$ . Thus we find that  $A_4^2(N, \lambda; \delta, 0)$  is bounded from above by an expression  $\tilde{A}_4^2(N, \lambda; \delta)$  for which  $A_4^2(r; N, \lambda) := r \lim_{\delta \rightarrow 0_+} \tilde{A}_4^2(N, \lambda; \delta)$  exists and is equal to

$$A_4^2(r; N, \lambda) = \sum_{\substack{\nu=\nu_1 \\ \nu-k \in 2\mathbb{Z}}}^{k-2} c_\nu r^{\frac{k-\nu+1}{2}},$$

where

$$c_\nu = \frac{1}{m_q \pi^k} \left| \Gamma \left( \frac{\nu - k + 1}{2} \right) \right| |A|^{-\frac{\nu - k + 1}{2}} \sum_{l=0}^{m_q-1} \left| \partial_y^{(\nu)} h(y; l) \right|_{y=0}$$

are independent of  $r$ . Finally we find that  $A_4^3(N, \lambda; \delta, 0)$  is bounded from above by an expression  $\tilde{A}_4^3(N, \lambda; \delta)$  for which  $A_4^3(r; N, \lambda) := r \lim_{\delta \rightarrow 0+} \sqrt{\delta} \lim_{\eta \rightarrow 0+} \tilde{A}_4^3(N, \lambda; \delta)$  exists and is equal to

$$A_4^3(r; N, \lambda) = \sum_{\substack{\nu=\nu_1 \\ \nu-k \in 2\mathbb{Z}}}^{k-2} c_\nu r^{-\frac{\nu}{2}} \Gamma \left( \frac{\nu+1}{2}, \frac{1}{8} A_0 r \right),$$

where

$$c_\nu = \frac{1}{4m_q} \left( \frac{2\sqrt{2}}{\pi} \right)^k \left( \frac{8}{|A|} \right)^{\frac{\nu}{2}+1} \sum_{l=0}^{m_q-1} \left| \partial_y^{(\nu)} h(y; l) \right|_{y=0}$$

are independent of  $r$ . Thus we can put

$$\varepsilon_2(N, \lambda; \delta) = \tilde{A}_4(N, \lambda; \delta) + \tilde{A}_5(N, \lambda; \delta) + A_6(\lambda; \delta),$$

where  $\tilde{A}_4(N, \lambda; \delta) = A_4^0(N, \lambda; \delta, 0)$  if  $\nu_1 \geq k$  and  $\tilde{A}_4(N, \lambda; \delta) = A_4^1(\lambda; \delta, 0) + \tilde{A}_4^2(N, \lambda; \delta) + \tilde{A}_4^3(N, \lambda; \delta)$  if  $\nu_1 < k$ . By the above we have

$$r \lim_{\delta \rightarrow 0+} \sqrt{\delta} \varepsilon_2(N, \lambda; \delta) = A_4(r; N, \lambda) + A_5(r; N, \lambda) + A_6(r; \lambda),$$

where  $A_4(r; N, \lambda) = A_4^0(r; N, \lambda)$  for  $\nu_1 \geq k$  and  $A_4(r; N, \lambda) = A_4^1(r; \lambda) + A_4^2(r; N, \lambda) + A_4^3(r; N, \lambda)$  for  $\nu_1 < k$ . By the remarks about the incomplete gamma function at the very end of the proof of Proposition 5.14 we see that this limit is  $O \left( r^{\frac{k-\nu_1+1}{2}} \right)$ .  $\square$

## 7.6. Appendix F. Proof of Lemma 6.1.

**Proof of Lemma 6.1** Let  $(m, \lambda) \in \mathbb{Z} \times \Lambda$  be arbitrary and put  $\nu = \nu(m, \lambda)$ . Then

$$\int_{\kappa_\nu^\rho} f(z; m, \lambda) dz = e^{\pi\delta\rho^2} e^{-2\pi r|\nu|\rho} \int_{-\infty}^{\infty} e^{-\pi\delta t^2} e^{-2\pi\delta i \text{sign}(\nu)\rho t} \frac{q_1(t + i \text{sign}(\nu)\rho) e^{2\pi i r \nu t}}{\sin^k(\pi(t + i \text{sign}(\nu)\rho - i\eta))} dt.$$

But then

$$\left| \int_{\kappa_\nu^\rho} f(z; m, \lambda) dz \right| \leq b'_0 e^{a'_0(m'_q + \rho)} e^{\pi\delta\rho^2} e^{-2\pi r|\nu|\rho} \int_{-\infty}^{\infty} e^{-\pi\delta t^2} \frac{1}{|\sin(\pi(t + i \text{sign}(\nu)\rho - i\eta))|^k} dt.$$

Here

$$|\sin(\pi(t + i \text{sign}(\nu)\rho - i\eta))| = \frac{1}{2} \sqrt{e^{2a} + e^{-2a} - 2 \cos(2\pi t)} \leq \frac{1}{2} \sqrt{e^{2a} + e^{-2a} - 2},$$

where  $a = \pi(\text{sign}(\nu)\rho - \eta)$ . For  $\eta \in ]0, \eta_0]$  we have  $|a| \geq \pi(\rho - \eta_0) > 0$ . If we put  $c_\rho = \sqrt{e^{2\pi(\rho - \eta_0)} + e^{-2\pi(\rho - \eta_0)} - 2}$  we get

$$\begin{aligned} \left| \int_{\kappa_\nu^\rho} f(z; m, \lambda) dz \right| &\leq b'_0 e^{a'_0(m'_q + \rho)} \left( \frac{2}{c_\rho} \right)^k e^{\pi\delta\rho^2} e^{-2\pi r|\nu|\rho} \int_{-\infty}^{\infty} e^{-\pi\delta t^2} dt \\ &= b'_0 e^{a'_0(m'_q + \rho)} \frac{1}{\sqrt{\delta}} \left( \frac{2}{c_\rho} \right)^k e^{\pi\delta\rho^2} e^{-2\pi r|\nu|\rho}. \end{aligned} \quad (156)$$

Therefore

$$\sum_{m \in \mathbb{Z}} \left| \int_{\kappa_\nu^\rho} f(z; m, \lambda) dz \right| \leq \frac{1}{\sqrt{\delta}} b'_0 e^{a'_0(m'_q + \rho)} \left( \frac{2}{c_\rho} \right)^k e^{\pi\delta\rho^2} \sum_{m \in \mathbb{Z}} e^{-2\pi r\rho|\nu(m, \lambda)|},$$



which is convergent since  $\nu(m, \lambda)$  is an affine function of  $m$ , showing that  $Z_{\text{int}}^J(\delta, \eta)$  is absolutely convergent.

For the sum  $Z_{\text{polar}}^J(\delta, \eta)$  we note that

$$\begin{aligned} & \text{Res}_{z=l+i\eta} \left\{ e^{-\pi\delta z^2} \frac{q_1(z)e^{2\pi i r \nu z}}{\sin^k(\pi(z-i\eta))} \right\} \\ &= (-1)^{kl} e^{2\pi i r \nu l} \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l)e^{2\pi i r \nu z}}{\sin^k(\pi(z-i\eta))} \right\}. \end{aligned}$$

We have to show that

$$\Sigma(\lambda) := \sum_{l \in \mathbb{Z}} \sum_{m \in \mathbb{Z}: \text{sign}(\nu(m, \lambda))=1} \left| \text{Res}_{z=i\eta} \left\{ e^{-\pi\delta(z+l)^2} \frac{q_1(z+l)e^{2\pi i r \nu z}}{\sin^k(\pi(z-i\eta))} \right\} \right|$$

is convergent for each  $\lambda \in \Lambda$ . But this follows exactly as in the 2nd part of the proof of Lemma 5.5. In fact we get by using notation from that proof that

$$\Sigma(\lambda) \leq \sum_{j=0}^{k-1} \sum_{l \in \mathbb{Z}} \sum_{m \in \mathbb{Z}: \nu(m, \lambda) \geq 0} |c_{-1-j}| |a_j|,$$

where

$$\begin{aligned} a_j &= \frac{1}{j!} \left. \frac{d^j e^{-\pi\delta(z+l)^2} q_1(z+l)e^{2\pi i r \nu z}}{dz^j} \right|_{z=i\eta} \\ &= \sum_{\substack{k_1, k_2, k_3 \geq 0 \\ k_1 + k_2 + k_3 = j}} \frac{1}{k_1! k_2! k_3!} P_{k_1}(i\eta + l, \delta) e^{-\pi\delta(i\eta+l)^2} (2\pi i r \nu)^{k_2} e^{2\pi i r \nu(i\eta+l)} q_1^{(k_3)}(i\eta + l). \end{aligned}$$

We therefore get

$$|a_j| \leq \sum_{\substack{k_1, k_2, k_3 \geq 0 \\ k_1 + k_2 + k_3 = j}} \frac{1}{k_1! k_2! k_3!} |P_{k_1}(i\eta + l, \delta)| e^{\pi\delta\eta^2} e^{-\pi\delta l^2} (2\pi r \nu)^{k_2} e^{-2\pi r \nu \eta} b'_{k_3} e^{a'_{k_3}(\eta + m'_q)}$$

showing the convergence of  $\Sigma(\lambda)$ . □

## REFERENCES

- [AM] S. Akbulut, J. McCarthy, *Casson's invariant for oriented homology 3-spheres - an exposition*, Princeton Mathematical Notes **36**, Princeton Univ. Press (1990).
- [An1] J. E. Andersen, *The Witten invariant of finite order mapping tori I*, to appear in J. Reine Angew. Math.
- [An2] J. E. Andersen, *The asymptotic expansion conjecture*, Sec. 7.2 in Problems on invariants of knots and 3-manifolds, ed. T. Ohtsuki, in *Invariants of knots and 3-manifolds (Kyoto 2001)*, Geometry and Topology Monographs **4** (2002), 377–572.
- [An3] J. E. Andersen, *Asymptotic faithfulness of the quantum  $\text{SU}(n)$  representations of the mapping class groups*, preprint arXiv:math.QA/0204084, to appear in Ann. of Math.
- [An4] J. E. Andersen, private communication.
- [AH] J. E. Andersen, S. K. Hansen, *Asymptotics of the quantum invariants for surgeries on the figure 8 knot*, to appear in J. Knot Theory Ramifications.
- [AU1] J. E. Andersen, K. Ueno, *Geometric construction of modular functors from conformal field theory*, preprint arXiv:math.DG/0306235
- [AU2] J. E. Andersen, K. Ueno, *Construction of the Reshetikhin–Turaev TQFT using conformal blocks*, in preparation.
- [At] M. F. Atiyah, *The geometry and physics of knots*, Cambridge University Press (1990).

- [Au1] D. R. Auckly, *Topological methods to compute Chern–Simons invariants*, Math. Proc. Cambridge Philos. Soc. **115** (1994), no. 2, 229–251.
- [Au2] D. R. Auckly, *Computing secondary and spectral invariants*, D. Phil. thesis, The University of Michigan (1991).
- [Ax] S. Axelrod, *Overview and warmup example for perturbation theory with instantons*, Geometry and physics (Aarhus, 1995), Lecture Notes in Pure and Appl. Math. **184**, Dekker, New York (1997), 321–338.
- [APW] S. Axelrod, S. Della Pietra, E. Witten, *Geometric quantization of Chern–Simons gauge theory*, J. Differential Geom. **33** (1991), 787–902.
- [AS1] S. Axelrod, I. M. Singer, *Chern–Simons perturbation theory*, Proceedings of the XXth International Conference on Differential Geometric Methods in Theoretical Physics, Vol. **1, 2**, World Sci. Publishing (1992), 3–45.
- [AS2] S. Axelrod, I. M. Singer, *Chern–Simons perturbation theory. II*, J. Differential Geom. **39** (1994), 173–213.
- [BK] B. Bakalov, A. Kirillov Jr., *Lectures on tensor categories and modular functors*, University Lecture Ser. **21**, Amer. Math. Soc. (2001).
- [BH] N. Bleistein, R. H. Handelsman, *Asymptotic expansions of integrals*, Holt, Rinehart and Winston 1974.
- [B] N. G. de Bruijn, *Asymptotic methods in analysis*, North-Holland Publishing Co. (1961).
- [Fa] G. Faltings, *Stable  $G$ -bundles and projective connections*, J. Algebraic Geom. **2** (1993), no. 3, 507–568.
- [FS] R. Fintushel, R. J. Stern, *Instanton homology of Seifert fibred homology three spheres*, Proc. London Math. Soc. **61** (1990), 109–137.
- [F] D. S. Freed, *Classical Chern–Simons theory, 1*, Adv. Math. **113** (1995), 237–303.
- [FG] D. S. Freed, R. E. Gompf, *Computer calculation of Witten’s 3-manifold invariant*, Comm. Math. Phys. **141** (1991), 79–117.
- [Ga1] S. Garoufalidis, *Relations among 3-manifold invariants*, D. Phil. thesis, The University of Chicago (1992).
- [Ga2] S. Garoufalidis, *Applications of quantum invariants in low dimensional topology*, Topology **37** (1998), 219–224.
- [Ha1] S. K. Hansen, *Reshetikhin–Turaev invariants of Seifert 3-manifolds, and their asymptotic expansions*, D. Phil. thesis, The University of Aarhus (1999).
- [Ha2] S. K. Hansen, *Reshetikhin–Turaev invariants of Seifert 3-manifolds and a rational surgery formula*, Algebr. Geom. Topol. **1** (2001), 627–686.
- [HT] S. K. Hansen, T. Takata, *Reshetikhin–Turaev invariants of Seifert 3-manifolds for classical simple Lie algebras*, J. Knot Theory Ramifications **13** (2004), no. 5, 617–668.
- [H1] K. Hikami, *Quantum invariants, modular form, and lattice points*, Int. Math. Res. Not. (2005), no. 3, 121–154.
- [H2] K. Hikami, *On the quantum invariants for the Brieskorn homology spheres*, Internat. J. Math. **16** (2005), no. 6, 661–685.
- [H3] K. Hikami, *On the quantum invariants for the spherical Seifert manifolds*, preprint arXiv:math-ph/0504082
- [Hi] N. Hitchin, *Flat connections and geometric quantization*, Comm. Math. Phys. **131** (1990), no. 2, 347–380.
- [Ho] L. Hörmander, *The analysis of linear partial differential operators. I*, 2nd ed., Grundlehren Math. Wiss. **256**, Springer-Verlag (1990).
- [JN] M. Jankins, W. D. Neumann, *Lectures on Seifert manifolds*, Brandeis Lecture Notes **2** (1981).
- [J1] L. C. Jeffrey, *Chern–Simons–Witten invariants of lens spaces and torus bundles, and the semiclassical approximation*, Comm. Math. Phys. **147** (1992), 563–604.
- [J2] L. C. Jeffrey, *On some aspects of Chern–Simons gauge theory*, D. Phil. thesis, University of Oxford (1991).
- [JL] G. W. Johnson, M. L. Lapidus, *The Feynman integral and Feynman’s operational calculus*, Oxford mathematical monographs, Clarendon Press, Oxford University Press, Oxford (2000).
- [Jo] V. F. R. Jones, *A polynomial invariant for knots via von Neumann algebras*, Bull. Amer. Math. Soc. **12** (1985), no. 1, 103–111.
- [KNTY] N. Kawamoto, Y. Namikawa, A. Tsuchiya, Y. Yamada, *Geometric realization of conformal field theory on Riemann surfaces*, Comm. Math. Phys. **116** (1988).
- [KK1] P. A. Kirk, E. P. Klassen, *Chern–Simons invariants of 3-manifolds and representation spaces of knot groups*, Math. Ann. **287** (1990), 343–367.
- [KK2] P. A. Kirk, E. P. Klassen, *Representation spaces of Seifert fibered homology spheres*, Topology **30** (1991), 77–95.
- [KM] P. B. Kronheimer, T. S. Mrowka, *Dehn surgery, the fundamental group and  $SU(2)$* , preprint math.GT/0312322
- [L] R. Lawrence, *On Ohtsuki’s invariants of homology 3-spheres*, J. Knot Theory Ramifications **8** (1999), 1049–1063.

- [LR] R. Lawrence, L. Rozansky, *Witten–Reshetikhin–Turaev invariants of Seifert manifolds*, Comm. Math. Phys. **205** (1999), no. 2, 287–314.
- [LZ] R. Lawrence, D. Zagier, *Modular forms and quantum invariants of 3-manifolds*, Asian J. Math. **3** (1999), 93–107.
- [La] Y. Laszlo, *Hitchin’s and WZW connections are the same*, J. Diff. Geom. **49** (1998), no. 3, 547–576.
- [Le] T. T. Q. Le, *Quantum invariants of 3-manifolds: Integrality, splitting, and perturbative expansions*, Proceedings of the Pacific Institute for the Mathematical Sciences Workshop “Invariants of Three-Manifolds” (Calgary, AB, 1999). Topology Appl. **127** (2003), 125–152.
- [Les] C. Lescop, *Global surgery formula for the Casson–Walker invariant*, Ann. of Math. Stud. **140** (1996).
- [M] J. M. Montesinos, *Classical tessellations and three-manifolds*, Universitext, Springer-Verlag (1987).
- [Oh1] T. Ohtsuki, *A polynomial invariant of integral homology 3-spheres*, Math. Proc. Cambridge Philos. Soc. **117** (1995), 83–112.
- [Oh2] T. Ohtsuki, *A polynomial invariant of rational homology 3-spheres*, Invent. Math. **123** (1996), 241–257.
- [Oh3] *Problems on invariants of knots and 3-manifolds*, ed. T. Ohtsuki, in *Invariants of knots and 3-manifolds (Kyoto 2001)*, Geometry and Topology Monographs **4** (2002), 377–572.
- [O] F. W. J. Olver, *Asymptotics and special functions*, Academic Press (1974).
- [Or] P. Orlik, *Seifert manifolds*, Lecture Notes Math. **291**, Springer-Verlag (1972).
- [RG] H. Rademacher, E. Grosswald, *Dedekind sums*, The Carus Mathematical Monographs **16**, The Mathematical Association of America (1972).
- [RT] N. Reshetikhin, V. G. Turaev, *Invariants of 3-manifolds via link polynomials and quantum groups*, Invent. Math. **103** (1991), 547–597.
- [Ro1] L. Rozansky, *A large  $k$  asymptotics of Witten’s invariant of Seifert manifolds*, Comm. Math. Phys. **171** (1995), 279–322.
- [Ro2] L. Rozansky, *Witten’s invariant of 3-dimensional manifolds: loop expansion and surgery calculus*, Knots and applications, Ser. Knots Everything **6**, World Sci. Publishing, River Edge, NJ (1995), 271–299.
- [Ro3] L. Rozansky, *Residue formulas for the large  $k$  asymptotics of Witten’s invariants of Seifert manifolds. The case of  $SU(2)$* , Comm. Math. Phys. **178** (1996), 27–60.
- [Ro4] L. Rozansky, *On  $p$ -adic convergence of perturbative invariants of some rational homology spheres*, Duke Math. J. **91** (1998), 353–379.
- [Ro5] L. Rozansky, *On  $p$ -adic properties of the Witten–Reshetikhin–Turaev invariant*, preprint arXiv:math.QA/9806075
- [Sc] P. Scott, *The geometries of 3-manifolds*, Bull. London Math. Soc. **15** (1983), 401–487.
- [Se1] H. Seifert, *Topologie dreidimensionaler gefaserter räume*, Acta. Math. **60** (1933), 147–238.
- [Se2] H. Seifert, *Topology of 3-dimensional fibered spaces* (english translation by W. Heil of: *Topologie dreidimensionaler gefaserter räume*, Acta. Math. **60** (1933), 147–238). In H. Seifert, W. Threlfall, *A textbook of topology*, Pure Appl. Math. **89**, Academic Press (1980), 359–422.
- [T] E. C. Titchmarsh, *The theory of functions*, 2nd ed., Oxford University Press (1952).
- [TUY] A. Tsuchiya, K. Ueno, Y. Yamada, *Conformal field theory on universal family of stable curves with gauge symmetries*, Advanced Studies in Pure Mathematics **19** (1989), 459–566.
- [Tu] V. G. Turaev, *Quantum invariants of knots and 3-manifolds*, de Gruyter Stud. Math. **18**, Walter de Gruyter (1994).
- [U] K. Ueno, *Introduction to conformal field theory with gauge symmetries*, Geometry and Physics (Aarhus 1995), Lecture Notes in Pure and Appl. Math. **184**, Marcel Dekker, Inc. (1997), 603–745.
- [Y1] T. Yoshida, *An abelianization of  $SU(2)$  WZW model*, to appear in Ann. of Math.
- [Y2] T. Yoshida,  *$SU(2)$  WZW model and topological invariants of 3-manifolds*, preprint
- [W] K. Walker, *An extension of Casson’s invariant*, Ann. of Math. Stud. **126** (1992).
- [Wi] E. Witten, *Quantum field theory and the Jones polynomial*, Comm. Math. Phys. **121** (1989), 351–399.
- [Wo] R. Wong, *Asymptotic approximations of integrals*, Academic Press, Inc. (1989).
- [Z] A. Zygmund, *Trigonometric series* Vol. **I**, 2nd ed., Cambridge University Press (1959).

DEPARTMENT OF MATHEMATICS, KANSAS STATE UNIVERSITY, MANHATTAN, KS 66506, USA  
 E-mail address: hansen@math.ksu.edu